

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
**WO 01/36597 A1**

(51) International Patent Classification<sup>7</sup>: C12N 5/04, 5/10,  
15/00, 15/09, 15/63, 15/70, 15/74, 15/82, 15/87, C07H  
21/02, 21/04, A01H 1/00, 9/00, 11/00

(21) International Application Number: PCT/US00/31344

(22) International Filing Date:

14 November 2000 (14.11.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/166,228	17 November 1999 (17.11.1999)	US
60/197,899	17 April 2000 (17.04.2000)	US
60/227,439	22 August 2000 (22.08.2000)	US

SAMAHA, Raymond [US/US]; 2224 Albert Lane,  
Capitola, CA 95010 (US). PILGRIM, Marsha [US/US];  
790 Saltillo Place, Fremont, CA 94536 (US). PINEDA,  
Omaira [CO/US]; 19563 Helen Place, Castro Valley,  
CA 94546 (US). JIANG, Cai-Zhong [CN/US]; 34495  
Heathrow Terrace, Fremont, CA 94555 (US).

(74) Agent: GUERRERO, Karen; Mendel Biotechnology,  
Inc., 21375 Cabot Boulevard, Hayward, CA 94545 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):  
MENDEL BIOTECHNOLOGY, INC. [US/US]; 21375  
Cabot Boulevard, Hayward, CA 94541 (US).

(71) Applicants and

(72) Inventors: CREELMAN, Robert [US/US]; 2801  
Jennifer Drive, Castro Valley, CA 94546 (US). YU,  
Guo-Liang [CN/US]; 242 Gravatt Drive, Berkeley, CA  
94705 (US). ADAM, Luc [CA/US]; 25800 Industrial  
Boulevard, L403, Hayward, CA 94545 (US). RIECH-  
MANN, Jose Luis [ES/US]; 115 Moss Avenue #308,  
Oakland, CA 94611 (US). HEARD, Jacqueline [US/US];  
810 Guildford Avenue, San Mateo, CA 94402 (US).

Published:

— With international search report.

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PLANT BIOCHEMISTRY-RELATED GENES

(57) Abstract: Recombinant polynucleotides and methods for modifying the phenotype of a plant are provided. In particular, the phenotype that is being modified is a plant's biochemical characteristic.



WO 01/36597 A1

## PLANT BIOCHEMISTRY-RELATED GENES

### RELATED APPLICATION INFORMATION

The present invention claims the benefit from US Provisional Patent Application Serial  
5 Nos. 60/166,228 filed November 17, 1999 and 60/197,899 filed April 17, 2000 and "Plant Trait  
Modification III" filed August 22, 2000.

### FIELD OF THE INVENTION

This invention relates to the field of plant biology. More particularly, the present  
invention pertains to compositions and methods for phenotypically modifying a plant.

### 10 BACKGROUND OF THE INVENTION

Transcription factors can modulate gene expression, either increasing or decreasing  
(inducing or repressing) the rate of transcription. This modulation results in differential levels of  
gene expression at various developmental stages, in different tissues and cell types, and in  
response to different exogenous (e.g., environmental) and endogenous stimuli throughout the life  
15 cycle of the organism.

Because transcription factors are key controlling elements of biological pathways,  
altering the expression levels of one or more transcription factors can change entire biological  
pathways in an organism. For example, manipulation of the levels of selected transcription  
factors may result in increased expression of economically useful proteins or metabolic chemicals  
20 in plants or to improve other agriculturally relevant characteristics. Conversely, blocked or  
reduced expression of a transcription factor may reduce biosynthesis of unwanted compounds or  
remove an undesirable trait. Therefore, manipulating transcription factor levels in a plant offers  
tremendous potential in agricultural biotechnology for modifying a plant's traits.

The present invention provides novel transcription factors useful for modifying a plant's  
25 phenotype in desirable ways, such as modifying a plant's biochemical traits.

### SUMMARY OF THE INVENTION

In a first aspect, the invention relates to a recombinant polynucleotide comprising a  
nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a  
polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-22, or a  
30 complementary nucleotide sequence thereof; (b) a nucleotide sequence encoding a polypeptide  
comprising a conservatively substituted variant of a polypeptide of (a); (c) a nucleotide sequence  
comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-22, or a  
complementary nucleotide sequence thereof; (d) a nucleotide sequence comprising silent

substitutions in a nucleotide sequence of (c); (e) a nucleotide sequence which hybridizes under stringent conditions over substantially the entire length of a nucleotide sequence of one or more of: (a), (b), (c), or (d); (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e); (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide having a biological activity that modifies a plant's biochemical characteristic; (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g); (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g); (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-22; (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-22; and (l) a nucleotide sequence which encodes a conserved domain of a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-22. The recombinant polynucleotide may further comprise a constitutive, inducible, or tissue-active promoter operably linked to the nucleotide sequence. The invention also relates to compositions comprising at least two of the above described polynucleotides.

In a second aspect, the invention is an isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide described above.

In another aspect, the invention is a transgenic plant comprising one or more of the above described recombinant polynucleotides. In yet another aspect, the invention is a plant with altered expression levels of a polynucleotide described above or a plant with altered expression or activity levels of an above described polypeptide. Further, the invention is a plant lacking a nucleotide sequence encoding a polypeptide described above. The plant may be a soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, or vegetable brassicas plant.

In a further aspect, the invention relates to a cloning or expression vector comprising the isolated or recombinant polynucleotide described above or cells comprising the cloning or expression vector.

In yet a further aspect, the invention relates to a composition produced by incubating a polynucleotide of the invention with a nuclease, a restriction enzyme, a polymerase; a polymerase and a primer; a cloning vector, or with a cell.

Furthermore, the invention relates to a method for producing a plant having a modified  
5 biochemical trait. The method comprises altering the expression of an isolated or recombinant polynucleotide of the invention or altering the expression or activity of a polypeptide of the invention in a plant to produce a modified plant, and selecting the modified plant for a modified biochemical trait.

In another aspect, the invention relates to a method of identifying a factor that is  
10 modulated by or interacts with a polypeptide encoded by a polynucleotide of the invention. The method comprises expressing a polypeptide encoded by the polynucleotide in a plant; and identifying at least one factor that is modulated by or interacts with the polypeptide. In one embodiment the method for identifying modulating or interacting factors is by detecting binding by the polypeptide to a promoter sequence, or by detecting interactions between an additional  
15 protein and the polypeptide in a yeast two hybrid system, or by detecting expression of a factor by hybridization to a microarray, subtractive hybridization or differential display.

In yet another aspect, the invention is a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest. The method comprises placing the molecule in contact with a plant comprising the polynucleotide or polypeptide  
20 encoded by the polynucleotide of the invention and monitoring one or more of the expression level of the polynucleotide in the plant, the expression level of the polypeptide in the plant, and modulation of an activity of the polypeptide in the plant.

In yet another aspect, the invention relates to an integrated system, computer or computer readable medium comprising one or more character strings corresponding to a polynucleotide of  
25 the invention, or to a polypeptide encoded by the polynucleotide. The integrated system, computer or computer readable medium may comprise a link between one or more sequence strings to a modified plant biochemical trait.

In yet another aspect, the invention is a method for identifying a sequence similar or homologous to one or more polynucleotides of the invention, or one or more polypeptides  
30 encoded by the polynucleotides. The method comprises providing a sequence database; and, querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.

The method may further comprise of linking the one or more of the polynucleotides of the invention, or encoded polypeptides, to a modified plant biochemical phenotype.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a table of exemplary polynucleotide and polypeptide sequences of the invention. The table includes from left to right for each sequence: the SEQ ID No., the internal code reference number (GID), whether the sequence is a polynucleotide or polypeptide sequence, and identification of any conserved domains for the polypeptide sequences.

Figure 2 provides a table of exemplary sequences that are homologous to other sequences provided in the Sequence Listing and that are derived from *Arabidopsis thaliana*. The table includes from left to right: the SEQ ID No., the internal code reference number (GID), identification of the homologous sequence, whether the sequence is a polynucleotide or polypeptide sequence, and identification of any conserved domains for the polypeptide sequences.

Figure 3 provides a table of exemplary sequences that are homologous to the sequences provided in Figures 1 and 2 and that are derived from plants other than *Arabidopsis thaliana*. The table includes from left to right: the SEQ ID No., the internal code reference number (GID), the unique GenBank sequence ID No. (NID), the probability that the comparison was generated by chance (P-value), and the species from which the homologous gene was identified.

#### DETAILED DESCRIPTION

The present invention relates to polynucleotides and polypeptides, e.g. for modifying phenotypes of plants.

In particular, the polynucleotides or polypeptides are useful for modifying traits associated with a plant's biochemical characteristic when the expression levels of the polynucleotides or expression levels or activity levels of the polypeptides are altered.

The polynucleotides of the invention encode plant transcription factors. The plant transcription factors are derived, e.g., from *Arabidopsis thaliana* and can belong, e.g., to one or more of the following transcription factor families: the AP2 (APETALA2) domain transcription factor family (Riechmann and Meyerowitz (1998) J. Biol. Chem. 379:633-646); the MYB transcription factor family (Martin and Paz-Ares (1997) Trends Genet. 13:67-73); the MADS domain transcription factor family (Riechmann and Meyerowitz (1997) J. Biol. Chem. 378:1079-1101); the WRKY protein family (Ishiguro and Nakamura (1994) Mol. Gen. Genet. 244:563-571); the ankyrin-repeat protein family (Zhang et al. (1992) Plant Cell 4:1575-1588); the

miscellaneous protein (MISC) family (Kim et al. (1997) Plant J. 11:1237-1251); the zinc finger protein (Z) family (Klug and Schwabe (1995) FASEB J. 9: 597-604); the homeobox (HB) protein family (Duboule (1994) Guidebook to the Homeobox Genes, Oxford University Press); the CAAT-element binding proteins (Forsburg and Guarente (1989) Genes Dev. 3:1166-1178); the squamosa promoter binding proteins (SPB) (Klein et al. (1996) Mol. Gen. Genet. 1996 250:7-16); the NAM protein family; the IAA/AUX proteins (Rouse et al. (1998) Science 279:1371-1373); the HLH/MYC protein family (Littlewood et al. (1994) Prot. Profile 1:639-709); the DNA-binding protein (DBP) family (Tucker et al. (1994) EMBO J. 13:2994-3002); the bZIP family of transcription factors (Foster et al. (1994) FASEB J. 8:192-200); the BPF-1 protein (Box P-binding factor) family (da Costa e Silva et al. (1993) Plant J. 4:125-135); and the golden protein (GLD) family (Hall et al. (1998) Plant Cell 10:925-936).

In addition to methods for modifying a plant phenotype by employing one or more polynucleotides and polypeptides of the invention described herein, the polynucleotides and polypeptides of the invention have a variety of additional uses. These uses include their use in the recombinant production (i.e., expression) of proteins; as regulators of plant gene expression, as diagnostic probes for the presence of complementary or partially complementary nucleic acids (including for detection of natural coding nucleic acids); as substrates for further reactions, e.g., mutation reactions, PCR reactions, or the like, of as substrates for cloning e.g., including digestion or ligation reactions, and for identifying exogenous or endogenous modulators of the transcription factors.

### DEFINITIONS

A "polynucleotide" is a nucleic acid sequence comprising a plurality of polymerized nucleotide residues, e.g., at least about 15 consecutive polymerized nucleotide residues, optionally at least about 30 consecutive nucleotides, at least about 50 consecutive nucleotides. In many instances, a polynucleotide comprises a nucleotide sequence encoding a polypeptide (or protein) or a domain or fragment thereof. Additionally, the polynucleotide may comprise a promoter, an intron, an enhancer region, a polyadenylation site, a translation initiation site, 5' or 3' untranslated regions, a reporter gene, a selectable marker, or the like. The polynucleotide can be single stranded or double stranded DNA or RNA. The polynucleotide optionally comprises modified bases or a modified backbone. The polynucleotide can be, e.g., genomic DNA or RNA, a transcript (such as an mRNA), a cDNA, a PCR product, a cloned DNA, a synthetic DNA or RNA, or the like. The polynucleotide can comprise a sequence in either sense or antisense orientations.



A "recombinant polynucleotide" is a polynucleotide that is not in its native state, e.g., the polynucleotide comprises a nucleotide sequence not found in nature, or the polynucleotide is in a context other than that in which it is naturally found, e.g., separated from nucleotide sequences with which it typically is in proximity in nature, or adjacent (or contiguous with) nucleotide sequences with which it typically is not in proximity. For example, the sequence at issue can be cloned into a vector, or otherwise recombined with one or more additional nucleic acid.

An "isolated polynucleotide" is a polynucleotide whether naturally occurring or recombinant, that is present outside the cell in which it is typically found in nature, whether purified or not. Optionally, an isolated polynucleotide is subject to one or more enrichment or purification procedures, e.g., cell lysis, extraction, centrifugation, precipitation, or the like.

A "recombinant polypeptide" is a polypeptide produced by translation of a recombinant polynucleotide. An "isolated polypeptide," whether a naturally occurring or a recombinant polypeptide, is more enriched in (or out of) a cell than the polypeptide in its natural state in a wild type cell, e.g., more than about 5% enriched, more than about 10% enriched, or more than about 20%, or more than about 50%, or more, enriched, i.e., alternatively denoted: 105%, 110%, 120%, 150% or more, enriched relative to wild type standardized at 100%. Such an enrichment is not the result of a natural response of a wild type plant. Alternatively, or additionally, the isolated polypeptide is separated from other cellular components with which it is typically associated, e.g., by any of the various protein purification methods herein.

The term "transgenic plant" refers to a plant that contains genetic material, not found in a wild type plant of the same species, variety or cultivar. The genetic material may include a transgene, an insertional mutagenesis event (such as by transposon or T-DNA insertional mutagenesis), an activation tagging sequence, a mutated sequence, a homologous recombination event or a sequence modified by chimeraplasty. Typically, the foreign genetic material has been introduced into the plant by human manipulation.

A transgenic plant may contain an expression vector or cassette. The expression cassette typically comprises a polypeptide-encoding sequence operably linked (i.e., under regulatory control of) to appropriate inducible or constitutive regulatory sequences that allow for the expression of polypeptide. The expression cassette can be introduced into a plant by transformation or by breeding after transformation of a parent plant. A plant refers to a whole plant as well as to a plant part, such as seed, fruit, leaf, or root, plant tissue, plant cells or any other plant material, e.g., a plant explant, as well as to progeny thereof, and to *in vitro* systems that mimic biochemical or cellular components or processes in a cell.

The phrase "ectopically expression or altered expression" in reference to a polynucleotide indicates that the pattern of expression in, e.g., a transgenic plant or plant tissue, is different from the expression pattern in a wild type plant or a reference plant of the same species. For example, the polynucleotide or polypeptide is expressed in a cell or tissue type other than a cell or tissue type in which the sequence is expressed in the wild type plant, or by expression at a time other than at the time the sequence is expressed in the wild type plant, or by a response to different inducible agents, such as hormones or environmental signals, or at different expression levels (either higher or lower) compared with those found in a wild type plant. The term also refers to altered expression patterns that are produced by lowering the levels of expression to below the detection level or completely abolishing expression. The resulting expression pattern can be transient or stable, constitutive or inducible. In reference to a polypeptide, the term "ectopic expression or altered expression" further may relate to altered activity levels resulting from the interactions of the polypeptides with exogenous or endogenous modulators or from interactions with factors or as a result of the chemical modification of the polypeptides.

The term "fragment" or "domain," with respect to a polypeptide, refers to a subsequence of the polypeptide. In some cases, the fragment or domain, is a subsequence of the polypeptide which performs at least one biological function of the intact polypeptide in substantially the same manner, or to a similar extent, as does the intact polypeptide. For example, a polypeptide fragment can comprise a recognizable structural motif or functional domain such as a DNA binding domain that binds to a DNA promoter region, an activation domain or a domain for protein-protein interactions. Fragments can vary in size from as few as 6 amino acids to the full length of the intact polypeptide, but are preferably at least about 30 amino acids in length and more preferably at least about 60 amino acids in length. In reference to a nucleotide sequence, "a fragment" refers to any subsequence of a polynucleotide, typically, of at least consecutive about 15 nucleotides, preferably at least about 30 nucleotides, more preferably at least about 50, of any of the sequences provided herein.

The term "trait" refers to a physiological, morphological, biochemical or physical characteristic of a plant or particular plant material or cell. In some instances, this characteristic is visible to the human eye, such as seed or plant size, or can be measured by available biochemical techniques, such as the protein, starch or oil content of seed or leaves or by the observation of the expression level of genes, e.g., by employing Northern analysis, RT-PCR, microarray gene expression assays or reporter gene expression systems, or by agricultural observations such as stress tolerance, yield or pathogen tolerance.

“Trait modification” refers to a detectable difference in a characteristic in a plant ectopically expressing a polynucleotide or polypeptide of the present invention relative to a plant not doing so, such as a wild type plant. In some cases, the trait modification can be evaluated quantitatively. For example, the trait modification can entail at least about a 2% increase or  
5 decrease in an observed trait (difference), at least a 5% difference, at least about a 10% difference, at least about a 20% difference, at least about a 30%, at least about a 50%, at least about a 70%, or at least about a 100%, or an even greater difference. It is known that there can be a natural variation in the modified trait. Therefore, the trait modification observed entails a change of the normal distribution of the trait in the plants compared with the distribution  
10 observed in wild type plant.

Trait modifications of particular interest include those to seed (such as embryo or endosperm), fruit, root, flower, leaf, stem, shoot, seedling or the like, including: enhanced tolerance to environmental conditions including freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved  
15 tolerance to pest infestations, including nematodes, mollicutes, parasitic higher plants or the like; decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals; improved growth under poor photoconditions (e.g., low light and/or short day length), or changes in expression levels of genes of interest. Other phenotype that can be modified relate to the production of plant metabolites, such as variations in the production of  
20 taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prenyl lipids (such as chlorophylls and carotenoids), glucosinolates, and terpenoids, enhanced or compositionally altered protein or oil production (especially in seeds), or modified sugar (insoluble or soluble) and/or starch composition.

Physical plant characteristics that can be modified include cell development (such as the number  
25 of trichomes), fruit and seed size and number, yields of plant parts such as stems, leaves and roots, the stability of the seeds during storage, characteristics of the seed pod (e.g., susceptibility to shattering), root hair length and quantity, internode distances, or the quality of seed coat. Plant growth characteristics that can be modified include growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time,  
30 flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, as well as plant architecture characteristics such as apical dominance, branching patterns, number of organs, organ identity, organ shape or size.

### POLYPEPTIDES AND POLYNUCLEOTIDES OF THE INVENTION

The present invention provides, among other things, transcription factors (TFs), and transcription factor homologue polypeptides, and isolated or recombinant polynucleotides encoding the polypeptides. These polypeptides and polynucleotides may be employed to modify a plant's biochemical characteristic.

Exemplary polynucleotides encoding the polypeptides of the invention were identified in the *Arabidopsis thaliana* GenBank database using publicly available sequence analysis programs and parameters. Sequences initially identified were then further characterized to identify sequences comprising specified sequence strings corresponding to sequence motifs present in families of known transcription factors. Polynucleotide sequences meeting such criteria were confirmed as transcription factors.

Additional polynucleotides of the invention were identified by screening *Arabidopsis thaliana* and/or other plant cDNA libraries with probes corresponding to known transcription factors under low stringency hybridization conditions. Additional sequences, including full length coding sequences were subsequently recovered by the rapid amplification of cDNA ends (RACE) procedure, using a commercially available kit according to the manufacturer's instructions. Where necessary, multiple rounds of RACE are performed to isolate 5' and 3' ends. The full length cDNA was then recovered by a routine end-to-end polymerase chain reaction (PCR) using primers specific to the isolated 5' and 3' ends. Exemplary sequences are provided in the Sequence Listing.

The polynucleotides of the invention were ectopically expressed in overexpressor or knockout plants and changes in the biochemical characteristics of the plants were observed. Therefore, the polynucleotides and polypeptides can be employed to improve the biochemical characteristics of plants.

#### Making polynucleotides

The polynucleotides of the invention include sequences that encode transcription factors and transcription factor homologue polypeptides and sequences complementary thereto, as well as unique fragments of coding sequence, or sequence complementary thereto. Such polynucleotides can be, e.g., DNA or RNA, e.g., mRNA, cRNA, synthetic RNA, genomic DNA, cDNA synthetic DNA, oligonucleotides, etc. The polynucleotides are either double-stranded or single-stranded, and include either, or both sense (i.e., coding) sequences and antisense (i.e., non-coding, complementary) sequences. The polynucleotides include the coding sequence of a transcription factor, or transcription factor homologue polypeptide, in isolation, in combination with additional coding sequences (e.g., a purification tag, a localization signal, as a fusion-

protein, as a pre-protein, or the like), in combination with non-coding sequences (e.g., introns or inteins, regulatory elements such as promoters, enhancers, terminators, and the like), and/or in a vector or host environment in which the polynucleotide encoding a transcription factor or transcription factor homologue polypeptide is an endogenous or exogenous gene.

- 5           A variety of methods exist for producing the polynucleotides of the invention. Procedures for identifying and isolating DNA clones are well known to those of skill in the art, and are described in, e.g., Berger and Kimmel, Guide to Molecular Cloning Techniques, Methods in Enzymology volume 152 Academic Press, Inc., San Diego, CA ("Berger"); Sambrook et al., Molecular Cloning - A Laboratory Manual (2nd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, 10 Cold Spring Harbor, New York, 1989 ("Sambrook") and Current Protocols in Molecular Biology, F.M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2000) ("Ausubel").

- Alternatively, polynucleotides of the invention, can be produced by a variety of in vitro amplification methods adapted to the present invention by appropriate selection of specific or 15 degenerate primers. Examples of protocols sufficient to direct persons of skill through in vitro amplification methods, including the polymerase chain reaction (PCR) the ligase chain reaction (LCR), Qbeta-replicase amplification and other RNA polymerase mediated techniques (e.g., NASBA), e.g., for the production of the homologous nucleic acids of the invention are found in Berger, Sambrook, and Ausubel, as well as Mullis et al., (1987) PCR Protocols A Guide to 20 Methods and Applications (Innis et al. eds) Academic Press Inc. San Diego, CA (1990) (Innis). Improved methods for cloning in vitro amplified nucleic acids are described in Wallace et al., U.S. Pat. No. 5,426,039. Improved methods for amplifying large nucleic acids by PCR are summarized in Cheng et al. (1994) Nature 369: 684-685 and the references cited therein, in which PCR amplicons of up to 40kb are generated. One of skill will appreciate that essentially any 25 RNA can be converted into a double stranded DNA suitable for restriction digestion, PCR expansion and sequencing using reverse transcriptase and a polymerase. See, e.g., Ausubel, Sambrook and Berger, *all supra*.

- Alternatively, polynucleotides and oligonucleotides of the invention can be assembled from fragments produced by solid-phase synthesis methods. Typically, fragments of up to 30 approximately 100 bases are individually synthesized and then enzymatically or chemically ligated to produce a desired sequence, e.g., a polynucleotide encoding all or part of a transcription factor. For example, chemical synthesis using the phosphoramidite method is described, e.g., by Beaucage et al. (1981) Tetrahedron Letters 22:1859-69; and Matthes et al. (1984) EMBO J. 3:801-5. According to such methods, oligonucleotides are synthesized, purified,

annealed to their complementary strand, ligated and then optionally cloned into suitable vectors. And if so desired, the polynucleotides and polypeptides of the invention can be custom ordered from any of a number of commercial suppliers.

#### HOMOLOGOUS SEQUENCES

5 Sequences homologous, i.e., that share significant sequence identity or similarity, to those provided in the Sequence Listing, derived from *Arabidopsis thaliana* or from other plants of choice are also an aspect of the invention. Homologous sequences can be derived from any plant including monocots and dicots and in particular agriculturally important plant species, including but not limited to, crops such as soybean, wheat, corn, potato, cotton, rice, oilseed rape (including  
10 canola), sunflower, alfalfa, sugarcane and turf; or fruits and vegetables, such as banana, blackberry, blueberry, strawberry, and raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits (such as apple, peach, pear, cherry and plum) and vegetable brassicas (such as broccoli, cabbage,  
15 cauliflower, brussel sprouts and kohlrabi). Other crops, fruits and vegetables whose phenotype can be changed include barley, rye, millet, sorghum, currant, avocado, citrus fruits such as oranges, lemons, grapefruit and tangerines, artichoke, cherries, nuts such as the walnut and peanut, endive, leek, roots, such as arrowroot, beet, cassava, turnip, radish, yam, and sweet potato, and beans. The homologous sequences may also be derived from woody species, such  
20 pine, poplar and eucalyptus.

Transcription factors that are homologous to the listed sequences will typically share at least about 30% amino acid sequence identity. More closely related transcription factors can share at least about 50%, about 60%, about 65%, about 70%, about 75% or about 80% or about 90% or about 95% or about 98% or more sequence identity with the listed sequences. Factors  
25 that are most closely related to the listed sequences share, e.g., at least about 85%, about 90% or about 95% or more % sequence identity to the listed sequences. At the nucleotide level, the sequences will typically share at least about 40% nucleotide sequence identity, preferably at least about 50%, about 60%, about 70% or about 80% sequence identity, and more preferably about 85%, about 90%, about 95% or about 97% or more sequence identity to one or more of the listed  
30 sequences. The degeneracy of the genetic code enables major variations in the nucleotide sequence of a polynucleotide while maintaining the amino acid sequence of the encoded protein. Conserved domains within a transcription factor family may exhibit a higher degree of sequence

homology, such as at least 65% sequence identity including conservative substitutions, and preferably at least 80% sequence identity.

Identifying Nucleic Acids by Hybridization

Polynucleotides homologous to the sequences illustrated in the Sequence Listing can be identified, e.g., by hybridization to each other under stringent or under highly stringent conditions. Single stranded polynucleotides hybridize when they associate based on a variety of well characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. The stringency of a hybridization reflects the degree of sequence identity of the nucleic acids involved, such that the higher the stringency, the more similar are the two polynucleotide strands. Stringency is influenced by a variety of factors, including temperature, salt concentration and composition, organic and non-organic additives, solvents, etc. present in both the hybridization and wash solutions and incubations (and number), as described in more detail in the references cited above.

An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Nucleic acid molecules that hybridize under stringent conditions will typically hybridize to a probe based on either the entire cDNA or selected portions, e.g., to a unique subsequence, of the cDNA under wash conditions of 0.2x SSC to 2.0 x SSC, 0.1% SDS at 50-65° C, for example 0.2 x SSC, 0.1% SDS at 65° C. For identification of less closely related homologues washes can be performed at a lower temperature, e.g., 50° C. In general, stringency is increased by raising the wash temperature and/or decreasing the concentration of SSC.

As another example, stringent conditions can be selected such that an oligonucleotide that is perfectly complementary to the coding oligonucleotide hybridizes to the coding oligonucleotide with at least about a 5-10x higher signal to noise ratio than the ratio for hybridization of the perfectly complementary oligonucleotide to a nucleic acid encoding a transcription factor known as of the filing date of the application. Conditions can be selected such that a higher signal to noise ratio is observed in the particular assay which is used, e.g., about 15x, 25x, 35x, 50x or more. Accordingly, the subject nucleic acid hybridizes to the unique coding oligonucleotide with at least a 2x higher signal to noise ratio as compared to hybridization of the coding oligonucleotide to a nucleic acid encoding known polypeptide. Again, higher signal to noise ratios can be selected, e.g., about 5x, 10x, 25x, 35x, 50x or more. The particular signal will

depend on the label used in the relevant assay, e.g., a fluorescent label, a colorimetric label, a radioactive label, or the like.

- Alternatively, transcription factor homologue polypeptides can be obtained by screening an expression library using antibodies specific for one or more transcription factors. With the provision herein of the disclosed transcription factor, and transcription factor homologue nucleic acid sequences, the encoded polypeptide(s) can be expressed and purified in a heterologous expression system (e.g., *E. coli*) and used to raise antibodies (monoclonal or polyclonal) specific for the polypeptide(s) in question. Antibodies can also be raised against synthetic peptides derived from transcription factor, or transcription factor homologue, amino acid sequences.
- 10 Methods of raising antibodies are well known in the art and are described in Harlow and Lane (1988) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York. Such antibodies can then be used to screen an expression library produced from the plant from which it is desired to clone additional transcription factor homologues, using the methods described above. The selected cDNAs can be confirmed by sequencing and enzymatic activity.

#### 15 SEQUENCE VARIATIONS

- It will readily be appreciated by those of skill in the art, that any of a variety of polynucleotide sequences are capable of encoding the transcription factors and transcription factor homologue polypeptides of the invention. Due to the degeneracy of the genetic code, many different polynucleotides can encode identical and/or substantially similar polypeptides in addition to those sequences illustrated in the Sequence Listing.
- 20

For example, Table 1 illustrates, e.g., that the codons AGC, AGT, TCA, TCC, TCG, and TCT all encode the same amino acid: serine. Accordingly, at each position in the sequence where there is a codon encoding serine, any of the above trinucleotide sequences can be used without altering the encoded polypeptide.



**Table 1**

Amino acids			Codon							
Alanine	Ala	A	GCA	GCC	GCG	GCU				
Cysteine	Cys	C	TGC	TGT						
Aspartic acid	Asp	D	GAC	GAT						
Glutamic acid	Glu	E	GAA	GAG						
Phenylalanine	Phe	F	TTC	TTT						
Glycine	Gly	G	GGA	GGC	GGG	GGT				
Histidine	His	H	CAC	CAT						
Isoleucine	Ile	I	ATA	ATC	ATT					
Lysine	Lys	K	AAA	AAG						
Leucine	Leu	L	TTA	TTG	CTA	CTC	CTG	CTT		
Methionine	Met	M	ATG							
Asparagine	Asn	N	AAC	AAT						
Proline	Pro	P	CCA	CCC	CCG	CCT				
Glutamine	Gln	Q	CAA	CAG						
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGT		
Serine	Ser	S	AGC	AGT	TCA	TCC	TCG	TCT		
Threonine	Thr	T	ACA	ACC	ACG	ACT				
Valine	Val	V	GTA	GTC	GTG	GTT				
Tryptophan	Trp	W	TGG							
Tyrosine	Tyr	Y	TAC	TAT						

Sequence alterations that do not change the amino acid sequence encoded by the polynucleotide are termed "silent" variations. With the exception of the codons ATG and TGG, encoding methionine and tryptophan, respectively, any of the possible codons for the same amino acid can be substituted by a variety of techniques, e.g., site-directed mutagenesis, available in the art. Accordingly, any and all such variations of a sequence selected from the above table are a feature of the invention.

In addition to silent variations, other conservative variations that alter one, or a few amino acids in the encoded polypeptide, can be made without altering the function of the polypeptide, these conservative variants are, likewise, a feature of the invention.

For example, substitutions, deletions and insertions introduced into the sequences provided in the Sequence Listing are also envisioned by the invention. Such sequence modifications can be engineered into a sequence by site-directed mutagenesis (Wu (ed.) Meth. Enzymol. (1993) vol. 217, Academic Press) or the other methods noted below. Amino acid substitutions are typically of single residues; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. In preferred embodiments, deletions or insertions are made in adjacent pairs, e.g., a deletion of two residues or insertion of two residues. Substitutions, deletions, insertions or any combination thereof can be

combined to arrive at a sequence. The mutations that are made in the polynucleotide encoding the transcription factor should not place the sequence out of reading frame and should not create complementary regions that could produce secondary mRNA structure. Preferably, the polypeptide encoded by the DNA performs the desired function.

- 5       Conservative substitutions are those in which at least one residue in the amino acid sequence has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the Table 2 when it is desired to maintain the activity of the protein. Table 2 shows amino acids which can be substituted for an amino acid in a protein and which are typically regarded as conservative substitutions.

10

**Table 2**

<b>Residue</b>	<b>Conservative Substitutions</b>
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Gln	Asn
Cys	Ser
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; Gln
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr; Gly
Thr	Ser; Val
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

Substitutions that are less conservative than those in Table 2 can be selected by picking residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in protein properties will be those in which (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

#### FURTHER MODIFYING SEQUENCES OF THE INVENTION—MUTATION/ FORCED EVOLUTION

In addition to generating silent or conservative substitutions as noted, above, the present invention optionally includes methods of modifying the sequences of the Sequence Listing. In the methods, nucleic acid or protein modification methods are used to alter the given sequences to produce new sequences and/or to chemically or enzymatically modify given sequences to change the properties of the nucleic acids or proteins.

Thus, in one embodiment, given nucleic acid sequences are modified, e.g., according to standard mutagenesis or artificial evolution methods to produce modified sequences. For example, Ausubel, *supra*, provides additional details on mutagenesis methods. Artificial forced evolution methods are described, e.g., by Stemmer (1994) *Nature* 370:389-391, and Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751. Many other mutation and evolution methods are also available and expected to be within the skill of the practitioner.

Similarly, chemical or enzymatic alteration of expressed nucleic acids and polypeptides can be performed by standard methods. For example, sequence can be modified by addition of lipids, sugars, peptides, organic or inorganic compounds, by the inclusion of modified nucleotides or amino acids, or the like. For example, protein modification techniques are illustrated in Ausubel, *supra*. Further details on chemical and enzymatic modifications can be found herein. These modification methods can be used to modify any given sequence, or to modify any sequence produced by the various mutation and artificial evolution modification methods noted herein.

Accordingly, the invention provides for modification of any given nucleic acid by mutation, evolution, chemical or enzymatic modification, or other available methods, as well as

for the products produced by practicing such methods, e.g., using the sequences herein as a starting substrate for the various modification approaches.

For example, optimized coding sequence containing codons preferred by a particular prokaryotic or eukaryotic host can be used e.g., to increase the rate of translation or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, as compared with transcripts produced using a non-optimized sequence. Translation stop codons can also be modified to reflect host preference. For example, preferred stop codons for *S. cerevisiae* and mammals are TAA and TGA, respectively. The preferred stop codon for monocotyledonous plants is TGA, whereas insects and *E. coli* prefer to use TAA as the stop codon.

The polynucleotide sequences of the present invention can also be engineered in order to alter a coding sequence for a variety of reasons, including but not limited to, alterations which modify the sequence to facilitate cloning, processing and/or expression of the gene product. For example, alterations are optionally introduced using techniques which are well known in the art, e.g., site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to introduce splice sites, etc.

Furthermore, a fragment or domain derived from any of the polypeptides of the invention can be combined with domains derived from other transcription factors or synthetic domains to modify the biological activity of a transcription factor. For instance, a DNA binding domain derived from a transcription factor of the invention can be combined with the activation domain of another transcription factor or with a synthetic activation domain. A transcription activation domain assists in initiating transcription from a DNA binding site. Examples include the transcription activation region of VP16 or GAL4 (Moore et al. (1998) Proc. Natl. Acad. Sci. USA 95: 376-381; and Aoyama et al. (1995) Plant Cell 7:1773-1785), peptides derived from bacterial sequences (Ma and Ptashne (1987) Cell 51: 113-119) and synthetic peptides (Giniger and Ptashne, (1987) Nature 330:670-672).

#### EXPRESSION AND MODIFICATION OF POLYPEPTIDES

Typically, polynucleotide sequences of the invention are incorporated into recombinant DNA (or RNA) molecules that direct expression of polypeptides of the invention in appropriate host cells, transgenic plants, in vitro translation systems, or the like. Due to the inherent degeneracy of the genetic code, nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence can be substituted for any listed sequence to provide for cloning and expressing the relevant homologue.

### Vectors, Promoters and Expression Systems

The present invention includes recombinant constructs comprising one or more of the nucleic acid sequences herein. The constructs typically comprise a vector, such as a plasmid, a cosmid, a phage, a virus (e.g., a plant virus), a bacterial artificial chromosome (BAC), a yeast  
5 artificial chromosome (YAC), or the like, into which a nucleic acid sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

10 General texts which describe molecular biological techniques useful herein, including the use and production of vectors, promoters and many other relevant topics, include Berger, Sambrook and Ausubel, *supra*. Any of the identified sequences can be incorporated into a cassette or vector, e.g., for expression in plants. A number of expression vectors suitable for stable transformation of plant cells or for the establishment of transgenic plants have been described  
15 including those described in Weissbach and Weissbach, (1989) Methods for Plant Molecular Biology, Academic Press, and Gelvin et al., (1990) Plant Molecular Biology Manual, Kluwer Academic Publishers. Specific examples include those derived from a Ti plasmid of *Agrobacterium tumefaciens*, as well as those disclosed by Herrera-Estrella et al. (1983) Nature 303: 209, Bevan (1984) Nucl Acid Res. 12: 8711-8721, Klee (1985) Bio/Technology 3: 637-642,  
20 for dicotyledonous plants.

Alternatively, non-Ti vectors can be used to transfer the DNA into monocotyledonous plants and cells by using free DNA delivery techniques. Such methods can involve, for example, the use of liposomes, electroporation, microprojectile bombardment, silicon carbide whiskers, and viruses. By using these methods transgenic plants such as wheat, rice (Christou (1991)  
25 Bio/Technology 9: 957-962) and corn (Gordon-Kamm (1990) Plant Cell 2: 603-618) can be produced. An immature embryo can also be a good target tissue for monocots for direct DNA delivery techniques by using the particle gun (Weeks et al. (1993) Plant Physiol 102: 1077-1084; Vasil (1993) Bio/Technology 10: 667-674; Wan and Lemeaux (1994) Plant Physiol 104: 37-48, and for *Agrobacterium*-mediated DNA transfer (Ishida et al. (1996) Nature Biotech 14: 745-750).

30 Typically, plant transformation vectors include one or more cloned plant coding sequence (genomic or cDNA) under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant transformation vectors typically also contain a promoter (e.g., a regulatory region controlling inducible or constitutive, environmentally-or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start

site, an RNA processing signal (such as intron splice sites), a transcription termination site, and/or a polyadenylation signal.

Examples of constitutive plant promoters which can be useful for expressing the TF sequence include: the cauliflower mosaic virus (CaMV) 35S promoter, which confers  
 5 constitutive, high-level expression in most plant tissues (*see, e.g.,* Odel et al. (1985) Nature 313:810); the nopaline synthase promoter (An et al. (1988) Plant Physiol 88:547); and the octopine synthase promoter (Fromm et al. (1989) Plant Cell 1: 977).

A variety of plant gene promoters that regulate gene expression in response to environmental, hormonal, chemical, developmental signals, and in a tissue-active manner can be  
 10 used for expression of a TF sequence in plants. Choice of a promoter is based largely on the phenotype of interest and is determined by such factors as tissue (e.g., seed, fruit, root, pollen, vascular tissue, flower, carpel, etc.), inducibility (e.g., in response to wounding, heat, cold, drought, light, pathogens, etc.), timing, developmental stage, and the like. Numerous known promoters have been characterized and can favorably be employed to promote expression of a  
 15 polynucleotide of the invention in a transgenic plant or cell of interest. For example, tissue specific promoters include: seed-specific promoters (such as the napin, phaseolin or DC3 promoter described in US Pat. No. 5,773,697), fruit-specific promoters that are active during fruit ripening (such as the *dru 1* promoter (US Pat. No. 5,783,393), or the 2A11 promoter (US Pat. No. 4,943,674) and the tomato polygalacturonase promoter (Bird et al. (1988) Plant Mol Biol 11:651),  
 20 root-specific promoters, such as those disclosed in US Patent Nos. 5,618,988, 5,837,848 and 5,905,186, pollen-active promoters such as PTA29, PTA26 and PTA13 (US Pat. No. 5,792,929), promoters active in vascular tissue (Ringli and Keller (1998) Plant Mol Biol 37:977-988), flower-specific (Kaiser et al. (1995) Plant Mol Biol 28:231-243), pollen (Baerson et al. (1994) Plant Mol Biol 26:1947-1959), carpels (Ohl et al. (1990) Plant Cell 2:837-848), pollen and ovules (Baerson et al. (1993) Plant Mol Biol 22:255-267), auxin-inducible promoters (such as that described in van der Kop et al. (1999) Plant Mol Biol 39:979-990 or Baumann et al. (1999) Plant Cell 11:323-334), cytokinin-inducible promoter (Guevara-Garcia (1998) Plant Mol Biol 38:743-753),  
 25 promoters responsive to gibberellin (Shi et al. (1998) Plant Mol Biol 38:1053-1060, Willmott et al. (1998) 38:817-825) and the like. Additional promoters are those that elicit expression in response to heat (Ainley et al. (1993) Plant Mol Biol 22: 13-23), light (e.g., the pea *rbcS-3A* promoter, Kuhlmeier et al. (1989) Plant Cell 1:471, and the maize *rbcS* promoter, Schaffner and Sheen (1991) Plant Cell 3: 997); wounding (e.g., *wun1*, Siebertz et al. (1989) Plant Cell 1: 961); pathogens (such as the PR-1 promoter described in Buchel et al. (1999) Plant Mol. Biol. 40:387-396, and the PDF1.2 promoter described in Manners et al. (1998) Plant Mol. Biol. 38:1071-80),  
 30

and chemicals such as methyl jasmonate or salicylic acid (Gatz et al. (1997) Plant Mol Biol 48: 89-108). In addition, the timing of the expression can be controlled by using promoters such as those acting at senescence (An and Amazon (1995) Science 270: 1986-1988); or late seed development (Odell et al. (1994) Plant Physiol 106:447-458).

5 Plant expression vectors can also include RNA processing signals that can be positioned within, upstream or downstream of the coding sequence. In addition, the expression vectors can include additional regulatory sequences from the 3'-untranslated region of plant genes, e.g., a 3' terminator region to increase mRNA stability of the mRNA, such as the PI-II terminator region of potato or the octopine or nopaline synthase 3' terminator regions.

#### 10 Additional Expression Elements

Specific initiation signals can aid in efficient translation of coding sequences. These signals can include, e.g., the ATG initiation codon and adjacent sequences. In cases where a coding sequence, its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases  
15 where only coding sequence (e.g., a mature protein coding sequence), or a portion thereof, is inserted, exogenous transcriptional control signals including the ATG initiation codon can be separately provided. The initiation codon is provided in the correct reading frame to facilitate transcription. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of  
20 enhancers appropriate to the cell system in use.

#### Expression Hosts

The present invention also relates to host cells which are transduced with vectors of the invention, and the production of polypeptides of the invention (including fragments thereof) by recombinant techniques. Host cells are genetically engineered (i.e, nucleic acids are introduced,  
25 e.g., transduced, transformed or transfected) with the vectors of this invention, which may be, for example, a cloning vector or an expression vector comprising the relevant nucleic acids herein. The vector is optionally a plasmid, a viral particle, a phage, a naked nucleic acids, *etc.* The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the relevant gene. The culture  
30 conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, Sambrook and Ausubel.

The host cell can be a eukaryotic cell, such as a yeast cell, or a plant cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Plant protoplasts are also suitable for some

applications. For example, the DNA fragments are introduced into plant tissues, cultured plant cells or plant protoplasts by standard methods including electroporation (Fromm et al., (1985) Proc. Natl. Acad. Sci. USA 82, 5824, infection by viral vectors such as cauliflower mosaic virus (CaMV) (Hohn et al., (1982) Molecular Biology of Plant Tumors, (Academic Press, New York) pp. 549-560; US 4,407,956), high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface (Klein et al., (1987) Nature 327, 70-73), use of pollen as vector (WO 85/01856), or use of *Agrobacterium tumefaciens* or *A. rhizogenes* carrying a T-DNA plasmid in which DNA fragments are cloned. The T-DNA plasmid is transmitted to plant cells upon infection by *Agrobacterium tumefaciens*, and a portion is stably integrated into the plant genome (Horsch et al. (1984) Science 233:496-498; Fraley et al. (1983) Proc. Natl. Acad. Sci. USA 80, 4803).

The cell can include a nucleic acid of the invention which encodes a polypeptide, wherein the cells expresses a polypeptide of the invention. The cell can also include vector sequences, or the like. Furthermore, cells and transgenic plants which include any polypeptide or nucleic acid above or throughout this specification, e.g., produced by transduction of a vector of the invention, are an additional feature of the invention.

For long-term, high-yield production of recombinant proteins, stable expression can be used. Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The protein or fragment thereof produced by a recombinant cell may be secreted, membrane-bound, or contained intracellularly, depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides encoding mature proteins of the invention can be designed with signal sequences which direct secretion of the mature polypeptides through a prokaryotic or eukaryotic cell membrane.

#### Modified Amino Acids

Polypeptides of the invention may contain one or more modified amino acids. The presence of modified amino acids may be advantageous in, for example, increasing polypeptide half-life, reducing polypeptide antigenicity or toxicity, increasing polypeptide storage stability, or the like. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production or modified by synthetic or chemical means.

Non-limiting examples of a modified amino acid include incorporation or other use of acetylated amino acids, glycosylated amino acids, sulfated amino acids, prenylated (e.g., farnesylated, geranylgeranylated) amino acids, PEG modified (e.g., "PEGylated") amino acids,



biotinylated amino acids, carboxylated amino acids, phosphorylated amino acids, etc. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature.

#### IDENTIFICATION OF ADDITIONAL FACTORS

5 A transcription factor provided by the present invention can also be used to identify additional endogenous or exogenous molecules that can affect a phenotype or trait of interest. On the one hand, such molecules include organic (small or large molecules) and/or inorganic compounds that affect expression of (i.e., regulate) a particular transcription factor. Alternatively, such molecules include endogenous molecules that are acted upon either at a transcriptional level by a transcription factor of the invention to modify a phenotype as desired. For example, the transcription factors can be employed to identify one or more downstream gene with which is subject to a regulatory effect of the transcription factor. In one approach, a transcription factor or transcription factor homologue of the invention is expressed in a host cell, e.g., a transgenic plant cell, tissue or explant, and expression products, either RNA or protein, of likely or random targets are monitored, e.g., by hybridization to a microarray of nucleic acid probes corresponding to genes expressed in a tissue or cell type of interest, by two-dimensional gel electrophoresis of protein products, or by any other method known in the art for assessing expression of gene products at the level of RNA or protein. Alternatively, a transcription factor of the invention can be used to identify promoter sequences (i.e., binding sites) involved in the regulation of a downstream target. After identifying a promoter sequence, interactions between the transcription factor and the promoter sequence can be modified by changing specific nucleotides in the promoter sequence or specific amino acids in the transcription factor that interact with the promoter sequence to alter a plant trait. Typically, transcription factor DNA binding sites are identified by gel shift assays. After identifying the promoter regions, the promoter region sequences can be employed in double-stranded DNA arrays to identify molecules that affect the interactions of the transcription factors with their promoters (Bulyk et al. (1999) Nature Biotechnology 17:573-577).

30 The identified transcription factors are also useful to identify proteins that modify the activity of the transcription factor. Such modification can occur by covalent modification, such as by phosphorylation, or by protein-protein (homo or-heteropolymer) interactions. Any method suitable for detecting protein-protein interactions can be employed. Among the methods that can be employed are co-immunoprecipitation, cross-linking and co-purification through gradients or chromatographic columns, and the two-hybrid yeast system.

The two-hybrid system detects protein interactions in vivo and is described in Chien, et al., (1991), Proc. Natl. Acad. Sci. USA 88, 9578-9582 and is commercially available from Clontech (Palo Alto, Calif.). In such a system, plasmids are constructed that encode two hybrid proteins: one consists of the DNA-binding domain of a transcription activator protein fused to the TF polypeptide and the other consists of the transcription activator protein's activation domain fused to an unknown protein that is encoded by a cDNA that has been recombined into the plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the cDNA library are transformed into a strain of the yeast *Saccharomyces cerevisiae* that contains a reporter gene (e.g., lacZ) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene. Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product. Then, the library plasmids responsible for reporter gene expression are isolated and sequenced to identify the proteins encoded by the library plasmids. After identifying proteins that interact with the transcription factors, assays for compounds that interfere with the TF protein-protein interactions can be preformed.

#### IDENTIFICATION OF MODULATORS

In addition to the intracellular molecules described above, extracellular molecules that alter activity or expression of a transcription factor, either directly or indirectly, can be identified. For example, the methods can entail first placing a candidate molecule in contact with a plant or plant cell. The molecule can be introduced by topical administration, such as spraying or soaking of a plant, and then the molecule's effect on the expression or activity of the TF polypeptide or the expression of the polynucleotide monitored. Changes in the expression of the TF polypeptide can be monitored by use of polyclonal or monoclonal antibodies, gel electrophoresis or the like. Changes in the expression of the corresponding polynucleotide sequence can be detected by use of microarrays, Northern, quantitative PCR, or any other technique for monitoring changes in mRNA expression. These techniques are exemplified in Ausubel et al. (eds) Current Protocols in Molecular Biology, John Wiley & Sons (1998). Such changes in the expression levels can be correlated with modified plant traits and thus identified molecules can be useful for soaking or spraying on fruit, vegetable and grain crops to modify traits in plants.

Essentially any available composition can be tested for modulatory activity of expression or activity of any nucleic acid or polypeptide herein. Thus, available libraries of compounds such as chemicals, polypeptides, nucleic acids and the like can be tested for modulatory activity.

Often, potential modulator compounds can be dissolved in aqueous or organic (e.g., DMSO-based) solutions for easy delivery to the cell or plant of interest in which the activity of the modulator is to be tested. Optionally, the assays are designed to screen large modulator composition libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays).

In one embodiment, high throughput screening methods involve providing a combinatorial library containing a large number of potential compounds (potential modulator compounds). Such "combinatorial chemical libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as target compounds.

A combinatorial chemical library can be, e.g., a collection of diverse chemical compounds generated by chemical synthesis or biological synthesis. For example, a combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (e.g., in one example, amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound of a set length). Exemplary libraries include peptide libraries, nucleic acid libraries, antibody libraries (see, e.g., Vaughn et al. (1996) Nature Biotechnology, 14(3):309-314 and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al. Science (1996) 274:1520-1522 and U.S. Patent 5,593,853), peptide nucleic acid libraries (see, e.g., U.S. Patent 5,539,083), and small organic molecule libraries (see, e.g., benzodiazepines, Baum C&EN Jan 18, page 33 (1993); isoprenoids, U.S. Patent 5,569,588; thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pyrrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337) and the like.

Preparation and screening of combinatorial or other libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Patent 5,010,175, Furka, Int. J. Pept. Prot. Res. 37:487-493 (1991) and Houghton et al. Nature 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used.

In addition, as noted, compound screening equipment for high-throughput screening is generally available, e.g., using any of a number of well known robotic systems that have also been developed for solution phase chemistries useful in assay systems. These systems include automated workstations including an automated synthesis apparatus and robotic systems utilizing robotic arms. Any of the above devices are suitable for use with the present invention, e.g., for

high-throughput screening of potential modulators. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art.

Indeed, entire high throughput screening systems are commercially available. These systems typically automate entire procedures including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. Similarly, microfluidic implementations of screening are also commercially available.

The manufacturers of such systems provide detailed protocols the various high throughput. Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like. The integrated systems herein, in addition to providing for sequence alignment and, optionally, synthesis of relevant nucleic acids, can include such screening apparatus to identify modulators that have an effect on one or more polynucleotides or polypeptides according to the present invention.

In some assays it is desirable to have positive controls to ensure that the components of the assays are working properly. At least two types of positive controls are appropriate. That is, known transcriptional activators or inhibitors can be incubated with cells/plants/ etc. in one sample of the assay, and the resulting increase/decrease in transcription can be detected by measuring the resulting increase in RNA/ protein expression, etc., according to the methods herein. It will be appreciated that modulators can also be combined with transcriptional activators or inhibitors to find modulators which inhibit transcriptional activation or transcriptional repression. Either expression of the nucleic acids and proteins herein or any additional nucleic acids or proteins activated by the nucleic acids or proteins herein, or both, can be monitored.

In an embodiment, the invention provides a method for identifying compositions that modulate the activity or expression of a polynucleotide or polypeptide of the invention. For example, a test compound, whether a small or large molecule, is placed in contact with a cell, plant (or plant tissue or explant), or composition comprising the polynucleotide or polypeptide of interest and a resulting effect on the cell, plant, (or tissue or explant) or composition is evaluated by monitoring, either directly or indirectly, one or more of: expression level of the polynucleotide or polypeptide, activity (or modulation of the activity) of the polynucleotide or polypeptide. In some cases, an alteration in a plant phenotype can be detected following contact of a plant (or

plant cell, or tissue or explant) with the putative modulator, e.g., by modulation of expression or activity of a polynucleotide or polypeptide of the invention.

### SUBSEQUENCES

5 Also contemplated are uses of polynucleotides, also referred to herein as oligonucleotides, typically having at least 12 bases, preferably at least 15, more preferably at least 20, 30, or 50 bases, which hybridize under at least highly stringent (or ultra-high stringent or ultra-ultra- high stringent conditions) conditions to a polynucleotide sequence described above. The polynucleotides may be used as probes, primers, sense and antisense agents, and the like,  
10 according to methods as noted *supra*.

Subsequences of the polynucleotides of the invention, including polynucleotide fragments and oligonucleotides are useful as nucleic acid probes and primers. An oligonucleotide suitable for use as a probe or primer is at least about 15 nucleotides in length, more often at least about 18 nucleotides, often at least about 21 nucleotides, frequently at least about 30 nucleotides,  
15 or about 40 nucleotides, or more in length. A nucleic acid probe is useful in hybridization protocols, e.g., to identify additional polypeptide homologues of the invention, including protocols for microarray experiments. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then extended along the target DNA strand by a DNA polymerase enzyme. Primer  
20 pairs can be used for amplification of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other nucleic-acid amplification methods. See Sambrook and Ausubel, *supra*.

In addition, the invention includes an isolated or recombinant polypeptide including a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotides of the invention. For example, such polypeptides, or domains or fragments  
25 thereof, can be used as immunogens, e.g., to produce antibodies specific for the polypeptide sequence, or as probes for detecting a sequence of interest. A subsequence can range in size from about 15 amino acids in length up to and including the full length of the polypeptide.

### PRODUCTION OF TRANSGENIC PLANTS

#### Modification of Traits

30 The polynucleotides of the invention are favorably employed to produce transgenic plants with various traits, or characteristics, that have been modified in a desirable manner, e.g., to improve the seed characteristics of a plant. For example, alteration of expression levels or patterns (e.g., spatial or temporal expression patterns) of one or more of the transcription factors

(or transcription factor homologues) of the invention, as compared with the levels of the same protein found in a wild type plant, can be used to modify a plant's traits. An illustrative example of trait modification, improved biochemical characteristics, by altering expression levels of a particular transcription factor is described further in the Examples and the Sequence Listing.

5

#### Antisense and Cosuppression Approaches

In addition to expression of the nucleic acids of the invention as gene replacement or plant phenotype modification nucleic acids, the nucleic acids are also useful for sense and anti-sense suppression of expression, e.g., to down-regulate expression of a nucleic acid of the invention, e.g., as a further mechanism for modulating plant phenotype. That is, the nucleic acids  
10 of the invention, or subsequences or anti-sense sequences thereof, can be used to block expression of naturally occurring homologous nucleic acids. A variety of sense and anti-sense technologies are known in the art, e.g., as set forth in Lichtenstein and Nellen (1997) Antisense Technology: A Practical Approach IRL Press at Oxford University, Oxford, England. In general, sense or anti-sense sequences are introduced into a cell, where they are optionally amplified, e.g., by  
15 transcription. Such sequences include both simple oligonucleotide sequences and catalytic sequences such as ribozymes.

For example, a reduction or elimination of expression (i.e., a "knock-out") of a transcription factor or transcription factor homologue polypeptide in a transgenic plant, e.g., to modify a plant trait, can be obtained by introducing an antisense construct corresponding to the  
20 polypeptide of interest as a cDNA. For antisense suppression, the transcription factor or homologue cDNA is arranged in reverse orientation (with respect to the coding sequence) relative to the promoter sequence in the expression vector. The introduced sequence need not be the full length cDNA or gene, and need not be identical to the cDNA or gene found in the plant type to be transformed. Typically, the antisense sequence need only be capable of hybridizing to the target  
25 gene or RNA of interest. Thus, where the introduced sequence is of shorter length, a higher degree of homology to the endogenous transcription factor sequence will be needed for effective antisense suppression. While antisense sequences of various lengths can be utilized, preferably, the introduced antisense sequence in the vector will be at least 30 nucleotides in length, and improved antisense suppression will typically be observed as the length of the antisense sequence  
30 increases. Preferably, the length of the antisense sequence in the vector will be greater than 100 nucleotides. Transcription of an antisense construct as described results in the production of RNA molecules that are the reverse complement of mRNA molecules transcribed from the endogenous transcription factor gene in the plant cell.

Suppression of endogenous transcription factor gene expression can also be achieved using a ribozyme. Ribozymes are RNA molecules that possess highly specific endoribonuclease activity. The production and use of ribozymes are disclosed in U.S. Patent No. 4,987,071 and U.S. Patent No. 5,543,508. Synthetic ribozyme sequences including antisense RNAs can be used to confer RNA cleaving activity on the antisense RNA, such that endogenous mRNA molecules that hybridize to the antisense RNA are cleaved, which in turn leads to an enhanced antisense inhibition of endogenous gene expression.

Vectors in which RNA encoded by a transcription factor or transcription factor homologue cDNA is over-expressed can also be used to obtain co-suppression of a corresponding endogenous gene, e.g., in the manner described in U.S. Patent No. 5,231,020 to Jorgensen. Such co-suppression (also termed sense suppression) does not require that the entire transcription factor cDNA be introduced into the plant cells, nor does it require that the introduced sequence be exactly identical to the endogenous transcription factor gene of interest. However, as with antisense suppression, the suppressive efficiency will be enhanced as specificity of hybridization is increased, e.g., as the introduced sequence is lengthened, and/or as the sequence similarity between the introduced sequence and the endogenous transcription factor gene is increased.

Vectors expressing an untranslatable form of the transcription factor mRNA, e.g., sequences comprising one or more stop codon, or nonsense mutation) can also be used to suppress expression of an endogenous transcription factor, thereby reducing or eliminating its activity and modifying one or more traits. Methods for producing such constructs are described in U.S. Patent No. 5,583,021. Preferably, such constructs are made by introducing a premature stop codon into the transcription factor gene. Alternatively, a plant trait can be modified by gene silencing using double-strand RNA (Sharp (1999) Genes and Development 13: 139-141).

Another method for abolishing the expression of a gene is by insertion mutagenesis using the T-DNA of *Agrobacterium tumefaciens*. After generating the insertion mutants, the mutants can be screened to identify those containing the insertion in a transcription factor or transcription factor homologue gene. Plants containing a single transgene insertion event at the desired gene can be crossed to generate homozygous plants for the mutation (Koncz et al. (1992) Methods in Arabidopsis Research, World Scientific).

Alternatively, a plant phenotype can be altered by eliminating an endogenous gene, such as a transcription factor or transcription factor homologue, e.g., by homologous recombination (Kempin et al. (1997) Nature 389:802).

A plant trait can also be modified by using the cre-lox system (for example, as described in US Pat. No. 5,658,772). A plant genome can be modified to include first and second lox sites

that are then contacted with a Cre recombinase. If the lox sites are in the same orientation, the intervening DNA sequence between the two sites is excised. If the lox sites are in the opposite orientation, the intervening sequence is inverted.

5 The polynucleotides and polypeptides of this invention can also be expressed in a plant in the absence of an expression cassette by manipulating the activity or expression level of the endogenous gene by other means. For example, by ectopically expressing a gene by T-DNA activation tagging (Ichikawa et al. (1997) Nature 390 698-701; Kakimoto et al. (1996) Science 274: 982-985). This method entails transforming a plant with a gene tag containing multiple transcriptional enhancers and once the tag has inserted into the genome, expression of a flanking  
10 gene coding sequence becomes deregulated. In another example, the transcriptional machinery in a plant can be modified so as to increase transcription levels of a polynucleotide of the invention (See, e.g., PCT Publications WO 96/06166 and WO 98/53057 which describe the modification of the DNA binding specificity of zinc finger proteins by changing particular amino acids in the DNA binding motif).

15 The transgenic plant can also include the machinery necessary for expressing or altering the activity of a polypeptide encoded by an endogenous gene, for example by altering the phosphorylation state of the polypeptide to maintain it in an activated state.

Transgenic plants (or plant cells, or plant explants, or plant tissues) incorporating the polynucleotides of the invention and/or expressing the polypeptides of the invention can be  
20 produced by a variety of well established techniques as described above. Following construction of a vector, most typically an expression cassette, including a polynucleotide, e.g., encoding a transcription factor or transcription factor homologue, of the invention, standard techniques can be used to introduce the polynucleotide into a plant, a plant cell, a plant explant or a plant tissue of interest. Optionally, the plant cell, explant or tissue can be regenerated to produce a transgenic  
25 plant.

The plant can be any higher plant, including gymnosperms, monocotyledonous and dicotyledonous plants. Suitable protocols are available for *Leguminosae* (alfalfa, soybean, clover, etc.), *Umbelliferae* (carrot, celery, parsnip), *Cruciferae* (cabbage, radish, rapeseed, broccoli, etc.), *Curcubitaceae* (melons and cucumber), *Gramineae* (wheat, corn, rice, barley, millet, etc.),  
30 *Solanaceae* (potato, tomato, tobacco, peppers, etc.), and various other crops. See protocols described in Ammirato et al. (1984) Handbook of Plant Cell Culture –Crop Species. Macmillan Publ. Co. Shimamoto et al. (1989) Nature 338:274-276; Fromm et al. (1990) Bio/Technology 8:833-839; and Vasil et al. (1990) Bio/Technology 8:429-434.



Transformation and regeneration of both monocotyledonous and dicotyledonous plant cells is now routine, and the selection of the most appropriate transformation technique will be determined by the practitioner. The choice of method will vary with the type of plant to be transformed; those skilled in the art will recognize the suitability of particular methods for given plant types. Suitable methods can include, but are not limited to: electroporation of plant protoplasts; liposome-mediated transformation; polyethylene glycol (PEG) mediated transformation; transformation using viruses; micro-injection of plant cells; micro-projectile bombardment of plant cells; vacuum infiltration; and *Agrobacterium tumefaciens* mediated transformation. Transformation means introducing a nucleotide sequence in a plant in a manner to cause stable or transient expression of the sequence.

Successful examples of the modification of plant characteristics by transformation with cloned sequences which serve to illustrate the current knowledge in this field of technology, and which are herein incorporated by reference, include: U.S. Patent Nos. 5,571,706; 5,677,175; 5,510,471; 5,750,386; 5,597,945; 5,589,615; 5,750,871; 5,268,526; 5,780,708; 5,538,880; 5,773,269; 5,736,369 and 5,610,042.

Following transformation, plants are preferably selected using a dominant selectable marker incorporated into the transformation vector. Typically, such a marker will confer antibiotic or herbicide resistance on the transformed plants, and selection of transformants can be accomplished by exposing the plants to appropriate concentrations of the antibiotic or herbicide.

After transformed plants are selected and grown to maturity, those plants showing a modified trait are identified. The modified trait can be any of those traits described above. Additionally, to confirm that the modified trait is due to changes in expression levels or activity of the polypeptide or polynucleotide of the invention can be determined by analyzing mRNA expression using Northern blots, RT-PCR or microarrays, or protein expression using immunoblots or Western blots or gel shift assays.

#### INTEGRATED SYSTEMS—SEQUENCE IDENTITY

Additionally, the present invention may be an integrated system, computer or computer readable medium that comprises an instruction set for determining the identity of one or more sequences in a database. In addition, the instruction set can be used to generate or identify sequences that meet any specified criteria. Furthermore, the instruction set may be used to associate or link certain functional benefits, such improved biochemical characteristics, with one or more identified sequence.

For example, the instruction set can include, e.g., a sequence comparison or other alignment program, e.g., an available program such as, for example, the Wisconsin Package Version 10.0, such as BLAST, FASTA, PILEUP, FINDPATTERNS or the like (GCG, Madison, WI). Public sequence databases such as GenBank, EMBL, Swiss-Prot and PIR or private  
5 sequence databases such as PhytoSeq (Incyte Pharmaceuticals, Palo Alto, CA) can be searched.

Alignment of sequences for comparison can be conducted by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity method of Pearson and Lipman (1988) Proc. Natl. Acad. Sci. U.S.A. 85: 2444, by computerized  
10 implementations of these algorithms. After alignment, sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a comparison window to identify and compare local regions of sequence similarity. The comparison window can be a segment of at least about 20 contiguous positions, usually about 50 to about 200, more usually about 100 to about 150 contiguous positions. A  
15 description of the method is provided in Ausubel et al., *supra*.

A variety of methods of determining sequence relationships can be used, including manual alignment and computer assisted sequence alignment and analysis. This later approach is a preferred approach in the present invention, due to the increased throughput afforded by computer assisted methods. As noted above, a variety of computer programs for performing  
20 sequence alignment are available, or can be produced by one of skill.

One example algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al. J. Mol. Biol. 215:403-410 (1990). Software for performing BLAST analyses is publicly available, e.g., through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This  
25 algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them.  
30 The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each

direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment.

5 The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (*see* Henikoff & Henikoff (1989) Proc. Natl. Acad. Sci. USA 89:10915).

10 In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (*see*, e.g., Karlin & Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For  
15 example, a nucleic acid is considered similar to a reference sequence (and, therefore, in this context, homologous) if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, or less than about 0.01, and or even less than about 0.001. An additional example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using  
20 progressive, pairwise alignments. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters.

The integrated system, or computer typically includes a user input interface allowing a user to selectively view one or more sequence records corresponding to the one or more character strings, as well as an instruction set which aligns the one or more character strings with each other  
25 or with an additional character string to identify one or more region of sequence similarity. The system may include a link of one or more character strings with a particular phenotype or gene function. Typically, the system includes a user readable output element which displays an alignment produced by the alignment instruction set.

The methods of this invention can be implemented in a localized or distributed  
30 computing environment. In a distributed environment, the methods may implemented on a single computer comprising multiple processors or on a multiplicity of computers. The computers can be linked, e.g. through a common bus, but more preferably the computer(s) are nodes on a network. The network can be a generalized or a dedicated local or wide-area network and, in certain preferred embodiments, the computers may be components of an intra-net or an internet.

Thus, the invention provides methods for identifying a sequence similar or homologous to one or more polynucleotides as noted herein, or one or more target polypeptides encoded by the polynucleotides, or otherwise noted herein and may include linking or associating a given plant phenotype or gene function with a sequence. In the methods, a sequence database is provided (locally or across an inter or intra net) and a query is made against the sequence database using the relevant sequences herein and associated plant phenotypes or gene functions.

Any sequence herein can be entered into the database, before or after querying the database. This provides for both expansion of the database and, if done before the querying step, for insertion of control sequences into the database. The control sequences can be detected by the query to ensure the general integrity of both the database and the query. As noted, the query can be performed using a web browser based interface. For example, the database can be a centralized public database such as those noted herein, and the querying can be done from a remote terminal or computer across an internet or intranet.

### EXAMPLES

The following examples are intended to illustrate but not limit the present invention.

#### EXAMPLE I. FULL LENGTH GENE IDENTIFICATION AND CLONING

Putative transcription factor sequences (genomic or ESTs) related to known transcription factors were identified in the *Arabidopsis thaliana* GenBank database using the tblastn sequence analysis program using default parameters and a P-value cutoff threshold of  $-4$  or  $-5$  or lower, depending on the length of the query sequence. Putative transcription factor sequence hits were then screened to identify those containing particular sequence strings. If the sequence hits contained such sequence strings, the sequences were confirmed as transcription factors.

Alternatively, *Arabidopsis thaliana* cDNA libraries derived from different tissues or treatments, or genomic libraries were screened to identify novel members of a transcription family using a low stringency hybridization approach. Probes were synthesized using gene specific primers in a standard PCR reaction (annealing temperature  $60^{\circ}\text{C}$ ) and labeled with  $^{32}\text{P}$  dCTP using the High Prime DNA Labeling Kit (Boehringer Mannheim). Purified radiolabelled probes were added to filters immersed in Church hybridization medium (0.5 M  $\text{NaPO}_4$  pH 7.0, 7% SDS, 1 % w/v bovine serum albumin) and hybridized overnight at  $60^{\circ}\text{C}$  with shaking. Filters were washed two times for 45 to 60 minutes with 1xSSC, 1% SDS at  $60^{\circ}\text{C}$ .

To identify additional sequence 5' or 3' of a partial cDNA sequence in a cDNA library, 5' and 3' rapid amplification of cDNA ends (RACE) was performed using the Marathon<sup>TM</sup> cDNA amplification kit (Clontech, Palo Alto, CA). Generally, the method entailed first isolating

poly(A) mRNA, performing first and second strand cDNA synthesis to generate double stranded cDNA, blunting cDNA ends, followed by ligation of the Marathon™ Adaptor to the cDNA to form a library of adaptor-ligated ds cDNA.

Gene-specific primers were designed to be used along with adaptor specific primers for both 5' and 3' RACE reactions. Nested primers, rather than single primers, were used to increase PCR specificity. Using 5' and 3' RACE reactions, 5' and 3' RACE fragments were obtained, sequenced and cloned. The process can be repeated until 5' and 3' ends of the full-length gene were identified. Then the full-length cDNA was generated by PCR using primers specific to 5' and 3' ends of the gene by end-to-end PCR.

#### 10      EXAMPLE II. CONSTRUCTION OF EXPRESSION VECTORS

The sequence was amplified from a genomic or cDNA library using primers specific to sequences upstream and downstream of the coding region. The expression vector was pMEN20 or pMEN65, which are both derived from pMON316 (Sanders et al, (1987) Nucleic Acids Research 15:1543-58) and contain the CaMV 35S promoter to express transgenes. To clone the sequence into the vector, both pMEN20 and the amplified DNA fragment were digested separately with SalI and NotI restriction enzymes at 37° C for 2 hours. The digestion products were subject to electrophoresis in a 0.8% agarose gel and visualized by ethidium bromide staining. The DNA fragments containing the sequence and the linearized plasmid were excised and purified by using a Qiaquick gel extraction kit (Qiagen, CA). The fragments of interest were ligated at a ratio of 3:1 (vector to insert). Ligation reactions using T4 DNA ligase (New England Biolabs, MA) were carried out at 16° C for 16 hours. The ligated DNAs were transformed into competent cells of the *E. coli* strain DH5alpha by using the heat shock method. The transformations were plated on LB plates containing 50 mg/l kanamycin (Sigma).

Individual colonies were grown overnight in five milliliters of LB broth containing 50 mg/l kanamycin at 37° C. Plasmid DNA was purified by using Qiaquick Mini Prep kits (Qiagen, CA).

#### 25      EXAMPLE III. TRANSFORMATION OF AGROBACTERIUM WITH THE EXPRESSION VECTOR

After the plasmid vector containing the gene was constructed, the vector was used to transform *Agrobacterium tumefaciens* cells expressing the gene products. The stock of *Agrobacterium tumefaciens* cells for transformation were made as described by Nagel et al. (1990) FEMS Microbiol Letts. 67: 325-328. *Agrobacterium* strain ABI was grown in 250 ml LB medium (Sigma) overnight at 28°C with shaking until an absorbance ( $A_{600}$ ) of 0.5 – 1.0 was



reached. Cells were harvested by centrifugation at 4,000 x g for 15 min at 4° C. Cells were then resuspended in 250 µl chilled buffer (1 mM HEPES, pH adjusted to 7.0 with KOH). Cells were centrifuged again as described above and resuspended in 125 µl chilled buffer. Cells were then centrifuged and resuspended two more times in the same HEPES buffer as described above at a volume of 100 µl and 750 µl, respectively. Resuspended cells were then distributed into 40 µl aliquots, quickly frozen in liquid nitrogen, and stored at -80° C.

*Agrobacterium* cells were transformed with plasmids prepared as described above following the protocol described by Nagel et al. For each DNA construct to be transformed, 50 – 100 ng DNA (generally resuspended in 10 mM Tris-HCl, 1 mM EDTA, pH 8.0) was mixed with 40 µl of *Agrobacterium* cells. The DNA/cell mixture was then transferred to a chilled cuvette with a 2mm electrode gap and subject to a 2.5 kV charge dissipated at 25 µF and 200 µF using a Gene Pulser II apparatus (Bio-Rad). After electroporation, cells were immediately resuspended in 1.0 ml LB and allowed to recover without antibiotic selection for 2 – 4 hours at 28° C in a shaking incubator. After recovery, cells were plated onto selective medium of LB broth containing 100 µg/ml spectinomycin (Sigma) and incubated for 24-48 hours at 28° C. Single colonies were then picked and inoculated in fresh medium. The presence of the plasmid construct was verified by PCR amplification and sequence analysis.

#### EXAMPLE IV. TRANSFORMATION OF ARABIDOPSIS PLANTS WITH AGROBACTERIUM TUMEFACIENS WITH EXPRESSION VECTOR

After transformation of *Agrobacterium tumefaciens* with plasmid vectors containing the gene, single *Agrobacterium* colonies were identified, propagated, and used to transform *Arabidopsis* plants. Briefly, 500 ml cultures of LB medium containing 50 mg/l kanamycin were inoculated with the colonies and grown at 28° C with shaking for 2 days until an absorbance ( $A_{600}$ ) of > 2.0 is reached. Cells were then harvested by centrifugation at 4,000 x g for 10 min, and resuspended in infiltration medium (1/2 X Murashige and Skoog salts (Sigma), 1 X Gamborg's B-5 vitamins (Sigma), 5.0% (w/v) sucrose (Sigma), 0.044 µM benzylamino purine (Sigma), 200 µl/L Silwet L-77 (Lehle Seeds) until an absorbance ( $A_{600}$ ) of 0.8 was reached.

Prior to transformation, *Arabidopsis thaliana* seeds (ecotype Columbia) were sown at a density of ~10 plants per 4" pot onto Pro-Mix BX potting medium (Hummert International) covered with fiberglass mesh (18 mm X 16 mm). Plants were grown under continuous illumination (50-75 µE/m<sup>2</sup>/sec) at 22-23° C with 65-70% relative humidity. After about 4 weeks, primary inflorescence stems (bolts) are cut off to encourage growth of multiple secondary bolts. After flowering of the mature secondary bolts, plants were prepared for transformation by removal of all siliques and opened flowers.

The pots were then immersed upside down in the mixture of *Agrobacterium* infiltration medium as described above for 30 sec, and placed on their sides to allow draining into a 1' x 2' flat surface covered with plastic wrap. After 24 h, the plastic wrap was removed and pots are turned upright. The immersion procedure was repeated one week later, for a total of two  
5 immersions per pot. Seeds were then collected from each transformation pot and analyzed following the protocol described below.

#### EXAMPLE V. IDENTIFICATION OF ARABIDOPSIS PRIMARY TRANSFORMANTS

Seeds collected from the transformation pots were sterilized essentially as follows. Seeds  
10 were dispersed into in a solution containing 0.1% (v/v) Triton X-100 (Sigma) and sterile H<sub>2</sub>O and washed by shaking the suspension for 20 min. The wash solution was then drained and replaced with fresh wash solution to wash the seeds for 20 min with shaking. After removal of the second wash solution, a solution containing 0.1% (v/v) Triton X-100 and 70% ethanol (Equistar) was added to the seeds and the suspension was shaken for 5 min. After removal of the  
15 ethanol/detergent solution, a solution containing 0.1% (v/v) Triton X-100 and 30% (v/v) bleach (Clorox) was added to the seeds, and the suspension was shaken for 10 min. After removal of the bleach/detergent solution, seeds were then washed five times in sterile distilled H<sub>2</sub>O. The seeds were stored in the last wash water at 4° C for 2 days in the dark before being plated onto antibiotic selection medium (1 X Murashige and Skoog salts (pH adjusted to 5.7 with 1M KOH), 1 X  
20 Gamborg's B-5 vitamins, 0.9% phytagar (Life Technologies), and 50 mg/l kanamycin). Seeds were germinated under continuous illumination (50-75  $\mu\text{E}/\text{m}^2/\text{sec}$ ) at 22-23° C. After 7-10 days of growth under these conditions, kanamycin resistant primary transformants (T<sub>1</sub> generation) were visible and obtained. These seedlings were transferred first to fresh selection plates where the seedlings continued to grow for 3-5 more days, and then to soil (Pro-Mix BX potting  
25 medium).

Primary transformants were crossed and progeny seeds (T<sub>2</sub>) collected; kanamycin resistant seedlings were selected and analyzed. The expression levels of the recombinant polynucleotides in the transformants varies from about a 5% expression level increase to a least a  
30 100% expression level increase. Similar observations are made with respect to polypeptide level expression.



#### EXAMPLE VI. IDENTIFICATION OF ARABIDOPSIS PLANTS WITH TRANSCRIPTION FACTOR GENE KNOCKOUTS

The screening of insertion mutagenized *Arabidopsis* collections for null mutants in a known target gene was essentially as described in Krysan et al (1999) Plant Cell 11:2283-2290.

- 5 Briefly, gene-specific primers, nested by 5-250 base pairs to each other, were designed from the 5' and 3' regions of a known target gene. Similarly, nested sets of primers were also created specific to each of the T-DNA or transposon ends (the "right" and "left" borders). All possible combinations of gene specific and T-DNA/transposon primers were used to detect by PCR an insertion event within or close to the target gene. The amplified DNA fragments were then
- 10 sequenced which allows the precise determination of the T-DNA/transposon insertion point relative to the target gene. Insertion events within the coding or intervening sequence of the genes were deconvoluted from a pool comprising a plurality of insertion events to a single unique mutant plant for functional characterization. The method is described in more detail in Yu and Adam, US Application Serial No. 09/177,733 filed October 23, 1998.

#### EXAMPLE VII. IDENTIFICATION OF MODIFIED BIOCHEMICAL CHARACTERISTICS PHENOTYPE IN OVEREXPRESSOR OR GENE KNOCKOUT PLANTS

- Experiments were performed to identify those transformants or knockouts that exhibited modified biochemical characteristics. Among the biochemicals that were assayed were insoluble
- 20 sugars, such as arabinose, fucose, galactose, mannose, rhamnose or xylose or the like; prenyl lipids, such as lutein, beta-carotene, xanthophyll-1, xanthophyll-2, chlorophylls A or B, or alpha-, delta- or gamma-tocopherol or the like; fatty acids, such as 16:0 (palmitic acid), 16:1 (palmitoleic acid), 18:0 (stearic acid), 18:1 (oleic acid), 18:2 (linoleic acid), 20:0, 18:3 (linolenic acid), 20:1 (eicosenoic acid), 20:2, 22:1 (erucic acid) or the like; waxes, such as by altering the levels of C29,
- 25 C31, or C33 alkanes; sterols, such as brassicasterol, campesterol, stigmasterol, sitosterol or stigmastanol or the like, glucosinolates, protein or oil levels

- Fatty acids were measured using two methods depending on whether the tissue was from leaves or seeds. For leaves, lipids were extracted and esterified with hot methanolic H<sub>2</sub>SO<sub>4</sub> and partitioned into hexane from methanolic brine. For seed fatty acids, seeds were pulverized and
- 30 extracted in methanol:heptane:toluene:2,2-dimethoxypropane:H<sub>2</sub>SO<sub>4</sub> (39:34:20:5:2) for 90 minutes at 80°C. After cooling to room temperature the upper phase, containing the seed fatty acid esters, was subjected to GC analysis. Fatty acid esters from both seed and leaf tissues were analyzed with a Supelco SP-2330 column.

Glucosinolates were purified from seeds or leaves by first heating the tissue at 95°C for 10 minutes. Preheated ethanol:water (50:50) is and after heating at 95°C for a further 10 minutes, the extraction solvent is applied to a DEAE Sephadex column which had been previously equilibrated with 0.5 M pyridine acetate. Desulfoglucosinolates were eluted with 300 ul water and analyzed by reverse phase HPLC monitoring at 226 nm.

For wax alkanes, samples were extracted using an identical method as fatty acids and extracts were analyzed on a HP 5890 GC coupled with a 5973 MSD. Samples were chromatographed on a J&W DB35 mass spectrometer (J&W Scientific).

To measure prenyl lipids levels, seeds or leaves were pulverized with 1 to 2% pyrogallol as an antioxidant. For seeds, extracted samples were filtered and a portion removed for tocopherol and carotenoid/chlorophyll analysis by HPLC. The remaining material was saponified for sterol determination. For leaves, an aliquot was removed and diluted with methanol and chlorophyll A, chlorophyll B, and total carotenoids measured by spectrophotometry by determining absorbance at 665.2 nm, 652.5 nm, and 470 nm. An aliquot was removed for tocopherol and carotenoid/chlorophyll composition by HPLC using a Waters uBondapak C18 column (4.6 mm x 150 mm). The remaining methanolic solution was saponified with 10% KOH at 80°C for one hour. The samples were cooled and diluted with a mixture of methanol and water. A solution of 2% methylene chloride in hexane was mixed in and the samples were centrifuged. The aqueous methanol phase was again re-extracted 2% methylene chloride in hexane and, after centrifugation, the two upper phases were combined and evaporated. 2% methylene chloride in hexane was added to the tubes and the samples were then extracted with one ml of water. The upper phase was removed, dried, and resuspended in 400 ul of 2% methylene chloride in hexane and analyzed by gas chromatography using a 50 m DB-5ms (0.25 mm ID, 0.25 um phase, J&W Scientific).

Insoluble sugar levels were measured by the method essentially described by Reiter et al., Plant Journal 12:335-345. This method analyzes the neutral sugar composition of cell wall polymers found in *Arabidopsis* leaves. Soluble sugars were separated from sugar polymers by extracting leaves with hot 70% ethanol. The remaining residue containing the insoluble polysaccharides was then acid hydrolyzed with allose added as an internal standard. Sugar monomers generated by the hydrolysis were then reduced to the corresponding alditols by treatment with NaBH<sub>4</sub>, then were acetylated to generate the volatile alditol acetates which were then analyzed by GC-FID. Identity of the peaks was determined by comparing the retention times of known sugars converted to the corresponding alditol acetates with the retention times of peaks from wild-type plant extracts. Alditol acetates were analyzed on a Supelco SP-2330 capillary

column (30 m x 250  $\mu$ m x 0.2  $\mu$ m) using a temperature program beginning at 180° C for 2 minutes followed by an increase to 220° C in 4 minutes. After holding at 220° C for 10 minutes, the oven temperature is increased to 240° C in 2 minutes and held at this temperature for 10 minutes and brought back to room temperature.

- 5 To identify plants with alterations in total seed oil or protein content, 150mg of seeds from T2 progeny plants were subjected to analysis by Near Infrared Reflectance (NIR) using a Foss NirSystems Model 6500 with a spinning cup transport system.

- Table 3 shows the phenotypes observed for particular overexpressor or knockout plants and provides the SEQ ID No., the internal reference code (GID), whether a knockout or  
10 overexpressor plant was analyzed and the observed phenotype.

**Table 3**

SEQ ID No.	GID	Knockout (KO) or overexpressor (OE)	Phenotype observed
1	G214	OE	Increase in leaf fatty acids, for example 100% increase in 18:0 fatty acid. Also up to 100% increase in leaf chlorophyll and 100% increase in leaf carotenoids
3	G231	OE	Up to 5% increase in leaf 18:3 fatty acid
5	G274	OE	Up to 50% increase in leaf arabinose
7	G307	OE	Altered in leaf insoluble sugars, for example up to 44% decrease in mannose.
9	G346	OE	Altered leaf fatty acids, for example 25% increase in 16:3 and altered insoluble sugars, for example up to 25% increase in fucose
11	G598	OE	Altered in insoluble sugars, for example up to 20% decrease in rhamnose and up to 10% increase in galactose
13	G605	OE	Altered in leaf fatty acids, for example up to 20% increase in 16:1 fatty acid.
15	G777	OE	Altered in insoluble sugars, for example up to 60% increase in leaf rhamnose
17	G869	OE	Alteration in leaf fatty acids eg up to 39% decrease in 16:0 fatty acid; up to 43% increase in fucose
19	G1133	OE	Up to 34% decrease in leaf lutein
21	G1266	OE	Alteration in leaf fatty acids, for example up to 50% increase in 18:0 fatty acid. Alterations in leaf insoluble sugars, for example a 45% decrease in rhamnose
23	G1324	OE	Up to 65% decrease in leaf lutein and up to 84% increase in leaf xanthophyll

25	G1337	OE	Alteration in leaf fatty acids, for example up to 28% increase in 18:1 fatty acid
27	G975	OE	Up to 13-fold increase in wax in leaves

For a particular overexpressor that shows a less beneficial biochemical characteristic, it may be more useful to select a plant with a decreased expression of the particular transcription factor. For a particular knockout that shows a less beneficial biochemical characteristic, it may be more useful to select a plant with an increased expression of the particular transcription factor.

#### EXAMPLE VIII. IDENTIFICATION OF HOMOLOGOUS SEQUENCES

Homologous sequences from *Arabidopsis* and plant species other than *Arabidopsis* were identified using database sequence search tools, such as the Basic Local Alignment Search Tool (BLAST) (Altschul et al. (1990) *J. Mol. Biol.* 215:403-410; and Altschul et al. (1997) *Nucl. Acid Res.* 25: 3389-3402). The tblastx sequence analysis programs were employed using the BLOSUM-62 scoring matrix (Henikoff, S. and Henikoff, J. G. (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919).

Identified *Arabidopsis* homologous sequences are provided in Figure 2 and included in the Sequence Listing. The percent sequence identity among these sequences is as low as 47% sequence identity. Additionally, the entire NCBI GenBank database was filtered for sequences from all plants except *Arabidopsis thaliana* by selecting all entries in the NCBI GenBank database associated with NCBI taxonomic ID 33090 (Viridiplantae; all plants) and excluding entries associated with taxonomic ID 3701 (*Arabidopsis thaliana*). These sequences were compared to sequences representing genes of SEQ IDs Nos. 1-54 on 9/26/2000 using the Washington University TBLASTX algorithm (version 2.0a19MP). For each gene of SEQ IDs Nos. 1-54, individual comparisons were ordered by probability score (P-value), where the score reflects the probability that a particular alignment occurred by chance. For example, a score of  $3.6e-40$  is  $3.6 \times 10^{-40}$ . For up to ten species, the gene with the lowest P-value (and therefore the most likely homolog) is listed in Figure 3.

In addition to P-values, comparisons were also scored by percentage identity. Percentage identity reflects the degree to which two segments of DNA or protein are identical over a particular length. The ranges of percent identity between the non-*Arabidopsis* genes shown in Figure 3 and the *Arabidopsis* genes in the sequence listing are: SEQ ID No. 1: 38%-89%; SEQ ID No. 3: 64%-88%; SEQ ID No. 5: 44%-84%; SEQ ID No. 7: 35%-86%; SEQ ID No. 9: 43%-77%; SEQ ID No. 11: 43%-85%; SEQ ID No. 13: 41%-76%; SEQ ID No. 15: 34%-63%; SEQ ID No. 17: 31%-68%; SEQ ID No. 19: 26%-44%; SEQ ID No. 21: 52%-70%; SEQ ID No. 23: 37%-

93%; SEQ ID No. 25: 37%-58%; SEQ ID No. 27: 48%-92%; SEQ ID No. 29: 42%-88%; SEQ ID No. 31: 47%-90%; SEQ ID No. 33: 45%-69%; SEQ ID No. 35: 42%-94%; SEQ ID No. 37: 38%-85%; SEQ ID No. 39: 49%-93%; SEQ ID No. 41: 36%-64%; and SEQ ID No. 43: 36%-70%.

5 The polynucleotides and polypeptides in the Sequence Listing and the identified homologous sequences may be stored in a computer system and have associated or linked with the sequences a function, such as that the polynucleotides and polypeptides are useful for modifying the biochemical characteristics of a plant.

10 All references, publications, patents and other documents herein are incorporated by reference in their entirety for all purposes. Although the invention has been described with reference to the embodiments and examples above, it should be understood that various modifications can be made without departing from the spirit of the invention.

What is claimed is:

1. A transgenic plant with a modified biochemical characteristic, which plant comprises a recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - 5 (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-22, or a complementary nucleotide sequence thereof;
  - (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);
  - (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-22, or a complementary nucleotide sequence thereof;
  - 10 (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
  - (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
  - (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e);
  - 15 (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide that modifies a plant's biochemical characteristic;
  - (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
  - 20 (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g);
  - (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-22;
  - 25 (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-22; and
  - (l) a nucleotide sequence which encodes a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-22.
- 30 2. The transgenic plant of claim 1, further comprising a constitutive, inducible, or tissue-active promoter operably linked to said nucleotide sequence.
3. The transgenic plant of claim 1, wherein the plant is selected from the group consisting of: soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf,

banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, and vegetable brassicas.

5

4. An isolated or recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-22, or a complementary nucleotide sequence thereof;
- 10 (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);
- (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-22, or a complementary nucleotide sequence thereof;
- (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
- 15 (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
- (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e);
- (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which
- 20 subsequence or fragment encodes a polypeptide that modifies a plant's biochemical characteristic;
- (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
- (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide
- 25 sequence of any of (a)-(g);
- (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-22;
- (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-22; and
- 30 (l) a nucleotide sequence which encodes a conserved domain of a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-22.

5. The isolated or recombinant polynucleotide of claim 4, further comprising a constitutive, inducible, or tissue-active promoter operably linked to the nucleotide sequence.
6. A cloning or expression vector comprising the isolated or recombinant polynucleotide of claim 4.
7. A cell comprising the cloning or expression vector of claim 6.
8. A transgenic plant comprising the isolated or recombinant polynucleotide of claim 4.
9. A composition produced by one or more of:
- (a) incubating one or more polynucleotide of claim 4 with a nuclease;
  - (b) incubating one or more polynucleotide of claim 4 with a restriction enzyme;
  - (c) incubating one or more polynucleotide of claim 4 with a polymerase;
  - (d) incubating one or more polynucleotide of claim 4 with a polymerase and a primer;
  - (e) incubating one or more polynucleotide of claim 4 with a cloning vector, or
  - (f) incubating one or more polynucleotide of claim 4 with a cell.
10. A composition comprising two or more different polynucleotides of claim 4.
11. An isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4.
12. A plant ectopically expressing an isolated polypeptide of claim 11.
13. A method for producing a plant having a modified biochemical characteristic, the method comprising altering the expression of the isolated or recombinant polynucleotide of claim 4 or the expression levels or activity of a polypeptide of claim 11 in a plant, thereby producing a modified plant, and selecting the modified plant for a modified biochemical characteristic thereby providing the modified plant with a modified biochemical characteristic.
14. The method of claim 13, wherein the polynucleotide is a polynucleotide of claim 4.



15. A method of identifying a factor that is modulated by or interacts with a polypeptide encoded by a polynucleotide of claim 4, the method comprising:
- (a) expressing a polypeptide encoded by the polynucleotide in a plant; and
  - (b) identifying at least one factor that is modulated by or interacts with the polypeptide.
- 5
16. The method of claim 15, wherein the identifying is performed by detecting binding by the polypeptide to a promoter sequence, or detecting interactions between an additional protein and the polypeptide in a yeast two hybrid system.
- 10 17. The method of claim 15, wherein the identifying is performed by detecting expression of a factor by hybridization to a microarray, subtractive hybridization or differential display.
18. A method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest, the method comprising:
- (a) placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of claim 4; and,
  - (b) monitoring one or more of:
    - (i) expression level of the polynucleotide in the plant;
    - (ii) expression level of the polypeptide in the plant;
    - (iii) modulation of an activity of the polypeptide in the plant; or
    - (iv) modulation of an activity of the polynucleotide in the plant.
- 15
- 20
19. An integrated system, computer or computer readable medium comprising one or more character strings corresponding to a polynucleotide of claim 4, or to a polypeptide encoded by the polynucleotide.
- 25
20. The integrated system, computer or computer readable medium of claim 19, further comprising a link between said one or more sequence strings to a modified plant biochemical characteristics phenotype.
- 30
21. A method of identifying a sequence similar or homologous to one or more polynucleotides of claim 4, or one or more polypeptides encoded by the polynucleotides, the method comprising:
- (a) providing a sequence database; and,

(b) querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.

5

22. The method of claim 21, wherein the querying comprises aligning one or more of the target sequences with one or more of the one or more sequence members in the sequence database.

10

23. The method of claim 21, wherein the querying comprises identifying one or more of the one or more sequence members of the database that meet a user-selected identity criteria with one or more of the target sequences.

15

24. The method of claim 21, further comprising linking the one or more of the polynucleotides of claim 4, or encoded polypeptides, to a modified plant biochemical characteristics phenotype.

20

25. A plant comprising altered expression levels of an isolated or recombinant polynucleotide of claim 4.

26. A plant comprising altered expression levels or the activity of an isolated or recombinant polypeptide of claim 11.

25

27. A plant lacking a nucleotide sequence encoding a polynucleotide of claim 11.

Figure 1

SEQ ID No.	GID	cDNA or protein	conserved domain
1	G214	cDNA	
2	G214	protein	22-71
3	G231	cDNA	
4	G231	protein	14-118
5	G274	cDNA	
6	G274	protein	108-572
7	G307	cDNA	
8	G307	protein	323-339
9	G346	cDNA	
10	G346	protein	196-221
11	G598	cDNA	
12	G598	protein	205-263
13	G605	cDNA	
14	G605	protein	132-143
15	G777	cDNA	
16	G777	protein	47-101
17	G869	cDNA	
18	G869	protein	109-177
19	G1133	cDNA	
20	G1133	protein	256-326
21	G1266	cDNA	
22	G1266	protein	79-147
23	G1324	cDNA	
24	G1324	protein	20-118
25	G1337	cDNA	
26	G1337	protein	9-75
27	G975	cDNA	
28	G975	protein	4-71

Figure 2

SEQ ID No.	GID	homolog	cDNA or protein	conserved domain
29	G680	homolog of G214	cDNA	
30	G680	homolog of G214	protein	24-70
31	G883	homolog of G274	cDNA	
32	G883	homolog of G274	protein	245-302
33	G1855	homolog of G274	cDNA	
34	G1855	homolog of G274	protein	entire protein
35	G1190	homolog of G274	cDNA	
36	G1190	homolog of G274	protein	entire protein
37	G308	homolog of G307	cDNA	
38	G308	homolog of G307	protein	270-274
39	G1944	homolog of G605	cDNA	
40	G1944	homolog of G605	protein	87-100
41	G326	homolog of G1337	cDNA	
42	G326	homolog of G1337	protein	11-94, 354-400
43	G1387	homolog of G975	cDNA	
44	G1387	homolog of G975	protein	4-71

Figure 3A

SEQ ID No.	GID	Genbank NID	P-value	Species
1	G214	8170933	8.80E-35	Lycopersicon esculentum
1	G214	9205339	1.20E-27	Glycine max
1	G214	8577344	1.80E-23	Zea mays
1	G214	9119112	2.40E-18	Medicago truncatula
1	G214	7660673	4.80E-15	Sorghum bicolor
1	G214	8213273	4.40E-14	Oryza sativa
1	G214	3325786	4.70E-10	Gossypium hirsutum
1	G214	9435251	1.50E-09	Hordeum vulgare
1	G214	9411569	6.80E-09	Triticum aestivum
1	G214	7614730	3.00E-07	Lotus japonicus
3	G231	6651291	7.80E-71	Pimpinella brachycarpa
3	G231	1430845	1.90E-62	Lycopersicon esculentum
3	G231	5268844	1.40E-61	Zea mays
3	G231	7561750	3.90E-60	Medicago truncatula
3	G231	1945282	3.30E-59	Oryza sativa
3	G231	22637	9.80E-49	Physcomitrella patens
3	G231	437326	2.00E-48	Gossypium hirsutum
3	G231	20562	3.40E-48	Petunia x hybrida
3	G231	4886263	5.00E-48	Antirrhinum majus
3	G231	8379692	1.50E-47	Gossypium arboreum
5	G274	6752887	1.70E-231	Malus domestica
5	G274	5734616	1.20E-140	Oryza sativa
5	G274	8996178	5.40E-96	Suaeda maritima subsp. salsa
5	G274	6654657	1.50E-89	Medicago truncatula
5	G274	8105703	2.30E-88	Lycopersicon esculentum
5	G274	7625402	4.00E-87	Gossypium arboreum
5	G274	7588836	2.10E-82	Glycine max
5	G274	5045979	1.30E-76	Gossypium hirsutum
5	G274	7324635	1.90E-71	Lycopersicon pennellii
5	G274	8903627	3.60E-63	Hordeum vulgare
7	G307	5640156	3.80E-151	Triticum aestivum
7	G307	5640154	1.00E-101	Zea mays
7	G307	6970471	1.70E-97	Oryza sativa
7	G307	7718432	4.00E-82	Medicago truncatula
7	G307	8330344	7.90E-78	Mesembryanthemum crystallinum
7	G307	5047560	1.00E-72	Gossypium hirsutum
7	G307	7588689	2.70E-69	Glycine max
7	G307	7623983	2.20E-64	Gossypium arboreum
7	G307	7780253	9.30E-59	Lotus japonicus
7	G307	6733213	1.90E-51	Lycopersicon esculentum
9	G346	4387642	5.90E-28	Lycopersicon esculentum
9	G346	7627902	1.50E-27	Gossypium arboreum
9	G346	8335147	6.40E-27	Oryza sativa
9	G346	8529362	9.10E-27	Medicago truncatula
9	G346	403305	2.30E-26	Nicotiana tabacum
9	G346	9299618	2.50E-26	Sorghum bicolor
9	G346	5056246	7.80E-26	Brassica rapa subsp. pekinensis
9	G346	6827291	6.80E-25	Zea mays
9	G346	6567406	1.90E-24	Glycine max
9	G346	9425896	1.20E-21	Triticum turgidum subsp. durum
11	G598	8102670	1.30E-43	Zea mays
11	G598	4382198	9.80E-42	Lycopersicon esculentum

Figure 3B

SEQ ID No.	GID	Genbank NID	P-value	Species
11	G598	7553316	8.00E-38	<i>Sorghum bicolor</i>
11	G598	9445834	3.10E-36	<i>Triticum aestivum</i>
11	G598	7332502	8.80E-30	<i>Oryza sativa</i>
11	G598	9056816	1.70E-17	<i>Medicago truncatula</i>
11	G598	6644720	5.20E-15	<i>Mesembryanthemum crystallinum</i>
11	G598	3853398	2.20E-14	<i>Populus tremula</i> x <i>Populus tremuloides</i>
11	G598	9419408	6.80E-09	<i>Hordeum vulgare</i>
11	G598	6848223	1.40E-06	<i>Glycine max</i>
13	G605	7624850	4.40E-49	<i>Gossypium arboreum</i>
13	G605	9204125	6.50E-46	<i>Glycine max</i>
13	G605	2213533	5.50E-33	<i>Pisum sativum</i>
13	G605	7009437	1.40E-28	<i>Zea mays</i>
13	G605	8104258	3.50E-28	<i>Lycopersicon esculentum</i>
13	G605	7536402	4.10E-28	<i>Sorghum bicolor</i>
13	G605	3107210	1.60E-22	<i>Oryza sativa</i>
13	G605	7784135	9.20E-20	<i>Lotus japonicus</i>
13	G605	4165182	8.30E-18	<i>Antirrhinum majus</i>
13	G605	6555294	8.10E-17	<i>Pinus taeda</i>
15	G777	8172576	3.10E-29	<i>Medicago truncatula</i>
15	G777	8331320	4.60E-17	<i>Mesembryanthemum crystallinum</i>
15	G777	8106138	3.00E-16	<i>Lycopersicon esculentum</i>
15	G777	5046832	1.20E-14	<i>Gossypium hirsutum</i>
15	G777	6918785	1.70E-13	<i>Zea mays</i>
15	G777	5666914	1.30E-07	<i>Glycine max</i>
15	G777	8856987	0.98	<i>Oryza sativa</i>
15	G777	8404755	1	<i>Hordeum vulgare</i>
17	G869	2213784	1.30E-19	<i>Lycopersicon esculentum</i>
17	G869	3065894	7.30E-19	<i>Nicotiana tabacum</i>
17	G869	8570080	4.20E-18	<i>Oryza sativa</i>
17	G869	7560260	1.50E-17	<i>Medicago truncatula</i>
17	G869	7534890	5.20E-14	<i>Sorghum bicolor</i>
17	G869	6455322	1.10E-13	<i>Glycine max</i>
17	G869	9362061	2.70E-13	<i>Triticum aestivum</i>
17	G869	7788764	5.70E-13	<i>Lotus japonicus</i>
17	G869	7624302	2.50E-12	<i>Gossypium arboreum</i>
17	G869	3858036	2.80E-12	<i>Populus balsamifera</i> subsp. <i>trichocarpa</i>
19	G1133	8070726	1.30E-16	<i>Solanum tuberosum</i>
19	G1133	6848196	1.60E-16	<i>Glycine max</i>
19	G1133	7570922	3.60E-13	<i>Medicago truncatula</i>
19	G1133	9434859	1.90E-12	<i>Lycopersicon esculentum</i>
19	G1133	5704484	0.005	<i>Oryza sativa</i>
19	G1133	902661	0.0081	<i>Hordeum vulgare</i>
19	G1133	8666194	0.0086	<i>Pinus taeda</i>
19	G1133	5725018	0.14	<i>Brassica rapa</i> subsp. <i>pekinensis</i>
19	G1133	7501051	0.64	<i>Gossypium arboreum</i>
19	G1133	7747388	0.98	<i>Lotus japonicus</i>
21	G1266	1732405	1.50E-50	<i>Nicotiana tabacum</i>
21	G1266	7145976	2.50E-38	<i>Glycine max</i>
21	G1266	3326366	1.00E-37	<i>Gossypium hirsutum</i>
21	G1266	5762854	6.90E-37	<i>Lotus japonicus</i>
21	G1266	7560749	9.10E-34	<i>Medicago truncatula</i>
21	G1266	7934594	6.60E-33	<i>Euphorbia esula</i>
21	G1266	9431305	2.10E-28	<i>Lycopersicon esculentum</i>

Figure 3C

SEQ ID No.	GID	Genbank NID	P-value	Species
21	G1266	7528275	5.40E-21	Mesembryanthemum crystallinum
21	G1266	6478844	4.10E-20	Matricaria chamomilla
21	G1266	7627061	4.20E-20	Gossypium arboreum
23	G1324	2921337	2.30E-54	Gossypium hirsutum
23	G1324	5891412	3.50E-52	Lycopersicon esculentum
23	G1324	8528843	7.20E-50	Medicago truncatula
23	G1324	1002797	5.40E-49	Cratostigma plantagineum
23	G1324	5666961	3.90E-44	Glycine max
23	G1324	7244640	1.70E-42	Mentha x piperita
23	G1324	1841474	3.00E-42	Pisum sativum
23	G1324	4979554	1.30E-39	Oryza sativa
23	G1324	9363368	3.00E-32	Triticum aestivum
23	G1324	9296080	3.50E-32	Sorghum bicolor
25	G1337	7410432	2.60E-41	Lycopersicon esculentum
25	G1337	3618319	1.10E-32	Oryza sativa
25	G1337	7571599	1.00E-28	Medicago truncatula
25	G1337	7685955	5.10E-27	Glycine max
25	G1337	7323708	2.60E-25	Lycopersicon hirsutum
25	G1337	4091805	1.00E-18	Malus domestica
25	G1337	6917805	4.80E-18	Lycopersicon pennellii
25	G1337	3341722	1.60E-17	Raphanus sativus
25	G1337	2303680	4.50E-17	Brassica napus
25	G1337	4557092	9.10E-17	Pinus radiata
27	G975	8103850	8.50E-46	Lycopersicon esculentum
27	G975	7590215	1.50E-45	Glycine max
27	G975	5056299	2.20E-34	Brassica rapa subsp. pekinensis
27	G975	9278522	1.80E-26	Lotus japonicus
27	G975	1128767	2.70E-18	Brassica rapa
27	G975	5859978	5.50E-18	Pinus taeda
27	G975	9427282	2.40E-15	Triticum aestivum
27	G975	19506	4.70E-14	Lupinus polyphyllus
27	G975	6799584	5.30E-14	Medicago truncatula
27	G975	7324705	1.70E-12	Lycopersicon pennellii
29	G680	9258166	5.70E-36	Glycine max
29	G680	9255178	3.00E-29	Zea mays
29	G680	5274804	1.20E-27	Lycopersicon esculentum
29	G680	4974199	3.00E-22	Oryza sativa
29	G680	3325786	2.10E-21	Gossypium hirsutum
29	G680	9119112	1.30E-18	Medicago truncatula
29	G680	7660673	3.20E-17	Sorghum bicolor
29	G680	7243970	6.10E-16	Mentha x piperita
29	G680	3858093	2.10E-10	Populus balsamifera subsp. trichocarpa
29	G680	8845091	3.70E-10	Triticum aestivum
31	G883	4760595	2.40E-84	Nicotiana tabacum
31	G883	4894962	3.50E-45	Avena sativa
31	G883	6719425	1.70E-36	Glycine max
31	G883	5273248	2.80E-35	Lycopersicon esculentum
31	G883	9302479	3.00E-34	Sorghum bicolor
31	G883	6799932	1.40E-31	Medicago truncatula
31	G883	5456433	4.30E-31	Zea mays
31	G883	8706346	1.40E-30	Hordeum vulgare
31	G883	8404566	2.70E-30	Oryza sativa
31	G883	1432055	2.00E-27	Petroselinum crispum

Figure 3D

SEQ ID No.	GID	Genbank NID	P-value	Species
33	G1855	6752887	4.80E-181	Malus domestica
33	G1855	5734616	7.60E-154	Oryza sativa
33	G1855	4384552	3.80E-80	Lycopersicon esculentum
33	G1855	8996178	1.80E-78	Suaeda maritima subsp. salsa
33	G1855	7625402	1.60E-77	Gossypium arboreum
33	G1855	8903627	3.80E-74	Hordeum vulgare
33	G1855	6654657	2.20E-70	Medicago truncatula
33	G1855	8090141	4.50E-64	Sorghum bicolor
33	G1855	9028645	6.30E-64	Zea mays
33	G1855	7588836	6.70E-62	Glycine max
35	G1190	6752887	7.00E-111	Malus domestica
35	G1190	5734616	1.20E-98	Oryza sativa
35	G1190	7569650	2.60E-92	Medicago truncatula
35	G1190	4380101	5.50E-88	Lycopersicon esculentum
35	G1190	6567183	5.20E-81	Glycine max
35	G1190	8901706	7.40E-80	Hordeum vulgare
35	G1190	8070121	1.70E-76	Solanum tuberosum
35	G1190	8666639	5.50E-75	Pinus taeda
35	G1190	8088688	3.40E-72	Sorghum bicolor
35	G1190	6020980	6.50E-67	Zea mays
37	G308	5640156	3.50E-162	Triticum aestivum
37	G308	5640154	2.30E-134	Zea mays
37	G308	6970471	4.20E-120	Oryza sativa
37	G308	7718432	8.70E-80	Medicago truncatula
37	G308	8330344	3.90E-76	Mesembryanthemum crystallinum
37	G308	5047560	1.50E-71	Gossypium hirsutum
37	G308	7588689	1.90E-68	Glycine max
37	G308	7623983	2.90E-62	Gossypium arboreum
37	G308	7780253	1.10E-57	Lotus japonicus
37	G308	6733213	3.70E-48	Lycopersicon esculentum
39	G1944	9204125	5.50E-52	Glycine max
39	G1944	7624850	6.60E-45	Gossypium arboreum
39	G1944	7784135	7.20E-32	Lotus japonicus
39	G1944	9280727	2.60E-29	Oryza sativa
39	G1944	7009437	1.30E-28	Zea mays
39	G1944	7536402	1.30E-28	Sorghum bicolor
39	G1944	8104258	6.50E-27	Lycopersicon esculentum
39	G1944	2213533	3.50E-23	Pisum sativum
39	G1944	4165182	7.10E-17	Antirrhinum majus
39	G1944	6555294	2.90E-16	Pinus taeda
41	G326	7410432	1.10E-37	Lycopersicon esculentum
41	G326	3618319	2.90E-32	Oryza sativa
41	G326	7571599	4.90E-30	Medicago truncatula
41	G326	7232283	6.30E-28	Glycine max
41	G326	7323708	6.00E-27	Lycopersicon hirsutum
41	G326	4091805	2.30E-19	Malus domestica
41	G326	6917805	6.50E-19	Lycopersicon pennellii
41	G326	3341722	2.50E-18	Raphanus sativus
41	G326	4557092	7.50E-18	Pinus radiata
41	G326	2303680	4.70E-17	Brassica napus
43	G1387	8285738	1.40E-46	Glycine max
43	G1387	8103850	5.20E-46	Lycopersicon esculentum
43	G1387	5056299	1.10E-20	Brassica rapa subsp. pekinensis



Figure 3E

SEQ ID No.	GID	Genbank NID	P-value	Species
43	G1387	9278522	1.50E-18	Lotus japonicus
43	G1387	5859978	2.00E-15	Pinus taeda
43	G1387	7766740	4.70E-14	Medicago truncatula
43	G1387	9427282	1.40E-12	Triticum aestivum
43	G1387	3857766	3.40E-12	Populus balsamifera subsp. trichocarpa
43	G1387	19506	4.60E-12	Lupinus polyphyllus
43	G1387	7273843	2.20E-11	Oryza sativa

MBI-20 Sequence Listing.ST25  
SEQUENCE LISTING

<110> Creelman, Robert  
Yu, Guo-Liang  
Adam, Luc  
Riechmann, Jose Luis  
Heard, Jacqueline  
Samaha, Raymond  
Pilgrim, Marsha  
Pineda, Omaira  
Jiang, Cai-Zhong

<120> Plant Biochemistry-Related Genes

<130> MBI-0020

<150> 60/164,132  
<151> 1999-11-17

<150> 60/197,899  
<151> 2000-04-17

<150> Plant Trait Modification III  
<151> 2000-08-22

<160> 44

<170> PatentIn version 3.0

<210> 1  
<211> 2240  
<212> DNA  
<213> Arabidopsis thaliana

<220>  
<221> CDS  
<222> (238)..(2064)  
<223> G214

<400> 1  
tgagatttct ccatttccgt agcttctggt ctcttttctt tgtttcattg atcaaaagca 60  
aatcacttct tcttcttctt ctctctgatt tcttactggt ttcttatcca acgaaatctg 120  
gaattaaaaa tggaatcttt atcgaatcca agctgatttt gtttctttca ttgaatcatc 180  
tctctaaagt ggaattttgt aaagagaaga tctgaagttg tgtagaggag cttagtg 237  
atg gag aca aat tcg tct gga gaa gat ctg gtt att aag act cgg aag 285  
Met Glu Thr Asn Ser Ser Gly Glu Asp Leu Val Ile Lys Thr Arg Lys  
1 5 10 15  
cca tat acg ata aca aag caa cgt gaa agg tgg act gag gaa gaa cat 333  
Pro Tyr Thr Ile Thr Lys Gln Arg Glu Arg Trp Thr Glu Glu Glu His  
20 25 30  
aat aga ttc att gaa gct ttg agg ctt tat ggt aga gca tgg cag aag 381  
Asn Arg Phe Ile Glu Ala Leu Arg Leu Tyr Gly Arg Ala Trp Gln Lys  
35 40 45  
att gaa gaa cat gta gca aca aaa act gct gtc cag ata aga agt cac 429  
Ile Glu Glu His Val Ala Thr Lys Thr Ala Val Gln Ile Arg Ser His  
50 55 60  
gct cag aaa ttt ttc tcc aag gta gag aaa gag gct gaa gct aaa ggt 477  
Ala Gln Lys Phe Phe Ser Lys Val Glu Lys Glu Ala Glu Ala Lys Gly  
65 70 75 80  
gta gct atg ggt caa gcg cta gac ata gct att cct cct cca cgg cct 525  
Val Ala Met Gly Gln Ala Leu Asp Ile Ala Ile Pro Pro Pro Arg Pro  
85 90 95  
aag cgt aaa cca aac aat cct tat cct cga aag acg gga agt gga acg 573  
Lys Arg Lys Pro Asn Asn Pro Tyr Pro Arg Lys Thr Gly Ser Gly Thr

## MBI-20 Sequence Listing.ST25

100	105	110	
atc ctt atg tca aaa acg ggt gtg aat gat gga aaa gag tcc ctt gga Ile Leu Met Ser Lys Thr Gly Val Asn Asp Gly Lys Glu Ser Leu Gly 115 120 125			621
tca gaa aaa gtg tcg cat cct gag atg gcc aat gaa gat cga caa caa Ser Glu Lys Val Ser His Pro Glu Met Ala Asn Glu Asp Arg Gln Gln 130 135 140			669
tca aag cct gaa gag aaa act ctg cag gaa gac aac tgt tca gat tgt Ser Lys Pro Glu Glu Lys Thr Leu Gln Glu Asp Asn Cys Ser Asp Cys 145 150 155 160			717
ttc act cat cag tat ctc tct gct gca tcc tcc atg aat aaa agt tgt Phe Thr His Gln Tyr Leu Ser Ala Ala Ser Ser Met Asn Lys Ser Cys 165 170 175			765
ata gag aca tca aac gca agc act ttc cgc gag ttc ttg cct tca cgg Ile Glu Thr Ser Asn Ala Ser Thr Phe Arg Glu Phe Leu Pro Ser Arg 180 185 190			813
gaa gag gga agt cag aat aac agg gta aga aag gag tca aac tca gat Glu Glu Gly Ser Gln Asn Asn Arg Val Arg Lys Glu Ser Asn Ser Asp 195 200 205			861
ttg aat gca aaa tct ctg gaa aac ggt aat gag caa gga cct cag act Leu Asn Ala Lys Ser Leu Glu Asn Gly Asn Glu Gln Gly Pro Gln Thr 210 215 220			909
tat ccg atg cat atc cct gtg cta gtg cca ttg ggg agc tca ata aca Tyr Pro Met His Ile Pro Val Leu Val Pro Leu Gly Ser Ser Ile Thr 225 230 235 240			957
agt tct cta tca cat cct cct tca gag cca gat agt cat ccc cac aca Ser Ser Leu Ser His Pro Pro Ser Glu Pro Asp Ser His Pro His Thr 245 250 255			1005
gtt gca gga gat tat cag tcg ttt cct aat cat ata atg tca acc ctt Val Ala Gly Asp Tyr Gln Ser Phe Pro Asn His Ile Met Ser Thr Leu 260 265 270			1053
tta caa aca ccg gct ctt tat act gcc gca act ttc gcc tca tca ttt Leu Gln Thr Pro Ala Leu Tyr Thr Ala Ala Thr Phe Ala Ser Ser Phe 275 280 285			1101
tgg cct ccc gat tct agt ggt ggc tca cct gtt cca ggg aac tca cct Trp Pro Pro Asp Ser Ser Ser Gly Ser Pro Val Pro Gly Asn Ser Pro 290 295 300			1149
ccg aat ctg gct gcc atg gcc gca gcc act gtt gca gct gct agt gct Pro Asn Leu Ala Ala Met Ala Ala Ala Thr Val Ala Ala Ala Ser Ala 305 310 315 320			1197
tgg tgg gct gcc aat gga tta tta cct tta tgt gct cct ctt agt tca Trp Trp Ala Ala Asn Gly Leu Leu Pro Leu Cys Ala Pro Leu Ser Ser 325 330 335			1245
ggt ggt ttc act agt cat cct cca tct act ttt gga cca tca tgt gat Gly Gly Phe Thr Ser His Pro Pro Ser Thr Phe Gly Pro Ser Cys Asp 340 345 350			1293
gta gag tac aca aaa gca agc act tta caa cat ggt tct gtg cag agc Val Glu Tyr Thr Lys Ala Ser Thr Leu Gln His Gly Ser Val Gln Ser 355 360 365			1341
cga gag caa gaa cac tcc gag gca tca aag gct cga tct tca ctg gac Arg Glu Gln Glu His Ser Glu Ala Ser Lys Ala Arg Ser Ser Leu Asp 370 375 380			1389
tca gag gat gtt gaa aat aag agt aaa cca gtt tgt cat gag cag cct Ser Glu Asp Val Glu Asn Lys Ser Lys Pro Val Cys His Glu Gln Pro 385 390 395 400			1437
tct gca aca cct gag agt gat gca aag ggt tca gat gga gca gga gac			1485

## MBI-20 Sequence Listing.ST25

Ser Ala Thr Pro Glu Ser Asp Ala Lys Gly Ser Asp Gly Ala Gly Asp  
 405 410 415  
 aga aaa caa gtt gac cgg tcc tcg tgt ggc tca aac act ccg tcg agt 1533  
 Arg Lys Gln Val Asp Arg Ser Ser Cys Gly Ser Asn Thr Pro Ser Ser  
 420 425 430  
 agt gat gat gtt gag gcg gat gca tca gaa agg caa gag gat ggc acc 1581  
 Ser Asp Asp Val Glu Ala Asp Ala Ser Glu Arg Gln Glu Asp Gly Thr  
 435 440 445  
 aat ggt gag gtg aaa gaa acg aat gaa gac act aat aaa cct caa act 1629  
 Asn Gly Glu Val Lys Glu Thr Asn Glu Asp Thr Asn Lys Pro Gln Thr  
 450 455 460  
 tca gag tcc aat gca cgc cgc agt aga atc agc tcc aat ata acc gat 1677  
 Ser Glu Ser Asn Ala Arg Arg Ser Arg Ile Ser Ser Asn Ile Thr Asp  
 465 470 475 480  
 cca tgg aag tct gtg tct gac gag ggt cga att gcc ttc caa gct ctc 1725  
 Pro Trp Lys Ser Val Ser Asp Glu Gly Arg Ile Ala Phe Gln Ala Leu  
 485 490 495  
 ttc tcc aga gag gta ttg ccg caa agt ttt aca tat cga gaa gaa cac 1773  
 Phe Ser Arg Glu Val Leu Pro Gln Ser Phe Thr Tyr Arg Glu Glu His  
 500 505 510  
 aga gag gaa gaa caa caa caa gaa caa aga tat cca atg gca ctt 1821  
 Arg Glu Glu Glu Gln Gln Gln Glu Glu Gln Arg Tyr Pro Met Ala Leu  
 515 520 525  
 gat ctt aac ttc aca gct cag tta aca cca gtt gat gat caa gag gag 1869  
 Asp Leu Asn Phe Thr Ala Gln Leu Thr Pro Val Asp Asp Gln Glu Glu  
 530 535 540  
 aag aga aac aca gga ttt ctt gga atc gga tta gat gct tca aag cta 1917  
 Lys Arg Asn Thr Gly Phe Leu Gly Ile Gly Leu Asp Ala Ser Lys Leu  
 545 550 555 560  
 atg agt aga gga aga aca ggt ttt aaa cca tac aaa aga tgt tcc atg 1965  
 Met Ser Arg Gly Arg Thr Gly Phe Lys Pro Tyr Lys Arg Cys Ser Met  
 565 570 575  
 gaa gcc aaa gaa agt aga atc ctc aac aac aat cct atc att cat gtg 2013  
 Glu Ala Lys Glu Ser Arg Ile Leu Asn Asn Pro Ile Ile His Val  
 580 585 590  
 gaa cag aaa gat ccc aaa cgg atg cgg ttg gaa act caa gct tcc aca 2061  
 Glu Gln Lys Asp Pro Lys Arg Met Arg Leu Glu Thr Gln Ala Ser Thr  
 595 600 605  
 tga gactctatctt tcactctgac tgttgtttgt actctgtttt taagtgtttca 2114  
 agaccactgc tacatcttct ttttcttttg aggcctttgt atttgtttcc ttgtccatag 2174  
 tcttctgtga acatttgact ctgtattatt caacaaatca taaactgttt aatctttttt 2234  
 tttcca 2240  
 <210> 2  
 <211> 608  
 <212> PRT  
 <213> Arabidopsis thaliana  
 <400> 2  
 Met Glu Thr Asn Ser Ser Gly Glu Asp Leu Val Ile Lys Thr Arg Lys  
 1 5 10 15  
 Pro Tyr Thr Ile Thr Lys Gln Arg Glu Arg Trp Thr Glu Glu Glu His  
 20 25 30

## MBI-20 Sequence Listing.ST25

Asn Arg Phe Ile Glu Ala Leu Arg Leu Tyr Gly Arg Ala Trp Gln Lys  
 35 40 45

Ile Glu Glu His Val Ala Thr Lys Thr Ala Val Gln Ile Arg Ser His  
 50 55 60

Ala Gln Lys Phe Phe Ser Lys Val Glu Lys Glu Ala Glu Ala Lys Gly  
 65 70 75 80

Val Ala Met Gly Gln Ala Leu Asp Ile Ala Ile Pro Pro Pro Arg Pro  
 85 90 95

Lys Arg Lys Pro Asn Asn Pro Tyr Pro Arg Lys Thr Gly Ser Gly Thr  
 100 105 110

Ile Leu Met Ser Lys Thr Gly Val Asn Asp Gly Lys Glu Ser Leu Gly  
 115 120 125

Ser Glu Lys Val Ser His Pro Glu Met Ala Asn Glu Asp Arg Gln Gln  
 130 135 140

Ser Lys Pro Glu Glu Lys Thr Leu Gln Glu Asp Asn Cys Ser Asp Cys  
 145 150 155 160

Phe Thr His Gln Tyr Leu Ser Ala Ala Ser Ser Met Asn Lys Ser Cys  
 165 170 175

Ile Glu Thr Ser Asn Ala Ser Thr Phe Arg Glu Phe Leu Pro Ser Arg  
 180 185 190

Glu Glu Gly Ser Gln Asn Asn Arg Val Arg Lys Glu Ser Asn Ser Asp  
 195 200 205

Leu Asn Ala Lys Ser Leu Glu Asn Gly Asn Glu Gln Gly Pro Gln Thr  
 210 215 220

Tyr Pro Met His Ile Pro Val Leu Val Pro Leu Gly Ser Ser Ile Thr  
 225 230 235 240

Ser Ser Leu Ser His Pro Pro Ser Glu Pro Asp Ser His Pro His Thr  
 245 250 255

Val Ala Gly Asp Tyr Gln Ser Phe Pro Asn His Ile Met Ser Thr Leu  
 260 265 270

Leu Gln Thr Pro Ala Leu Tyr Thr Ala Ala Thr Phe Ala Ser Ser Phe  
 275 280 285

Trp Pro Pro Asp Ser Ser Gly Gly Ser Pro Val Pro Gly Asn Ser Pro  
 290 295 300

Pro Asn Leu Ala Ala Met Ala Ala Ala Thr Val Ala Ala Ala Ser Ala  
 305 310 315 320

Trp Trp Ala Ala Asn Gly Leu Leu Pro Leu Cys Ala Pro Leu Ser Ser  
 325 330 335

## MBI-20 Sequence Listing.ST25

Gly Gly Phe Thr Ser His Pro Pro Ser Thr Phe Gly Pro Ser Cys Asp  
 340 345 350  
 Val Glu Tyr Thr Lys Ala Ser Thr Leu Gln His Gly Ser Val Gln Ser  
 355 360 365  
 Arg Glu Gln Glu His Ser Glu Ala Ser Lys Ala Arg Ser Ser Leu Asp  
 370 375 380  
 Ser Glu Asp Val Glu Asn Lys Ser Lys Pro Val Cys His Glu Gln Pro  
 385 390 395 400  
 Ser Ala Thr Pro Glu Ser Asp Ala Lys Gly Ser Asp Gly Ala Gly Asp  
 405 410 415  
 Arg Lys Gln Val Asp Arg Ser Ser Cys Gly Ser Asn Thr Pro Ser Ser  
 420 425 430  
 Ser Asp Asp Val Glu Ala Asp Ala Ser Glu Arg Gln Glu Asp Gly Thr  
 435 440 445  
 Asn Gly Glu Val Lys Glu Thr Asn Glu Asp Thr Asn Lys Pro Gln Thr  
 450 455 460  
 Ser Glu Ser Asn Ala Arg Arg Ser Arg Ile Ser Ser Asn Ile Thr Asp  
 465 470 475 480  
 Pro Trp Lys Ser Val Ser Asp Glu Gly Arg Ile Ala Phe Gln Ala Leu  
 485 490 495  
 Phe Ser Arg Glu Val Leu Pro Gln Ser Phe Thr Tyr Arg Glu Glu His  
 500 505 510  
 Arg Glu Glu Glu Gln Gln Gln Gln Glu Gln Arg Tyr Pro Met Ala Leu  
 515 520 525  
 Asp Leu Asn Phe Thr Ala Gln Leu Thr Pro Val Asp Asp Gln Glu Glu  
 530 535 540  
 Lys Arg Asn Thr Gly Phe Leu Gly Ile Gly Leu Asp Ala Ser Lys Leu  
 545 550 555 560  
 Met Ser Arg Gly Arg Thr Gly Phe Lys Pro Tyr Lys Arg Cys Ser Met  
 565 570 575  
 Glu Ala Lys Glu Ser Arg Ile Leu Asn Asn Asn Pro Ile Ile His Val  
 580 585 590  
 Glu Gln Lys Asp Pro Lys Arg Met Arg Leu Glu Thr Gln Ala Ser Thr  
 595 600 605

<210> 3  
 <211> 916  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>

```
<221> CDS
<222> (88) .. (888)
<223> G231
```

[illegible]

## MBI-20 Sequence Listing.ST25

ttt tga ctgttcacca ttgacatatt ggcaacgc  
Phe

916

<210> 4  
<211> 266  
<212> PRT  
<213> Arabidopsis thaliana

<400> 4

Met Gly Arg Gln Pro Cys Cys Asp Lys Leu Gly Val Lys Lys Gly Pro  
1 5 10 15

Trp Thr Val Glu Glu Asp Lys Lys Leu Ile Asn Phe Ile Leu Thr Asn  
20 25 30

Gly His Cys Cys Trp Arg Ala Leu Pro Lys Leu Ala Gly Leu Arg Arg  
35 40 45

Cys Gly Lys Ser Cys Arg Leu Arg Trp Thr Asn Tyr Leu Arg Pro Gly  
50 55 60

Leu Lys Arg Gly Leu Leu Ser His Asp Glu Glu Gln Leu Val Ile Asp  
65 70 75 80

Leu His Ala Asn Leu Gly Asn Lys Trp Ser Lys Ile Ala Ser Arg Leu  
85 90 95

Pro Gly Arg Thr Asp Asn Glu Ile Lys Asn His Trp Asn Thr His Ile  
100 105 110

Lys Lys Lys Leu Leu Lys Met Gly Ile Asp Pro Met Thr His Gln Pro  
115 120 125

Leu Asn Gln Glu Pro Ser Asn Ile Asp Asn Ser Lys Thr Ile Pro Ser  
130 135 140

Asn Pro Asp Asp Val Ser Val Glu Pro Lys Thr Thr Asn Thr Lys Tyr  
145 150 155 160

Val Glu Ile Ser Val Thr Thr Thr Glu Glu Glu Ser Ser Ser Thr Val  
165 170 175

Thr Asp Gln Asn Ser Ser Met Asp Asn Glu Asn His Leu Ile Asp Asn  
180 185 190

Ile Tyr Asp Asp Asp Glu Leu Phe Ser Tyr Leu Trp Ser Asp Glu Thr  
195 200 205

Thr Lys Asp Glu Ala Ser Trp Ser Asp Ser Asn Phe Gly Val Gly Gly  
210 215 220

Thr Leu Tyr Asp His Asn Ile Ser Gly Ala Asp Ala Asp Phe Pro Ile  
225 230 235 240

Trp Ser Pro Glu Arg Ile Asn Asp Glu Lys Met Phe Leu Asp Tyr Cys  
245 250 255



## MBI-20 Sequence Listing.ST25

Gln Asp Phe Gly Val His Asp Phe Gly Phe  
 260 265

<210> 5  
 <211> 2371  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (172)..(2037)

<400> 5  
 gacattatatt taagtgtgtt ctctctctgt cacactcaca aagctttata ctttctggct 60  
 actgcaagct catcagttaa aagagcttaa accagagaga tctgataaga gaaatttttag 120  
 agtctctctg cttcaacaag atctacatcg accaggagat tagaaagaat c atg ggt 177  
 Met Gly  
 1  
 tct aag cat aac cca cca ggg aat aac aga tcg aga agt aca cta tct 225  
 Ser Lys His Asn Pro Pro Gly Asn Asn Arg Ser Arg Ser Thr Leu Ser  
 5 10 15  
 cta ctc gtt gtg gtt ggt tta tgt tgt ttc ttc tat ctt ctt gga gca 273  
 Leu Leu Val Val Val Gly Leu Cys Cys Phe Phe Tyr Leu Leu Gly Ala  
 20 25 30  
 tgg caa aag agt ggg ttt ggt aaa gga gat agc ata gct atg gag att 321  
 Trp Gln Lys Ser Gly Phe Gly Lys Gly Asp Ser Ile Ala Met Glu Ile  
 35 40 45 50  
 aca aag caa gcg cag tgt act gac att gtc act gat ctt gat ttt gaa 369  
 Thr Lys Gln Ala Gln Cys Thr Asp Ile Val Thr Asp Leu Asp Phe Glu  
 55 60 65  
 cct cat cac aac aca gtg aag atc cca cat aaa gct gat ccc aaa cct 417  
 Pro His His Thr Val Lys Ile Pro His Lys Ala Asp Pro Lys Pro  
 70 75 80  
 gtt tct ttc aaa ccg tgt gat gtg aag ctc aag gat tac acg cct tgt 465  
 Val Ser Phe Lys Pro Cys Asp Val Lys Leu Lys Asp Tyr Thr Pro Cys  
 85 90 95  
 caa gag caa gac cga gct atg aag ttc ccg aga gag aac atg att tac 513  
 Gln Glu Gln Asp Arg Ala Met Lys Phe Pro Arg Glu Asn Met Ile Tyr  
 100 105 110  
 aga gag aga cat tgt cct cct gat aat gag aag ctg cgt tgt ctt gtt 561  
 Arg Glu Arg His Cys Pro Pro Asp Asn Glu Lys Leu Arg Cys Leu Val  
 115 120 125 130  
 cca gct cct aaa ggg tat atg act cct ttc cct tgg cct aaa agc aga 609  
 Pro Ala Pro Lys Gly Tyr Met Thr Pro Phe Pro Trp Pro Lys Ser Arg  
 135 140 145  
 gat tat gtt cac tat gct aat gct cct ttc aag agc ttg act gtc gaa 657  
 Asp Tyr Val His Tyr Ala Asn Ala Pro Phe Lys Ser Leu Thr Val Glu  
 150 155 160  
 aaa gct gga cag aat tgg gtt cag ttt caa ggg aat gtg ttt aaa ttc 705  
 Lys Ala Gly Gln Asn Trp Val Gln Phe Gln Gly Asn Val Phe Lys Phe  
 165 170 175  
 cct ggt gga gga act atg ttt cct caa ggt gct gat gcg tat att gaa 753  
 Pro Gly Gly Gly Thr Met Phe Pro Gln Gly Ala Asp Ala Tyr Ile Glu  
 180 185 190  
 gag cta gct tct gtt atc cct atc aaa gat ggc tct gtt aga acc gca 801  
 Glu Leu Ala Ser Val Ile Pro Ile Lys Asp Gly Ser Val Arg Thr Ala  
 195 200 205 210

## MBI-20 Sequence Listing.ST25

ttg gac act gga tgt ggg gtt gct agt tgg ggt gct tat atg ctt aag Leu Asp Thr Gly Cys Gly Val Ala Ser Trp Gly Ala Tyr Met Leu Lys 215 220 225	849
agg aat gtt ttg act atg tcg ttt gcg cca agg gat aac cac gaa gca Arg Asn Val Leu Thr Met Ser Phe Ala Pro Arg Asp Asn His Glu Ala 230 235 240	897
caa gtc cag ttt gcg ctt gag aga ggt gtt cca gcg att atc gct gtt Gln Val Gln Phe Ala Leu Glu Arg Gly Val Pro Ala Ile Ile Ala Val 245 250 255	945
ctt gga tca atc ctt ctt cct tac cct gca aga gcc ttt gac atg gct Leu Gly Ser Ile Leu Leu Pro Tyr Pro Ala Arg Ala Phe Asp Met Ala 260 265 270	993
caa tgc tct cga tgc ttg ata cca tgg acc gca aac gag gga aca tac Gln Cys Ser Arg Cys Leu Ile Pro Trp Thr Ala Asn Glu Gly Thr Tyr 275 280 285 290	1041
tta atg gaa gta gat aga gtc ttg aga cct gga ggt tac tgg gtc tta Leu Met Glu Val Asp Arg Val Leu Arg Pro Gly Gly Tyr Trp Val Leu 295 300 305	1089
tcg ggt cct cca atc aac tgg aag aca tgg cac aag acg tgg aac cga Ser Gly Pro Pro Ile Asn Trp Lys Thr Trp His Lys Thr Trp Asn Arg 310 315 320	1137
act aaa gca gag cta aat gcc gag caa aag aga ata gag gga atc gca Thr Lys Ala Glu Leu Asn Ala Glu Gln Lys Arg Ile Glu Gly Ile Ala 325 330 335	1185
gag tcc tta tgc tgg gag aag aag tat gag aag gga gac att gca att Glu Ser Leu Cys Trp Glu Lys Lys Tyr Glu Lys Gly Asp Ile Ala Ile 340 345 350	1233
ttc aga aag aaa ata aac gat aga tca tgc gat aga tca aca ccg gtt Phe Arg Lys Lys Ile Asn Asp Arg Ser Cys Asp Arg Ser Thr Pro Val 355 360 365 370	1281
gac acc tgc aaa aga aag gac act gac gat gtc tgg tac aag gag ata Asp Thr Cys Lys Arg Lys Asp Thr Asp Asp Val Trp Tyr Lys Glu Ile 375 380 385	1329
gaa acg tgt gta aca cca ttc cct aaa gta tca aac gaa gaa gaa gtt Glu Thr Cys Val Thr Pro Phe Pro Lys Val Ser Asn Glu Glu Glu Val 390 395 400	1377
gct gga gga aag cta aag aag ttc ccc gag agg cta ttc gca gtg cct Ala Gly Gly Lys Leu Lys Lys Phe Pro Glu Arg Leu Phe Ala Val Pro 405 410 415	1425
cca agt atc tct aaa ggt ttg att aat ggc gtc gac gag gaa tca tac Pro Ser Ile Ser Lys Gly Leu Ile Asn Gly Val Asp Glu Glu Ser Tyr 420 425 430	1473
caa gaa gac atc aat cta tgg aag aag cga gtg acc gga tac aag aga Gln Glu Asp Ile Asn Leu Trp Lys Lys Arg Val Thr Gly Tyr Lys Arg 435 440 445 450	1521
att aac aga ctg ata ggt tcc acc aga tac cgt aat gtg atg gat atg Ile Asn Arg Leu Ile Gly Ser Thr Arg Tyr Arg Asn Val Met Asp Met 455 460 465	1569
aac gcc ggt ctt ggt gga ttc gct gct gcg ctt gaa tcg cct aaa tcg Asn Ala Gly Leu Gly Gly Phe Ala Ala Leu Glu Ser Pro Lys Ser 470 475 480	1617
tgg gtt atg aat gtg att cca acc att aac aag aac aca ttg agt gtt Trp Val Met Asn Val Ile Pro Thr Ile Asn Lys Asn Thr Leu Ser Val 485 490 495	1665
gtt tat gag aga ggt ctc att ggt atc tat cat gac tgg tgt gaa ggc Val Tyr Glu Arg Gly Leu Ile Gly Ile Tyr His Asp Trp Cys Glu Gly 500 505 510 515 520	1713

MBI-20 Sequence Listing.ST25

500	505	510	
ttt tca act tat cca aga aca tac gat ttc att cac gct agt ggt gtc			1761
Phe Ser Thr Tyr Pro Arg Thr Tyr Asp Phe Ile His Ala Ser Gly Val			
515	520	525	530
ttc agc ttg tat cag cac agc tgc aaa ctt gag gat att ctt ctt gaa			1809
Phe Ser Leu Tyr Gln His Ser Cys Lys Leu Glu Asp Ile Leu Leu Glu			
	535	540	545
act gat cgg att tta cga ccg gaa ggg att gtg att ttc cgg gat gag			1857
Thr Asp Arg Ile Leu Arg Pro Glu Gly Ile Val Ile Phe Arg Asp Glu			
	550	555	560
gtt gat gtt ttg aat gat gtg agg aag atc gtt gat gga atg aga tgg			1905
Val Asp Val Leu Asn Asp Val Arg Lys Ile Val Asp Gly Met Arg Trp			
	565	570	575
gat act aag tta atg gat cat gaa gac ggt cct ctc gtg ccg gag aag			1953
Asp Thr Lys Leu Met Asp His Glu Asp Gly Pro Leu Val Pro Glu Lys			
	580	585	590
att ctt gtc gcc acg aag cag tat tgg gta gcc ggc gac gat gga aac			2001
Ile Leu Val Ala Thr Lys Gln Tyr Trp Val Ala Gly Asp Asp Gly Asn			
	595	600	610
aat tct ccg tcg tct tct aat agt gaa gaa gaa taa aacaaaaaca			2047
Asn Ser Pro Ser Ser Ser Asn Ser Glu Glu Glu			
	615	620	
aaaaactcct caggttacta agcttgaagt gtagatctat tttacaacat ctggaaaatt			2107
cttatcaaaa aaggaaggaa tcagaatttc cattaaagaa aggtgtcaaa aaaaagttgt			2167
aaaactatat agtagtgatc aagacgaata tgtgcattta tgttttattt ttgttcccta			2227
gtttttaatt ttattttttt gaaggaagaa aaaattagtt ccatgtgttt ttgcaagata			2287
gttgaaacct tggacgcttg ttatgtatga tgcatcttg acatttttta ataacagtta			2347
ttttaaataa atttatgata taaa			2371
<210> 6			
<211> 621			
<212> PRT			
<213> Arabidopsis thaliana			
<400> 6			
Met Gly Ser Lys His Asn Pro Pro Gly Asn Asn Arg Ser Arg Ser Thr			
1	5	10	15
Leu Ser Leu Leu Val Val Val Gly Leu Cys Cys Phe Phe Tyr Leu Leu			
	20	25	30
Gly Ala Trp Gln Lys Ser Gly Phe Gly Lys Gly Asp Ser Ile Ala Met			
	35	40	45
Glu Ile Thr Lys Gln Ala Gln Cys Thr Asp Ile Val Thr Asp Leu Asp			
	50	55	60
Phe Glu Pro His His Asn Thr Val Lys Ile Pro His Lys Ala Asp Pro			
	65	70	75
Lys Pro Val Ser Phe Lys Pro Cys Asp Val Lys Leu Lys Asp Tyr Thr			
	85	90	95
Pro Cys Gln Glu Gln Asp Arg Ala Met Lys Phe Pro Arg Glu Asn Met			

MBI-20 Sequence Listing.ST25  
 100                      105                      110

```

Ile Tyr Arg Glu Arg His Cys Pro Pro Asp Asn Glu Lys Leu Arg Cys
  115                      120                      125

Leu Val Pro Ala Pro Lys Gly Tyr Met Thr Pro Phe Pro Trp Pro Lys
  130                      135                      140

Ser Arg Asp Tyr Val His Tyr Ala Asn Ala Pro Phe Lys Ser Leu Thr
  145                      150                      155                      160

Val Glu Lys Ala Gly Gln Asn Trp Val Gln Phe Gln Gly Asn Val Phe
                165                      170                      175

Lys Phe Pro Gly Gly Gly Thr Met Phe Pro Gln Gly Ala Asp Ala Tyr
                180                      185                      190

Ile Glu Glu Leu Ala Ser Val Ile Pro Ile Lys Asp Gly Ser Val Arg
  195                      200                      205

Thr Ala Leu Asp Thr Gly Cys Gly Val Ala Ser Trp Gly Ala Tyr Met
  210                      215                      220

Leu Lys Arg Asn Val Leu Thr Met Ser Phe Ala Pro Arg Asp Asn His
  225                      230                      235                      240

Glu Ala Gln Val Gln Phe Ala Leu Glu Arg Gly Val Pro Ala Ile Ile
                245                      250                      255

Ala Val Leu Gly Ser Ile Leu Leu Pro Tyr Pro Ala Arg Ala Phe Asp
  260                      265                      270

Met Ala Gln Cys Ser Arg Cys Leu Ile Pro Trp Thr Ala Asn Glu Gly
  275                      280                      285

Thr Tyr Leu Met Glu Val Asp Arg Val Leu Arg Pro Gly Gly Tyr Trp
  290                      295                      300

Val Leu Ser Gly Pro Pro Ile Asn Trp Lys Thr Trp His Lys Thr Trp
  305                      310                      315                      320

Asn Arg Thr Lys Ala Glu Leu Asn Ala Glu Gln Lys Arg Ile Glu Gly
                325                      330                      335

Ile Ala Glu Ser Leu Cys Trp Glu Lys Lys Tyr Glu Lys Gly Asp Ile
  340                      345                      350

Ala Ile Phe Arg Lys Lys Ile Asn Asp Arg Ser Cys Asp Arg Ser Thr
  355                      360                      365

Pro Val Asp Thr Cys Lys Arg Lys Asp Thr Asp Asp Val Trp Tyr Lys
  370                      375                      380

Glu Ile Glu Thr Cys Val Thr Pro Phe Pro Lys Val Ser Asn Glu Glu
  385                      390                      395                      400

```

## MBI-20 Sequence Listing.ST25

Glu Val Ala Gly Gly Lys Leu Lys Lys Phe Pro Glu Arg Leu Phe Ala  
405 410 415

Val Pro Pro Ser Ile Ser Lys Gly Leu Ile Asn Gly Val Asp Glu Glu  
420 425 430

Ser Tyr Gln Glu Asp Ile Asn Leu Trp Lys Lys Arg Val Thr Gly Tyr  
435 440 445

Lys Arg Ile Asn Arg Leu Ile Gly Ser Thr Arg Tyr Arg Asn Val Met  
450 455 460

Asp Met Asn Ala Gly Leu Gly Gly Phe Ala Ala Ala Leu Glu Ser Pro  
465 470 475 480

Lys Ser Trp Val Met Asn Val Ile Pro Thr Ile Asn Lys Asn Thr Leu  
485 490 495

Ser Val Val Tyr Glu Arg Gly Leu Ile Gly Ile Tyr His Asp Trp Cys  
500 505 510

Glu Gly Phe Ser Thr Tyr Pro Arg Thr Tyr Asp Phe Ile His Ala Ser  
515 520 525

Gly Val Phe Ser Leu Tyr Gln His Ser Cys Lys Leu Glu Asp Ile Leu  
530 535 540

Leu Glu Thr Asp Arg Ile Leu Arg Pro Glu Gly Ile Val Ile Phe Arg  
545 550 555 560

Asp Glu Val Asp Val Leu Asn Asp Val Arg Lys Ile Val Asp Gly Met  
565 570 575

Arg Trp Asp Thr Lys Leu Met Asp His Glu Asp Gly Pro Leu Val Pro  
580 585 590

Glu Lys Ile Leu Val Ala Thr Lys Gln Tyr Trp Val Ala Gly Asp Asp  
595 600 605

Gly Asn Asn Ser Pro Ser Ser Ser Asn Ser Glu Glu Glu  
610 615 620

<210> 7  
<211> 1764  
<212> DNA  
<213> Arabidopsis thaliana

<220>  
<221> CDS  
<222> (1)..(1764)  
<223> G307

<400> 7  
atg aag aga gat cat cac caa ttc caa ggt cga ttg tcc aac cac ggg 48  
Met Lys Arg Asp His His Gln Phe Gln Gly Arg Leu Ser Asn His Gly  
1 5 10 15

act tct tct tct tca tca tca atc tct aaa gat aag atg atg atg gtg 96  
Thr Ser Ser Ser Ser Ser Ser Ile Ser Lys Asp Lys Met Met Met Val  
20 25 30

## MBI-20 Sequence Listing.ST25

aaa aaa gaa gaa gac ggt gga ggt aac atg gac gac gag ctt ctc gct Lys Lys   Glu Glu Asp Gly Gly   Gly Asn Met Asp Asp   Glu Leu Leu Ala 35                  40                  45	144
ggt tta ggt tac aaa gtt agg tca tct gag atg gcg gag gtt gct ttg Val Leu   Gly Tyr Lys Val Arg Ser Ser Glu Met Ala Glu Val Ala Leu 50                  55                  60	192
aaa ctc gaa caa tta gag acg atg atg agt aat gtt caa gaa gat ggt Lys Leu Glu Gln Leu Glu Thr Met Met Ser Asn Val Gln Glu Asp Gly 65                  70                  75                  80	240
tta tct cat ctc gcg acg gat act gtt cat tat aat ccg tct gag ctt Leu Ser His Leu Ala Thr Asp Thr Val His Tyr Asn Pro Ser Glu Leu 85                  90                  95	288
tat tct tgg ctt gat aat atg ctc tct gag ctt aat cct cct cct ctt Tyr Ser Trp   Leu Asp Asn Met Leu Ser Glu Leu Asn Pro Pro Pro Leu 100                  105                  110	336
ccg gcg agt tct aac ggt tta gat ccg gtt ctt cct tct ccg gag att Pro Ala Ser Ser Asn Gly Leu Asp Pro Val Leu Pro Ser Pro Glu Ile 115                  120                  125	384
tgt ggt ttt ccg gct tct gat tat gac ctt aaa gtc att ccc gga aac Cys Gly Phe Pro Ala Ser Asp Tyr Asp Leu Lys Val Ile Pro Gly Asn 130                  135                  140	432
gcg att tat cag ttt ccg gcg att gat tct tct tct tct tct aat aat Ala Ile Tyr Gln Phe Pro Ala Ile Asp Ser Ser Ser Ser Ser Asn Asn 145                  150                  155                  160	480
cag aac aag cgt ttg aaa tca tgc tct agt cct gat tct atg gtt aca Gln Asn Lys Arg Leu Lys Ser Cys Ser Ser Pro Asp Ser Met Val Thr 165                  170                  175	528
tct act tct acg ggt acg cag att ggt gga gtc ata gga acg acg gtg Ser Thr Ser Thr Gly Thr Gln Ile Gly Gly Val Ile Gly Thr Thr Val 180                  185                  190	576
acg aca acc acc acg aca acg acg gcg gcg gct gag tca act cgt tct Thr Thr Thr Thr Thr Thr Thr Thr Ala Ala Ala Glu Ser Thr Arg Ser 195                  200                  205	624
ggt atc ctg gtt gac tct caa gag aac ggt gtt cgt tta gtc cac gcg Val Ile Leu Val Asp Ser Gln Glu Asn Gly Val Arg Leu Val His Ala 210                  215                  220	672
ctt atg gct tgt gca gaa gca atc cag cag aac aat ttg act cta gcg Leu Met Ala Cys Ala Glu Ala Ile Gln Gln Asn Asn Leu Thr Leu Ala 225                  230                  235                  240	720
gaa gct ctt gtg aag caa atc gga tgc tta gct gtg tct caa gcc gga Glu Ala Leu Val Lys Gln Ile Gly Cys Leu Ala Val Ser Gln Ala Gly 245                  250                  255	768
gct atg aga aaa gtg gct act tac ttc gcc gaa gct tta gct cgg cgg Ala Met Arg Lys Val Ala Thr Tyr Phe Ala Glu Ala Leu Ala Arg Arg 260                  265                  270	816
atc tac cgt ctc tct ccg ccg cag aat cag atc gat cat tgt ctc tcc Ile Tyr Arg Leu Ser Pro Pro Gln Asn Gln Ile Asp His Cys Leu Ser 275                  280                  285	864
gat act ctt cag atg cac ttt tac gag act tgt cct tat ctt aaa ttc Asp Thr Leu Gln Met His Phe Tyr Glu Thr Cys Pro Tyr Leu Lys Phe 290                  295                  300	912
gct cac ttc acg gcg aac caa gcg att ctc gaa gct ttt gaa ggt aag Ala His Phe Thr Ala Asn Gln Ala Ile Leu Glu Ala Phe Glu Gly Lys 305                  310                  315                  320	960
aag aga gta cac gtc att gat ttc tct atg aac caa ggt ctt caa tgg Lys Arg Val His Val Ile Asp Phe Ser Met Asn Gln Gly Leu Gln Trp	1008

## MBI-20 Sequence Listing.ST25

325	330	335	
cct gcg ctt atg caa gct ctt gcg ctt cga gaa gga ggt cct cca act Pro Ala Leu Met Gln Ala Leu Ala Leu Arg Glu Gly Gly Pro Pro Thr 340 345 350			1056
ttc cgg tta acc gga att ggt cca ccg gcg ccg gat aat tct gat cat Phe Arg Leu Thr Gly Ile Gly Pro Pro Ala Pro Asp Asn Ser Asp His 355 360 365			1104
ctt cat gaa gtt ggt tgt aaa tta gct cag ctt gcg gag gcg att cac Leu His Glu Val Gly Cys Lys Leu Ala Gln Leu Ala Glu Ala Ile His 370 375 380			1152
gta gaa ttc gaa tac cgt gga ttc gtt gct aac agc tta gcc gat ctc Val Glu Phe Glu Tyr Arg Gly Phe Val Ala Asn Ser Leu Ala Asp Leu 385 390 395 400			1200
gat gct tcg atg ctt gag ctt aga ccg agc gat acg gaa gct gtt gcg Asp Ala Ser Met Leu Glu Leu Arg Pro Ser Asp Thr Glu Ala Val Ala 405 410 415			1248
gtg aac tct gtt ttt gag cta cat aag ctc tta ggt cgt ccc ggt ggg Val Asn Ser Val Phe Glu Leu His Lys Leu Leu Gly Arg Pro Gly Gly 420 425 430			1296
ata gag aaa gtt ctc ggc gtt gtg aaa cag att aaa ccg gtg att ttc Ile Glu Lys Val Leu Gly Val Val Lys Gln Ile Lys Pro Val Ile Phe 435 440 445			1344
acg gtg gtt gag caa gaa tcg aac cat aac gga ccg gtt ttc tta gac Thr Val Val Glu Gln Glu Ser Asn His Asn Gly Pro Val Phe Leu Asp 450 455 460			1392
cgg ttt act gaa tcg tta cat tat tat tcg act ctg ttt gat tcg ttg Arg Phe Thr Glu Ser Leu His Tyr Tyr Ser Thr Leu Phe Asp Ser Leu 465 470 475 480			1440
gaa gga gtt ccg aat agt caa gac aaa gtc atg tct gaa gtt tac tta Glu Gly Val Pro Asn Ser Gln Asp Lys Val Met Ser Glu Val Tyr Leu 485 490 495			1488
ggg aaa cag att tgt aat ctg gtg gct tgt gaa ggt cct gac aga gtc Gly Lys Gln Ile Cys Asn Leu Val Ala Cys Glu Gly Pro Asp Arg Val 500 505 510			1536
gag aga cac gaa acg ttg agt caa tgg gga aac ccg ttt ggt tcg tcc Glu Arg His Glu Thr Leu Ser Gln Trp Gly Asn Arg Phe Gly Ser Ser 515 520 525			1584
ggt tta gcg ccg gca cat ctt ggg tct aac gcg ttt aag caa gcg agt Gly Leu Ala Pro Ala His Leu Gly Ser Asn Ala Phe Lys Gln Ala Ser 530 535 540			1632
atg ctt ttg tct gtg ttt aat agt ggc caa ggt tat cgt gtg gag gag Met Leu Leu Ser Val Phe Asn Ser Gly Gln Gly Tyr Arg Val Glu Glu 545 550 555 560			1680
agt aat gga tgt ttg atg ttg ggt tgg cac act cgc cca ctc att acc Ser Asn Gly Cys Leu Met Leu Gly Trp His Thr Arg Pro Leu Ile Thr 565 570 575			1728
acc tcc gct tgg aaa ctc tcg acg gcg gcg cac tga Thr Ser Ala Trp Lys Leu Ser Thr Ala Ala His 580 585			1764

<210> 8  
 <211> 587  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 8

Met Lys Arg Asp His His Gln Phe Gln Gly Arg Leu Ser Asn His Gly

## MBI-20 Sequence Listing.ST25

```

1           5           10           15
Thr Ser Ser Ser Ser Ser Ser Ile Ser Lys Asp Lys Met Met Met Val
      20      25      30
Lys Lys Glu Glu Asp Gly Gly Gly Asn Met Asp Asp Glu Leu Leu Ala
      35      40      45
Val Leu Gly Tyr Lys Val Arg Ser Ser Glu Met Ala Glu Val Ala Leu
      50      55      60
Lys Leu Glu Gln Leu Glu Thr Met Met Ser Asn Val Gln Glu Asp Gly
      65      70      75      80
Leu Ser His Leu Ala Thr Asp Thr Val His Tyr Asn Pro Ser Glu Leu
      85      90      95
Tyr Ser Trp Leu Asp Asn Met Leu Ser Glu Leu Asn Pro Pro Pro Leu
      100      105      110
Pro Ala Ser Ser Asn Gly Leu Asp Pro Val Leu Pro Ser Pro Glu Ile
      115      120      125
Cys Gly Phe Pro Ala Ser Asp Tyr Asp Leu Lys Val Ile Pro Gly Asn
      130      135      140
Ala Ile Tyr Gln Phe Pro Ala Ile Asp Ser Ser Ser Ser Ser Asn Asn
      145      150      155      160
Gln Asn Lys Arg Leu Lys Ser Cys Ser Ser Pro Asp Ser Met Val Thr
      165      170      175
Ser Thr Ser Thr Gly Thr Gln Ile Gly Gly Val Ile Gly Thr Thr Val
      180      185      190
Thr Thr Thr Thr Thr Thr Thr Thr Ala Ala Ala Glu Ser Thr Arg Ser
      195      200      205
Val Ile Leu Val Asp Ser Gln Glu Asn Gly Val Arg Leu Val His Ala
      210      215      220
Leu Met Ala Cys Ala Glu Ala Ile Gln Gln Asn Asn Leu Thr Leu Ala
      225      230      235      240
Glu Ala Leu Val Lys Gln Ile Gly Cys Leu Ala Val Ser Gln Ala Gly
      245      250      255
Ala Met Arg Lys Val Ala Thr Tyr Phe Ala Glu Ala Leu Ala Arg Arg
      260      265      270
Ile Tyr Arg Leu Ser Pro Pro Gln Asn Gln Ile Asp His Cys Leu Ser
      275      280      285
Asp Thr Leu Gln Met His Phe Tyr Glu Thr Cys Pro Tyr Leu Lys Phe
      290      295      300

```



## MBI-20 Sequence Listing.ST25

Ala His Phe Thr Ala Asn Gln Ala Ile Leu Glu Ala Phe Glu Gly Lys  
 305 310 315 320

Lys Arg Val His Val Ile Asp Phe Ser Met Asn Gln Gly Leu Gln Trp  
 325 330 335

Pro Ala Leu Met Gln Ala Leu Ala Leu Arg Glu Gly Gly Pro Pro Thr  
 340 345 350

Phe Arg Leu Thr Gly Ile Gly Pro Pro Ala Pro Asp Asn Ser Asp His  
 355 360 365

Leu His Glu Val Gly Cys Lys Leu Ala Gln Leu Ala Glu Ala Ile His  
 370 375 380

Val Glu Phe Glu Tyr Arg Gly Phe Val Ala Asn Ser Leu Ala Asp Leu  
 385 390 395 400

Asp Ala Ser Met Leu Glu Leu Arg Pro Ser Asp Thr Glu Ala Val Ala  
 405 410 415

Val Asn Ser Val Phe Glu Leu His Lys Leu Leu Gly Arg Pro Gly Gly  
 420 425 430

Ile Glu Lys Val Leu Gly Val Val Lys Gln Ile Lys Pro Val Ile Phe  
 435 440 445

Thr Val Val Glu Gln Glu Ser Asn His Asn Gly Pro Val Phe Leu Asp  
 450 455 460

Arg Phe Thr Glu Ser Leu His Tyr Tyr Ser Thr Leu Phe Asp Ser Leu  
 465 470 475 480

Glu Gly Val Pro Asn Ser Gln Asp Lys Val Met Ser Glu Val Tyr Leu  
 485 490 495

Gly Lys Gln Ile Cys Asn Leu Val Ala Cys Glu Gly Pro Asp Arg Val  
 500 505 510

Glu Arg His Glu Thr Leu Ser Gln Trp Gly Asn Arg Phe Gly Ser Ser  
 515 520 525

Gly Leu Ala Pro Ala His Leu Gly Ser Asn Ala Phe Lys Gln Ala Ser  
 530 535 540

Met Leu Leu Ser Val Phe Asn Ser Gly Gln Gly Tyr Arg Val Glu Glu  
 545 550 555 560

Ser Asn Gly Cys Leu Met Leu Gly Trp His Thr Arg Pro Leu Ile Thr  
 565 570 575

Thr Ser Ala Trp Lys Leu Ser Thr Ala Ala His  
 580 585

<210> 9  
 <211> 825  
 <212> DNA

## MBI-20 Sequence Listing.ST25

&lt;213&gt; Arabidopsis thaliana

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(825)

&lt;223&gt; G346

&lt;400&gt; 9

atg gaa atg gaa tca ttc atg gac gac ctt ttg aac ttc tct gta ccg	48
Met Glu Met Glu Ser Phe Met Asp Asp Leu Leu Asn Phe Ser Val Pro	
1 5 10 15	
gaa gag gaa gaa gac gac gac gaa cat acg caa cca ccg agg aat att	96
Glu Glu Glu Glu Asp Asp Asp Glu His Thr Gln Pro Pro Arg Asn Ile	
20 25 30	
act cgc cgg aaa act gga tta cgg cca aca gac tcc ttc ggt ctc ttt	144
Thr Arg Arg Lys Thr Gly Leu Arg Pro Thr Asp Ser Phe Gly Leu Phe	
35 40 45	
aat acc gac gac ctt gga gtg gtt gaa gaa gag gat ttg gaa tgg att	192
Asn Thr Asp Asp Leu Gly Val Val Glu Glu Glu Asp Leu Glu Trp Ile	
50 55 60	
tca aac aaa aat gct ttt ccg gtg att gaa aca ttc gtc ggt gta tta	240
Ser Asn Lys Asn Ala Phe Pro Val Ile Glu Thr Phe Val Gly Val Leu	
65 70 75 80	
ccg tcg gag cat ttt cct ata acg tct ctt ctg gaa aga gaa gcg act	288
Pro Ser Glu His Phe Pro Ile Thr Ser Leu Leu Glu Arg Glu Ala Thr	
85 90 95	
gag gta aaa cag ctg agt ccg gtt tca gta ctt gag acg agt agc cat	336
Glu Val Lys Gln Leu Ser Pro Val Ser Val Leu Glu Thr Ser Ser His	
100 105 110	
agc tcc aca acg act acc tca aac agt agc ggc gga agt aac gga agc	384
Ser Ser Thr Thr Thr Thr Ser Asn Ser Ser Gly Gly Ser Asn Gly Ser	
115 120 125	
acg gcc gtg gct acg acc acc acc act cca aca ata atg agc tgt tgc	432
Thr Ala Val Ala Thr Thr Thr Thr Thr Pro Thr Ile Met Ser Cys Cys	
130 135 140	
gtt ggt ttt aaa gcg ccg gct aaa gcg aga agc aag cgt cgt cgt aca	480
Val Gly Phe Lys Ala Pro Ala Lys Ala Arg Ser Lys Arg Arg Arg Thr	
145 150 155 160	
gga cgc cgt gat tta cga gtt ttg tgg aca gga aac gag caa gga gga	528
Gly Arg Arg Asp Leu Arg Val Leu Trp Thr Gly Asn Glu Gln Gly Gly	
165 170 175	
ata cag aag aag aag acg atg act gtg gcg gcg gct gcg ttg att atg	576
Ile Gln Lys Lys Lys Thr Met Thr Val Ala Ala Ala Ala Leu Ile Met	
180 185 190	
gga agg aag tgt caa cac tgt gga gcg gag aag act ccg caa tgg agg	624
Gly Arg Lys Cys Gln His Cys Gly Ala Glu Lys Thr Pro Gln Trp Arg	
195 200 205	
gca gga cca gcg ggg cct aag act ctg tgt aac gct tgt ggc gtg agg	672
Ala Gly Pro Ala Gly Pro Lys Thr Leu Cys Asn Ala Cys Gly Val Arg	
210 215 220	
tat aag tcc ggg agg cta gtt ccg gag tat cgt cca gcg aac agt cca	720
Tyr Lys Ser Gly Arg Leu Val Pro Glu Tyr Arg Pro Ala Asn Ser Pro	
225 230 235 240	
act ttc acg gcg gag tta cat tcg aat tct cac cgg aag att gta gag	768
Thr Phe Thr Ala Glu Leu His Ser Asn Ser His Arg Lys Ile Val Glu	
245 250 255	
atg agg aag cag tat cag tcc ggt gac ggt gac ggt gat cgg aaa gat	816
Met Arg Lys Gln Tyr Gln Ser Gly Asp Gly Asp Gly Asp Arg Lys Asp	

## 260

825

<400> 10

Thr Phe Thr Ala Glu Leu His Ser Asn Ser His Arg Lys Ile Val Glu

## MBI-20 Sequence Listing.ST25

245

250

255

Met Arg Lys Gln Tyr Gln Ser Gly Asp Gly Asp Gly Asp Arg Lys Asp  
 260 265 270

Cys Gly

<210> 11  
 <211> 1226  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (248)..(1039)  
 <223> G598

<400> 11  
 gtccggtgtc atattttaaa tttatcacct tcttgagaat tccacatttt tatccttttt 60  
 gtcattgtagt gtatattttt tcttctaacc taattaaaat caaaacaaaa tcctttgacc 120  
 caattagctt cgcgatatat cagaagagat caaactactt tgatcagacc atgatcttct 180  
 tcttcttctt cttcttcttc ttcttctttt tagacgatca caattcctaa accctatttc 240  
 tcagatt atg ctg act ctt tac cat caa gaa agg tca ccg gac gcc aca 289  
 Met Leu Thr Leu Tyr His Gln Glu Arg Ser Pro Asp Ala Thr  
 1 5 10  
 agt aat gat cgc gat gag acg cca gag act gtg gtt aga gaa gtc cac 337  
 Ser Asn Asp Arg Asp Glu Thr Pro Glu Thr Val Val Arg Glu Val His  
 15 20 25 30  
 gcg cta act cca gcg ccg gag gat aat tcc cgg acg atg acg gcg acg 385  
 Ala Leu Thr Pro Ala Pro Glu Asp Asn Ser Arg Thr Met Thr Ala Thr  
 35 40 45  
 cta cct cca ccg cct gct ttc cga ggc tat ttt tct cct cca agg tca 433  
 Leu Pro Pro Pro Pro Ala Phe Arg Gly Tyr Phe Ser Pro Pro Arg Ser  
 50 55 60  
 gcg acg acg atg agc gaa gga gag aac ttc aca act ata agc aga gag 481  
 Ala Thr Thr Met Ser Glu Gly Glu Asn Phe Thr Thr Ile Ser Arg Glu  
 65 70 75  
 ttc aac gct cta gtc atc gcc gga tcc tcc atg gag aac aac gaa cta 529  
 Phe Asn Ala Leu Val Ile Ala Gly Ser Ser Met Glu Asn Asn Glu Leu  
 80 85 90  
 atg act cgt gac gtc acg cag cgt gaa gat gag aga caa gac gag ttg 577  
 Met Thr Arg Asp Val Thr Gln Arg Glu Asp Glu Arg Gln Asp Glu Leu  
 95 100 105 110  
 atg aga atc cac gag gac acg gat cat gaa gag gaa acg aat cct tta 625  
 Met Arg Ile His Glu Asp Thr Asp His Glu Glu Glu Thr Asn Pro Leu  
 115 120 125  
 gca atc gtg ccg gat cag tat cct ggt tcg ggt ttg gat cct gga agt 673  
 Ala Ile Val Pro Asp Gln Tyr Pro Gly Ser Gly Leu Asp Pro Gly Ser  
 130 135 140  
 gat aat ggg ccg ggt cag agt cgg gtt ggg tcg acg gtg caa aga gtt 721  
 Asp Asn Gly Pro Gly Gln Ser Arg Val Gly Ser Thr Val Gln Arg Val  
 145 150 155  
 aag agg gaa gag gtg gaa gcg aag ata acg gcg tgg cag acg gca aaa 769  
 Lys Arg Glu Glu Val Glu Ala Lys Ile Thr Ala Trp Gln Thr Ala Lys  
 160 165 170

## MBI-20 Sequence Listing.ST25

ctg gct aag att aat aac agg ttt aag agg gaa gac gcc gtt att aac 817  
 Leu Ala Lys Ile Asn Asn Arg Phe Lys Arg Glu Asp Ala Val Ile Asn  
 175 180 185 190  
 ggt tgg ttt aat gaa caa gtt aac aag gcc aac tct tgg atg aag aaa 865  
 Gly Trp Phe Asn Glu Gln Val Asn Lys Ala Asn Ser Trp Met Lys Lys  
 195 200 205  
 att gag tat aat gta ggt tca ttc aac aat cgt cta aat gag gaa gct 913  
 Ile Glu Tyr Asn Val Gly Ser Phe Asn Asn Arg Leu Asn Glu Glu Ala  
 210 215 220  
 aga gga gag aaa agc aaa agc gat gga gaa aac gca aaa caa tgt ggc 961  
 Arg Gly Glu Lys Ser Lys Ser Asp Gly Glu Asn Ala Lys Gln Cys Gly  
 225 230 235  
 gaa agc gca gag gaa agc gga gga gag aag agc gac ggc aga ggc aaa 1009  
 Glu Ser Ala Glu Glu Ser Gly Gly Glu Lys Ser Asp Gly Arg Gly Lys  
 240 245 250  
 gag agg gac aga ggt tgc aaa agt agt tga agttgctaatt ctcattgagag 1059  
 Glu Arg Asp Arg Gly Cys Lys Ser Ser  
 255 260  
 cccttggaag tcttcctgcc aaacgctcct tcttctcttt ctcctaattt ttagttatat 1119  
 caaaccatta aattaaacag tactcgttat atatctagtt agtaaacaaa ggggcagttt 1179  
 tatagctcat gtacacataa ttgagagtgt agtactgttg tgtcaaa 1226

<210> 12  
 <211> 263  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 12

Met Leu Thr Leu Tyr His Gln Glu Arg Ser Pro Asp Ala Thr Ser Asn  
 1 5 10 15  
 Asp Arg Asp Glu Thr Pro Glu Thr Val Val Arg Glu Val His Ala Leu  
 20 25 30  
 Thr Pro Ala Pro Glu Asp Asn Ser Arg Thr Met Thr Ala Thr Leu Pro  
 35 40 45  
 Pro Pro Pro Ala Phe Arg Gly Tyr Phe Ser Pro Pro Arg Ser Ala Thr  
 50 55 60  
 Thr Met Ser Glu Gly Glu Asn Phe Thr Thr Ile Ser Arg Glu Phe Asn  
 65 70 75 80  
 Ala Leu Val Ile Ala Gly Ser Ser Met Glu Asn Asn Glu Leu Met Thr  
 85 90 95  
 Arg Asp Val Thr Gln Arg Glu Asp Glu Arg Gln Asp Glu Leu Met Arg  
 100 105 110  
 Ile His Glu Asp Thr Asp His Glu Glu Glu Thr Asn Pro Leu Ala Ile  
 115 120 125  
 Val Pro Asp Gln Tyr Pro Gly Ser Gly Leu Asp Pro Gly Ser Asp Asn  
 130 135 140  
 Gly Pro Gly Gln Ser Arg Val Gly Ser Thr Val Gln Arg Val Lys Arg

[illegible]

```
<220>
<221> CDS
<222> (72) .. (1076)
<223> G605
```

<400>	13																			
aattccatcc	taataatttt	caaagcttta	attctaagaa	ataatatcta	caagaaaata												60			
ttatctcatg	t	atg	gag	act	acc	gga	gaa	gtt	gtt	aaa	aca	acc	acc	ggg					110	
		Met	Glu	Thr	Thr	Gly	Glu	Val	Val	Lys	Thr	Thr	Thr	Gly						
	1					5					10									
agc	gac	gga	ggc	gtt	acg	gtg	gtg	aga	tcc	aac	gcg	ccg	tca	gac	ttc					158
Ser	Asp	Gly	Gly	Val	Thr	Val	Val	Arg	Ser	Asn	Ala	Pro	Ser	Asp	Phe					
	15					20					25									
cac	atg	gct	ccg	agg	tca	gaa	act	tca	aac	aca	cct	ccc	aac	tcc	gtc					206
His	Met	Ala	Pro	Arg	Ser	Glu	Thr	Ser	Asn	Ala	Pro	Pro	Asn	Ser	Val					
30					35					40					45					
gct	cct	cct	cct	cct	cca	ccg	ccg	caa	aac	tcc	ttt	act	ccg	tcg	gcg					254
Ala	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Gln	Asn	Ser	Phe	Thr	Pro	Ser	Ala					
				50					55					60						
gct	atg	gat	ggt	ttc	tca	agc	gga	ccg	ata	aag	aag	aga	cgt	ggg	cgc					302
Ala	Met	Asp	Gly	Phe	Ser	Ser	Gly	Pro	Ile	Lys	Lys	Arg	Arg	Gly	Arg					
			65				70					75								
cct	agg	aag	tac	gga	cac	gac	gga	gca	gcg	gtg	acg	cta	tct	ccg	aat					350
Pro	Arg	Lys	Tyr	Gly	His	Asp	Gly	Ala	Ala	Val	Thr	Leu	Ser	Pro	Asn					
		80					85					90								
ccg	ata	tca	tca	gcc	gca	cca	acg	act	tct	cac	gtc	atc	gat	ttc	tcg					398
Pro	Ile	Ser	Ser	Ala	Ala	Pro	Thr	Thr	Ser	His	Val	Ile	Asp	Phe	Ser					
	95					100					105									
acg	aca	tcg	gag	aaa	cgt	ggc	aaa	atg	aaa	cca	gca	act	cca	act	cca					446
Thr	Thr	Ser	Glu	Lys	Arg	Gly	Lys	Met	Lys	Pro	Ala	Thr	Pro	Thr	Pro					

MBI-20 Sequence Listing.ST25

110	115	120	125	
agc tca ttc atc agg cca aag tac cag gtc gag aat tta ggt gaa tgg				494
Ser Ser Phe Ile Arg Pro Lys Tyr Gln Val Glu Asn Leu Gly Glu Trp	130	135	140	
tct cct tcc tct gcc gcc gct aat ttc acg ccg cat att att acg gtg				542
Ser Pro Ser Ser Ala Ala Ala Asn Phe Thr Pro His Ile Ile Thr Val	145	150	155	
aat gca ggc gag gac gtt acg aag agg ata ata tca ttt tct caa caa				590
Asn Ala Gly Glu Asp Val Thr Lys Arg Ile Ile Ser Phe Ser Gln Gln	160	165	170	
ggg tct cta gct att tgc gtt tta tgc gca aac ggt gtc gtt tcg agc				638
Gly Ser Leu Ala Ile Cys Val Leu Cys Ala Asn Gly Val Val Ser Ser	175	180	185	
ggt aca ctt cgt cag cct gat tca tct ggt ggt aca ttg acc tat gag				686
Val Thr Leu Arg Gln Pro Asp Ser Ser Gly Thr Leu Thr Tyr Glu	190	195	200	205
ggt cgg ttt gag ata ttg tca cta tct gga aca ttc atg cct agt gac				734
Gly Arg Phe Glu Ile Leu Ser Leu Ser Gly Thr Phe Met Pro Ser Asp	210	215	220	
tca gac ggg aca cga agc aga aca ggc ggg atg agc gtg tcg ctt gct				782
Ser Asp Gly Thr Arg Ser Arg Thr Gly Gly Met Ser Val Ser Leu Ala	225	230	235	
agc cct gat gga cgt gta gta ggt ggt ggt gtt gct ggc ttg ctg gtt				830
Ser Pro Asp Gly Arg Val Val Gly Gly Gly Val Ala Gly Leu Leu Val	240	245	250	
gca gcc act cct att caa gtg gtt gta gga act ttc tta ggt gga aca				878
Ala Ala Thr Pro Ile Gln Val Val Gly Thr Phe Leu Gly Gly Thr	255	260	265	
aac cag caa gaa cag aca ccg aag ccg cat aac cac aac ttc atg tct				926
Asn Gln Gln Glu Gln Thr Pro Lys Pro His Asn His Asn Phe Met Ser	270	275	280	285
tct cca tta atg cca act tct tcg aat gta gct gat cat cga acc atc				974
Ser Pro Leu Met Pro Thr Ser Ser Asn Val Ala Asp His Arg Thr Ile	290	295	300	
cgt ccc atg aca tct agt ctc ccg atc agt aca tgg aca ccg tct ttt				1022
Arg Pro Met Thr Ser Ser Leu Pro Ile Ser Thr Trp Thr Pro Ser Phe	305	310	315	
cct tct gat tca cga cac aag cat tct cat gac ttt aat atc act ttg				1070
Pro Ser Asp Ser Arg His Lys His Ser His Asp Phe Asn Ile Thr Leu	320	325	330	
acg tga tttcttctct gaagaactcg tagatcctct gtattttggt ttccagttta				1126
Thr				
gggctctaca tgtagactc tcaaagtcta ggtgttatgt tggctgtca cttaggattg				1186
tcacttagga ttgtagacc atctccatca atggtttctc attgagaaac tgttcaatat				1246
aaaaataaaa tataatc				1263

<210> 14  
 <211> 334  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 14

Met	Glu	Thr	Thr	Gly	Glu	Val	Val	Lys	Thr	Thr	Thr	Gly	Ser	Asp	Gly
1				5					10					15	

## MBI-20 Sequence Listing.ST25

Gly Val Thr Val Val Arg Ser Asn Ala Pro Ser Asp Phe His Met Ala  
 20 25 30  
 Pro Arg Ser Glu Thr Ser Asn Thr Pro Pro Asn Ser Val Ala Pro Pro  
 35 40 45  
 Pro Pro Pro Pro Pro Gln Asn Ser Phe Thr Pro Ser Ala Ala Met Asp  
 50 55 60  
 Gly Phe Ser Ser Gly Pro Ile Lys Lys Arg Arg Gly Arg Pro Arg Lys  
 65 70 75 80  
 Tyr Gly His Asp Gly Ala Ala Val Thr Leu Ser Pro Asn Pro Ile Ser  
 85 90 95  
 Ser Ala Ala Pro Thr Thr Ser His Val Ile Asp Phe Ser Thr Thr Ser  
 100 105 110  
 Glu Lys Arg Gly Lys Met Lys Pro Ala Thr Pro Thr Pro Ser Ser Phe  
 115 120 125  
 Ile Arg Pro Lys Tyr Gln Val Glu Asn Leu Gly Glu Trp Ser Pro Ser  
 130 135 140  
 Ser Ala Ala Ala Asn Phe Thr Pro His Ile Ile Thr Val Asn Ala Gly  
 145 150 155 160  
 Glu Asp Val Thr Lys Arg Ile Ile Ser Phe Ser Gln Gln Gly Ser Leu  
 165 170 175  
 Ala Ile Cys Val Leu Cys Ala Asn Gly Val Val Ser Ser Val Thr Leu  
 180 185 190  
 Arg Gln Pro Asp Ser Ser Gly Gly Thr Leu Thr Tyr Glu Gly Arg Phe  
 195 200 205  
 Glu Ile Leu Ser Leu Ser Gly Thr Phe Met Pro Ser Asp Ser Asp Gly  
 210 215 220  
 Thr Arg Ser Arg Thr Gly Gly Met Ser Val Ser Leu Ala Ser Pro Asp  
 225 230 235 240  
 Gly Arg Val Val Gly Gly Gly Val Ala Gly Leu Leu Val Ala Ala Thr  
 245 250 255  
 Pro Ile Gln Val Val Val Gly Thr Phe Leu Gly Gly Thr Asn Gln Gln  
 260 265 270  
 Glu Gln Thr Pro Lys Pro His Asn His Asn Phe Met Ser Ser Pro Leu  
 275 280 285  
 Met Pro Thr Ser Ser Asn Val Ala Asp His Arg Thr Ile Arg Pro Met  
 290 295 300  
 Thr Ser Ser Leu Pro Ile Ser Thr Trp Thr Pro Ser Phe Pro Ser Asp  
 305 310 315 320



Ser Arg His Lys His Ser His Asp Phe Asn Ile Thr Leu Thr  
325 330

```
<220>
<221> CDS
<222> (54) .. (914)
<223> G777
```

<400>	15																	56				
gtggctctct	ctttatcttt	cttggagttt	agtttagagat	tttaacgttg	caa	atg																
						Met																
						1																
gat	caa	cca	atg	aaa	cca	aaa	act	tgc	tct	gaa	tct	gat	ttt	gct	gat	104						
Asp	Gln	Pro	Met	Lys	Pro	Lys	Thr	Cys	Ser	Glu	Ser	Asp	Phe	Ala	Asp							
						5							15									
gat	tcc	tct	gct	tct	tct	tct	tct	tct	tgc	gga	caa	aat	ctc	aga	gga	152						
Asp	Ser	Ser	Ala	Ser	Ser	Ser	Ser	Ser	Ser	Gly	Gln	Asn	Leu	Arg	Gly							
						20							30									
gct	gag	atg	gtg	gtg	gaa	gtg	aag	aag	gaa	gca	gtt	tgt	tcc	cag	aaa	200						
Ala	Glu	Met	Val	Val	Glu	Val	Lys	Lys	Glu	Ala	Val	Cys	Ser	Gln	Lys							
						35							45									
gca	gag	cga	gag	aag	ctt	cgt	aga	gat	aag	ctt	aag	gaa	cag	ttt	ctt	248						
Ala	Glu	Arg	Glu	Lys	Leu	Arg	Arg	Asp	Lys	Leu	Lys	Glu	Gln	Phe	Leu							
						55							60									
gag	ctt	gga	aat	gca	ctt	gat	ccg	aat	agg	cct	aag	agt	gac	aaa	gcc	296						
Glu	Leu	Gly	Asn	Ala	Leu	Asp	Pro	Asn	Arg	Pro	Lys	Ser	Asp	Lys	Ala							
						70							75									
tca	gtt	ctc	act	gat	aca	ata	caa	atg	ctc	aag	gat	gta	atg	aac	caa	344						
Ser	Val	Leu	Thr	Asp	Thr	Ile	Gln	Met	Leu	Lys	Asp	Val	Met	Asn	Gln							
						85							95									
gtt	gat	aga	cta	aaa	gct	gag	tat	gaa	aca	cta	tct	caa	gag	tct	cgt	392						
Val	Asp	Arg	Leu	Lys	Ala	Glu	Tyr	Glu	Thr	Leu	Ser	Gln	Glu	Ser	Arg							
						100							110									
gag	cta	att	caa	gag	aag	agt	gag	ctg	aga	gag	gag	aaa	gcg	act	tta	440						
Glu	Leu	Ile	Gln	Glu	Lys	Ser	Glu	Leu	Arg	Glu	Glu	Lys	Ala	Thr	Leu							
						115							125									
aag	tct	gat	atc	gag	att	ctt	aat	gct	caa	tat	cag	cat	aga	atc	aaa	488						
Lys	Ser	Asp	Ile	Glu	Ile	Leu	Asn	Ala	Gln	Tyr	Gln	His	Arg	Ile	Lys							
						135							140									
acc	atg	gtt	cca	tgg	gta	cct	cat	tac	agt	tat	cat	atc	ccc	ttc	gta	536						
Thr	Met	Val	Pro	Trp	Val	Pro	His	Tyr	Ser	Tyr	His	Ile	Pro	Phe	Val							
						150							155									
gcc	ata	act	cag	ggt	cag	tcc	agt	ttt	ata	cct	tat	tca	gcc	tct	gtc	584						
Ala	Ile	Thr	Gln	Gly	Gln	Ser	Ser	Phe	Ile	Pro	Tyr	Ser	Ala	Ser	Val							
						165							175									
aat	cct	cta	acc	gaa	caa	caa	gca	tgc	gtt	cag	cag	cat	tct	tct	tct	632						
Asn	Pro	Leu	Thr	Glu	Gln	Gln	Ala	Ser	Val	Gln	Gln	His	Ser	Ser	Ser							
						180							185									
tct	gcc	gat	gct	tca	atg	aaa	caa	gat	tcc	aaa	atc	aag	ccg	tta	gat	680						
Ser	Ala	Asp	Ala	Ser	Met	Lys	Gln	Asp	Ser	Lys	Ile	Lys	Pro	Leu	Asp							
						195							205									
ttg	gat	ctg	atg	atg	aac	agt	aac	cat	tca	ggt	caa	gga	aat	gat	caa	728						

## MBI-20 Sequence Listing.ST25

Leu Asp Leu Met Met Asn Ser Asn His Ser Gly Gln Gly Asn Asp Gln  
 210 215 220 225  
 aaa gat gat gtt cgt tta aag ctc gag ctt aaa atc cat gcc tct tct 776  
 Lys Asp Asp Val Arg Leu Lys Leu Glu Leu Lys Ile His Ala Ser Ser  
 230 235 240  
 tta gct caa cag gat gtt tct gga aaa gag aag aaa gta agc ttg aca 824  
 Leu Ala Gln Gln Asp Val Ser Gly Lys Glu Lys Lys Val Ser Leu Thr  
 245 250 255  
 acc act gca agc tca tcg aat agt tac tca tta tct caa gct gtt caa 872  
 Thr Thr Ala Ser Ser Ser Asn Ser Tyr Ser Leu Ser Gln Ala Val Gln  
 260 265 270  
 gat agt tcc ccc ggt acc gta aat gac atg ttg aag cca taa 914  
 Asp Ser Ser Pro Gly Thr Val Asn Asp Met Leu Lys Pro  
 275 280 285  
 accaataaac atattcccct gaacttgtgt ttaataccgt gattgagaag gtaccatgat 974  
 taaacttggt gtagattatc cacatgatta acgatgtatt cttatcacia gcaaataaaa 1034  
 cacaaaagca ttgcttaaa aaa 1057

<210> 16  
 <211> 286  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 16

Met Asp Gln Pro Met Lys Pro Lys Thr Cys Ser Glu Ser Asp Phe Ala  
 1 5 10 15  
 Asp Asp Ser Ser Ala Ser Ser Ser Ser Ser Ser Gly Gln Asn Leu Arg  
 20 25 30  
 Gly Ala Glu Met Val Val Glu Val Lys Lys Glu Ala Val Cys Ser Gln  
 35 40 45  
 Lys Ala Glu Arg Glu Lys Leu Arg Arg Asp Lys Leu Lys Glu Gln Phe  
 50 55 60  
 Leu Glu Leu Gly Asn Ala Leu Asp Pro Asn Arg Pro Lys Ser Asp Lys  
 65 70 75 80  
 Ala Ser Val Leu Thr Asp Thr Ile Gln Met Leu Lys Asp Val Met Asn  
 85 90 95  
 Gln Val Asp Arg Leu Lys Ala Glu Tyr Glu Thr Leu Ser Gln Glu Ser  
 100 105 110  
 Arg Glu Leu Ile Gln Glu Lys Ser Glu Leu Arg Glu Glu Lys Ala Thr  
 115 120 125  
 Leu Lys Ser Asp Ile Glu Ile Leu Asn Ala Gln Tyr Gln His Arg Ile  
 130 135 140  
 Lys Thr Met Val Pro Trp Val Pro His Tyr Ser Tyr His Ile Pro Phe  
 145 150 155 160  
 Val Ala Ile Thr Gln Gly Gln Ser Ser Phe Ile Pro Tyr Ser Ala Ser  
 165 170 175

## MBI-20 Sequence Listing.ST25

Val Asn Pro Leu Thr Glu Gln Gln Ala Ser Val Gln Gln His Ser Ser  
 180 185 190

Ser Ser Ala Asp Ala Ser Met Lys Gln Asp Ser Lys Ile Lys Pro Leu  
 195 200 205

Asp Leu Asp Leu Met Met Asn Ser Asn His Ser Gly Gln Gly Asn Asp  
 210 215 220

Gln Lys Asp Asp Val Arg Leu Lys Leu Glu Leu Lys Ile His Ala Ser  
 225 230 235 240

Ser Leu Ala Gln Gln Asp Val Ser Gly Lys Glu Lys Lys Val Ser Leu  
 245 250 255

Thr Thr Thr Ala Ser Ser Ser Asn Ser Tyr Ser Leu Ser Gln Ala Val  
 260 265 270

Gln Asp Ser Ser Pro Gly Thr Val Asn Asp Met Leu Lys Pro  
 275 280 285

<210> 17  
 <211> 1571  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (428)..(1402)  
 <223> G869

<400> 17  
 aggaacagtg aaagggttcgg ttttttggggt ttcgatctga taatcaacaa gaaaaaaggg 60  
 tttgatttat gtcggctggg tttgaatcga ctgtgatttt gtctttgatt catatctctt 120  
 ctccgatttc atcatcatct tccccatcat cgtcgtcttt gaaatcttgt cttctcaacg 180  
 ctcttcactt ctgctgtaat aagcagaggc ttgttctgga gactccttct cttccatgc 240  
 gcttaagacc caaaaggact tgttctagtg ttgaagtctt tggggggttt cacataaagc 300  
 agcaaaagtt tctttttttc atagttcgct gagagttttg agttttgata ccaaaaaagt 360  
 tttgaccttt tagagtgatt ttttgttctt tctgttttct ggggtatttt gaggagtggg 420  
 tttaaca atg gtt gcg att aga aag gaa cag tct ttg agt ggt gtt agt 469  
 Met Val Ala Ile Arg Lys Glu Gln Ser Leu Ser Gly Val Ser  
 1 5 10

agc gag att aag aag aga gct aag aga aac act cta tcg tcc ctt cct 517  
 Ser Glu Ile Lys Lys Arg Ala Lys Arg Asn Thr Leu Ser Ser Leu Pro  
 15 20 25 30

caa gaa acc caa cct ttg agg aaa gtc cgt att att gtg aat gat cct 565  
 Gln Glu Thr Gln Pro Leu Arg Lys Val Arg Ile Ile Val Asn Asp Pro  
 35 40 45

tat gct act gat gat tcc tct agt gat gag gaa gag ctt aag gtt cct 613  
 Tyr Ala Thr Asp Asp Ser Ser Ser Asp Glu Glu Glu Leu Lys Val Pro  
 50 55 60

aag cca agg aaa atg aaa cgt atc gtt cgt gag att aac ttt cct tct 661  
 Lys Pro Arg Lys Met Lys Arg Ile Val Arg Glu Ile Asn Phe Pro Ser  
 65 70 75

## MBI-20 Sequence Listing.ST25

atg gaa gtt tct gaa cag cct tct gag agt tct tct cag gac agt act	709
Met Glu Val Ser Glu Gln Pro Ser Glu Ser Ser Ser Gln Asp Ser Thr	
80 85 90	
aaa act gat ggc aag ata gct gtg tca gct tct cct gct gtt cct agg	757
Lys Thr Asp Gly Lys Ile Ala Val Ser Ala Ser Pro Ala Val Pro Arg	
95 100 105 110	
aag aag cct gtt ggt gtt agg caa agg aaa tgg ggg aaa tgg gct gct	805
Lys Lys Pro Val Gly Val Arg Gln Arg Lys Trp Gly Lys Trp Ala Ala	
115 120 125	
gag att aga gat cct att aag aaa act agg act tgg ttg ggt act ttt	853
Glu Ile Arg Asp Pro Ile Lys Lys Thr Arg Thr Trp Leu Gly Thr Phe	
130 135 140	
gat act ctt gaa gaa gct gct aaa gct tat gat gct aag aag ctt gag	901
Asp Thr Leu Glu Glu Ala Ala Lys Ala Tyr Asp Ala Lys Lys Leu Glu	
145 150 155	
ttt gat gct att gtt gct gga aat gtg tcc act act aaa cgt gat gtt	949
Phe Asp Ala Ile Val Ala Gly Asn Val Ser Thr Thr Lys Arg Asp Val	
160 165 170	
tct tca tct gag act agc caa tgc tct cgt tct tca cct gtt gtt cct	997
Ser Ser Ser Glu Thr Ser Gln Cys Ser Arg Ser Ser Pro Val Val Pro	
175 180 185 190	
gtt gag caa gat gac act tct gca tca gct ctc act tgt gtc aac aac	1045
Val Glu Gln Asp Val Thr Ser Ala Ser Ala Leu Thr Cys Val Asn Asn	
195 200 205	
cct gat gac gtc tcg acc gtt gct cca act gct cca act cca aat gtt	1093
Pro Asp Asp Val Ser Thr Val Ala Pro Thr Ala Pro Thr Pro Asn Val	
210 215 220	
cct gct ggt gga aac aag gaa acg ttg ttc gat ttc gac ttt act aat	1141
Pro Ala Gly Gly Asn Lys Glu Thr Leu Phe Asp Phe Asp Phe Thr Asn	
225 230 235	
cta cag atc cct gat ttt ggt ttc ttg gca gag gag caa caa gac cta	1189
Leu Gln Ile Pro Asp Phe Gly Phe Leu Ala Glu Glu Gln Gln Asp Leu	
240 245 250	
gac ttc gat tgt ttc ctc gcg gat gat cag ttt gat gat ttc ggc ttg	1237
Asp Phe Asp Cys Phe Ala Asp Asp Gln Phe Asp Asp Phe Gly Leu	
255 260 265 270	
ctt gat gac att caa gga ttc gaa gat aac ggt cca agt gcg tta cca	1285
Leu Asp Asp Ile Gln Gly Phe Glu Asp Asn Gly Pro Ser Ala Leu Pro	
275 280 285	
gat ttc gac ttt gcg gat gtt gaa gat ctt cag cta gct gac tct agt	1333
Asp Phe Asp Phe Ala Asp Val Glu Asp Leu Gln Leu Ala Asp Ser Ser	
290 295 300	
ttc ggt ttc ctt gat caa ctt gct cct atc aac atc tct tgc cca tta	1381
Phe Gly Phe Leu Asp Gln Leu Ala Pro Ile Asn Ile Ser Cys Pro Leu	
305 310 315	
aaa agt ttt gca gct tca tag gatcttgctt agtaagtta agtgagaaga	1432
Lys Ser Phe Ala Ala Ser	
320	
gtgttttggt ttttcgttta tgcttttagta atttaagaca tacaaaagtg tgtgttccgg	1492
attgtagtaa gatcttaaga cataaagccg gggttttgcaa ttaggaaatcg agttttaatg	1552
aagtttttagt ttatgtttg	1571

<210> 18  
 <211> 324  
 <212> PRT

## MBI-20 Sequence Listing.ST25

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 18

```

Met Val Ala Ile Arg Lys Glu Gln Ser Leu Ser Gly Val Ser Ser Glu
1      5      10      15

Ile Lys Lys Arg Ala Lys Arg Asn Thr Leu Ser Ser Leu Pro Gln Glu
20      25      30

Thr Gln Pro Leu Arg Lys Val Arg Ile Ile Val Asn Asp Pro Tyr Ala
35      40      45

Thr Asp Asp Ser Ser Ser Asp Glu Glu Glu Leu Lys Val Pro Lys Pro
50      55      60

Arg Lys Met Lys Arg Ile Val Arg Glu Ile Asn Phe Pro Ser Met Glu
65      70      75      80

Val Ser Glu Gln Pro Ser Glu Ser Ser Ser Gln Asp Ser Thr Lys Thr
85      90      95

Asp Gly Lys Ile Ala Val Ser Ala Ser Pro Ala Val Pro Arg Lys Lys
100     105     110

Pro Val Gly Val Arg Gln Arg Lys Trp Gly Lys Trp Ala Ala Glu Ile
115     120     125

Arg Asp Pro Ile Lys Lys Thr Arg Thr Trp Leu Gly Thr Phe Asp Thr
130     135     140

Leu Glu Glu Ala Ala Lys Ala Tyr Asp Ala Lys Lys Leu Glu Phe Asp
145     150     155     160

Ala Ile Val Ala Gly Asn Val Ser Thr Thr Lys Arg Asp Val Ser Ser
165     170     175

Ser Glu Thr Ser Gln Cys Ser Arg Ser Ser Pro Val Val Pro Val Glu
180     185     190

Gln Asp Asp Thr Ser Ala Ser Ala Leu Thr Cys Val Asn Asn Pro Asp
195     200     205

Asp Val Ser Thr Val Ala Pro Thr Ala Pro Thr Pro Asn Val Pro Ala
210     215     220

Gly Gly Asn Lys Glu Thr Leu Phe Asp Phe Asp Phe Thr Asn Leu Gln
225     230     235     240

Ile Pro Asp Phe Gly Phe Leu Ala Glu Glu Gln Gln Asp Leu Asp Phe
245     250     255

Asp Cys Phe Leu Ala Asp Asp Gln Phe Asp Asp Phe Gly Leu Leu Asp
260     265     270

Asp Ile Gln Gly Phe Glu Asp Asn Gly Pro Ser Ala Leu Pro Asp Phe
275     280     285

```



## MBI-20 Sequence Listing.ST25

Asp Phe Ala Asp Val Glu Asp Leu Gln Leu Ala Asp Ser Ser Phe Gly  
 290 295 300

Phe Leu Asp Gln Leu Ala Pro Ile Asn Ile Ser Cys Pro Leu Lys Ser  
 305 310 315 320

Phe Ala Ala Ser

<210> 19  
 <211> 1322  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (104)..(1084)  
 <223> G1133

<400> 19  
 ttcaagaaag aatcaccaag tgttgcgttc cacacatttg agcaacagct tccacaatcg 60  
 tattgtattc ctgtaaagtt cccttggtt aaactgcaag agc atg cct ctt gat 115  
 Met Pro Leu Asp  
 1  
 acc aaa cag cag aaa tgg ttg cca tta ggc tta aat cct caa gct tgt 163  
 Thr Lys Gln Gln Lys Trp Leu Pro Leu Gly Leu Asn Pro Gln Ala Cys  
 5 10 15 20  
 gtc cag gac aag gcg act gag tat ttc cgt cct gga att cct ttt ccg 211  
 Val Gln Asp Lys Ala Thr Glu Tyr Phe Arg Pro Gly Ile Pro Phe Pro  
 25 30 35  
 gaa ctc ggt aaa gtt tat gca gct gag cat cag ttt cgc tat ttg cag 259  
 Glu Leu Gly Lys Val Tyr Ala Ala Glu His Gln Phe Arg Tyr Leu Gln  
 40 45 50  
 cca ccg ttc caa gcc tta ttg tct aga tat gat cag cag tct tgt gga 307  
 Pro Pro Phe Gln Ala Leu Leu Ser Arg Tyr Asp Gln Gln Ser Cys Gly  
 55 60 65  
 aaa caa gtt tca tgt ttg aat ggg cga tct agc aac ggt gct gct cca 355  
 Lys Gln Val Ser Cys Leu Asn Gly Arg Ser Ser Asn Gly Ala Ala Pro  
 70 75 80  
 gag ggg gca ctc aag tct tct cgg aaa aga ttt ata gta ttc gat cag 403  
 Glu Gly Ala Leu Lys Ser Ser Arg Lys Arg Phe Ile Val Phe Asp Gln  
 85 90 95 100  
 tcg gga gag cag act cgt ttg tta caa tgt gga ttt cct ctg cgg ttt 451  
 Ser Gly Glu Gln Thr Arg Leu Leu Gln Cys Gly Phe Pro Leu Arg Phe  
 105 110 115  
 cct tct tct atg gat gca gag cga ggg aac att ctc ggt gcc cta cac 499  
 Pro Ser Ser Met Asp Ala Glu Arg Gly Asn Ile Leu Gly Ala Leu His  
 120 125 130  
 cca gag aaa ggg ttt agt aaa gat cat gcc att caa gaa aag ata ttg 547  
 Pro Glu Lys Gly Phe Ser Lys Asp His Ala Ile Gln Glu Lys Ile Leu  
 135 140 145  
 caa cat gaa gat cat gaa aat ggc gaa gaa gac tcg gaa atg cac gaa 595  
 Gln His Glu Asp His Glu Asn Gly Glu Glu Asp Ser Glu Met His Glu  
 150 155 160  
 gac act gag gaa atc aac gcg tta ctg tat tct gat gat gac gat aat 643  
 Asp Thr Glu Glu Ile Asn Ala Leu Leu Tyr Ser Asp Asp Asp Asp Asn  
 165 170 175 180

## MBI-20 Sequence Listing.ST25

```

gat gat tgg gaa agt gat gat gaa gta atg agc act ggt cac tct cca      691
Asp Asp Trp Glu Ser Asp Asp Glu Val Met Ser Thr Gly His Ser Pro
185                               190                               195

ttc aca gtt gaa caa caa gcg tgc aac ata aca aca gaa gag ctg gat      739
Phe Thr Val Glu Gln Gln Ala Cys Asn Ile Thr Thr Glu Glu Leu Asp
200                               205                               210

gaa act gaa agc act gtt gat ggt cca ctt ctt aaa aga cag aaa cta      787
Glu Thr Glu Ser Thr Val Asp Gly Pro Leu Leu Lys Arg Gln Lys Leu
215                               220                               225

ctg gac cat tcg tac aga gac tca tca cca tcc ctt gtg ggc acc act      835
Leu Asp His Ser Tyr Arg Asp Ser Ser Pro Ser Leu Val Gly Thr Thr
230                               235                               240

aaa gtc aaa ggc tta tca gat gaa aac ctt cct gaa tca aac att tca      883
Lys Val Lys Gly Leu Ser Asp Glu Asn Leu Pro Glu Ser Asn Ile Ser
245                               250                               255

agc aaa caa gaa acg ggt tct ggt ttg agc gac gag cag tca aga aaa      931
Ser Lys Gln Glu Thr Gly Ser Gly Leu Ser Asp Glu Gln Ser Arg Lys
265                               270                               275

gac aag att cac acc gct ctg aga atc ctg gag agt gta gtt cca ggg      979
Asp Lys Ile His Thr Ala Leu Arg Ile Leu Glu Ser Val Val Pro Gly
280                               285                               290

gca aag gga aaa gaa gct ctt tta cta cta gac gaa gcc att gat tac     1027
Ala Lys Gly Lys Glu Ala Leu Leu Leu Leu Asp Glu Ala Ile Asp Tyr
295                               300                               305

ctc aag ttg ctg aag caa agc tta aac tca tca aag ggt ttg aat aac     1075
Leu Lys Leu Leu Lys Gln Ser Leu Asn Ser Ser Lys Gly Leu Asn Asn
310                               315                               320

cat tgg tga aaaacctaca accccttttg tcctattgat aaggcatgtt          1124
His Trp
325

tggttggtta aagagaagac atgggacaaa agataatcaa tgaggtaaag gactgatgaa    1184

gaagattctc tcaaattcat taacgtgggt ttgaacaat tagaacacgc ctggtgaccc    1244

tagtgggacc gtatccactg ttcattctagc tggatcaata gtggtttact tttggatttg    1304

gcatgctctc tcaaaaaa                                                1322

<210> 20
<211> 326
<212> PRT
<213> Arabidopsis thaliana

<400> 20
Met Pro Leu Asp Thr Lys Gln Gln Lys Trp Leu Pro Leu Gly Leu Asn
1                               5                               10                               15

Pro Gln Ala Cys Val Gln Asp Lys Ala Thr Glu Tyr Phe Arg Pro Gly
20                               25                               30

Ile Pro Phe Pro Glu Leu Gly Lys Val Tyr Ala Ala Glu His Gln Phe
35                               40                               45

Arg Tyr Leu Gln Pro Pro Phe Gln Ala Leu Leu Ser Arg Tyr Asp Gln
50                               55                               60

Gln Ser Cys Gly Lys Gln Val Ser Cys Leu Asn Gly Arg Ser Ser Asn
65                               70                               75                               80

```



## MBI-20 Sequence Listing.ST25

Gly Ala Ala Pro Glu Gly Ala Leu Lys Ser Ser Arg Lys Arg Phe Ile  
                     85                    90                    95  
 Val Phe Asp Gln Ser Gly Glu Gln Thr Arg Leu Leu Gln Cys Gly Phe  
                     100                    105                    110  
 Pro Leu Arg Phe Pro Ser Ser Met Asp Ala Glu Arg Gly Asn Ile Leu  
                     115                    120                    125  
 Gly Ala Leu His Pro Glu Lys Gly Phe Ser Lys Asp His Ala Ile Gln  
                     130                    135                    140  
 Glu Lys Ile Leu Gln His Glu Asp His Glu Asn Gly Glu Glu Asp Ser  
                     145                    150                    155                    160  
 Glu Met His Glu Asp Thr Glu Glu Ile Asn Ala Leu Leu Tyr Ser Asp  
                     165                    170                    175  
 Asp Asp Asp Asn Asp Asp Trp Glu Ser Asp Asp Glu Val Met Ser Thr  
                     180                    185                    190  
 Gly His Ser Pro Phe Thr Val Glu Gln Gln Ala Cys Asn Ile Thr Thr  
                     195                    200                    205  
 Glu Glu Leu Asp Glu Thr Glu Ser Thr Val Asp Gly Pro Leu Leu Lys  
                     210                    215                    220  
 Arg Gln Lys Leu Leu Asp His Ser Tyr Arg Asp Ser Ser Pro Ser Leu  
                     225                    230                    235                    240  
 Val Gly Thr Thr Lys Val Lys Gly Leu Ser Asp Glu Asn Leu Pro Glu  
                     245                    250                    255  
 Ser Asn Ile Ser Ser Lys Gln Glu Thr Gly Ser Gly Leu Ser Asp Glu  
                     260                    265                    270  
 Gln Ser Arg Lys Asp Lys Ile His Thr Ala Leu Arg Ile Leu Glu Ser  
                     275                    280                    285  
 Val Val Pro Gly Ala Lys Gly Lys Glu Ala Leu Leu Leu Leu Asp Glu  
                     290                    295                    300  
 Ala Ile Asp Tyr Leu Lys Leu Leu Lys Gln Ser Leu Asn Ser Ser Lys  
                     305                    310                    315                    320  
 Gly Leu Asn Asn His Trp  
                     325

<210> 21  
 <211> 859  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (62)..(718)  
 <223> G1266

## MBI-20 Sequence Listing.ST25

<400> 21  
 caatccacta acgatacccta accgaaaaca gagtagtcaa gaaacagagt attttttcta 60  
 c atg gat cca ttt tta att cag tcc cca ttc tcc ggc ttc tca ccg gaa 109  
 Met Asp Pro Phe Leu Ile Gln Ser Pro Phe Ser Gly Phe Ser Pro Glu  
 1 5 10 15  
 tat tct atc gga tct tct cca gat tct ttc tca tcc tct tct tct aac 157  
 Tyr Ser Ile Gly Ser Ser Pro Asp Ser Phe Ser Ser Ser Ser Ser Asn  
 20 25 30  
 aat tac tct ctt ccc ttc aac gag aac gac tca gag gaa atg ttt ctc 205  
 Asn Tyr Ser Leu Pro Phe Asn Glu Asn Asp Ser Glu Glu Met Phe Leu  
 35 40 45  
 tac ggt cta atc gag cag tcc acg caa caa acc tat att gac tcg gat 253  
 Tyr Gly Leu Ile Glu Gln Ser Thr Gln Gln Thr Tyr Ile Asp Ser Asp  
 50 55 60  
 agt caa gac ctt ccg atc aaa tcc gta agc tca aga aag tca gag aag 301  
 Ser Gln Asp Leu Pro Ile Lys Ser Val Ser Ser Arg Lys Ser Glu Lys  
 65 70 75 80  
 tct tac aga ggc gta aga cga cgg cca tgg ggg aaa ttc gcg gcg gag 349  
 Ser Tyr Arg Gly Val Arg Arg Arg Pro Trp Gly Lys Phe Ala Ala Glu  
 85 90 95  
 ata aga gat tcg act aga aac ggt att agg gtt tgg ctc ggg acg ttc 397  
 Ile Arg Asp Ser Thr Arg Asn Gly Ile Arg Val Trp Leu Gly Thr Phe  
 100 105 110  
 gaa agc gcg gaa gag gcg gct tta gcc tac gat caa gct gct ttc tcg 445  
 Glu Ser Ala Glu Glu Ala Ala Leu Ala Tyr Asp Gln Ala Ala Phe Ser  
 115 120 125  
 atg aga ggg tcc tcg gcg att ctc aat ttt tcg gcg gag aga gtt caa 493  
 Met Arg Gly Ser Ser Ala Ile Leu Asn Phe Ser Ala Glu Arg Val Gln  
 130 135 140  
 gag tcg ctt tcg gag att aaa tat acc tac gag gat ggt tgt tct ccg 541  
 Glu Ser Leu Ser Glu Ile Lys Tyr Thr Tyr Glu Asp Gly Cys Ser Pro  
 145 150 155 160  
 gtt gtg gcg ttg aag agg aaa cac tcg atg aga cgg aga atg acc aat 589  
 Val Val Ala Leu Lys Arg Lys His Ser Met Arg Arg Arg Met Thr Asn  
 165 170 175  
 aag aag acg aaa gat agt gac ttt gat cac cgc tcc gtg aag tta gat 637  
 Lys Lys Thr Lys Asp Ser Asp Phe Asp His Arg Ser Val Lys Leu Asp  
 180 185 190  
 aat gta gtt gtc ttt gag gat ttg gga gaa cag tac ctt gag gag ctt 685  
 Asn Val Val Val Phe Glu Asp Leu Gly Glu Gln Tyr Leu Glu Glu Leu  
 195 200 205  
 ttg ggg tct tct gaa aat agt ggg act tgg tga aagattagga tttgtattag 738  
 Leu Gly Ser Ser Glu Asn Ser Gly Thr Trp  
 210 215  
 ggaccttaag tttgaagtgg ttgattaatt ttaaccctaa tatgtttttt gtttgcttaa 798  
 atatttgatt ctattgagaa acatcgaaaa cagtttgtat gtacttttgt gatacttggc 858  
 g 859

<210> 22  
 <211> 218  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 22  
 Met Asp Pro Phe Leu Ile Gln Ser Pro Phe Ser Gly Phe Ser Pro Glu

## MBI-20 Sequence Listing.ST25

```

1           5           10           15
Tyr Ser Ile Gly Ser Ser Pro Asp Ser Phe Ser Ser Ser Ser Ser Asn
20           25           30
Asn Tyr Ser Leu Pro Phe Asn Glu Asn Asp Ser Glu Glu Met Phe Leu
35           40           45
Tyr Gly Leu Ile Glu Gln Ser Thr Gln Gln Thr Tyr Ile Asp Ser Asp
50           55           60
Ser Gln Asp Leu Pro Ile Lys Ser Val Ser Ser Arg Lys Ser Glu Lys
65           70           75           80
Ser Tyr Arg Gly Val Arg Arg Arg Pro Trp Gly Lys Phe Ala Ala Glu
85           90           95
Ile Arg Asp Ser Thr Arg Asn Gly Ile Arg Val Trp Leu Gly Thr Phe
100          105          110
Glu Ser Ala Glu Glu Ala Ala Leu Ala Tyr Asp Gln Ala Ala Phe Ser
115          120          125
Met Arg Gly Ser Ser Ala Ile Leu Asn Phe Ser Ala Glu Arg Val Gln
130          135          140
Glu Ser Leu Ser Glu Ile Lys Tyr Thr Tyr Glu Asp Gly Cys Ser Pro
145          150          155          160
Val Val Ala Leu Lys Arg Lys His Ser Met Arg Arg Arg Met Thr Asn
165          170          175
Lys Lys Thr Lys Asp Ser Asp Phe Asp His Arg Ser Val Lys Leu Asp
180          185          190
Asn Val Val Val Phe Glu Asp Leu Gly Glu Gln Tyr Leu Glu Glu Leu
195          200          205
Leu Gly Ser Ser Glu Asn Ser Gly Thr Trp
210          215

```

```

<210> 23
<211> 1137
<212> DNA
<213> Arabidopsis thaliana

```

```

<220>
<221> CDS
<222> (54)..(914)
<223> G1324

```

```

<400> 23
cgaaaaacacc acaaaccaaa tatcattaag taattaggaa acttaaacta agt atg
Met
1

```

```

gaa aat tcg atg aag aag aag aag agc ttc aaa gaa agt gaa gat gaa
Glu Asn Ser Met Lys Lys Lys Lys Ser Phe Lys Glu Ser Glu Asp Glu
5           10           15

```

## MBI-20 Sequence Listing.ST25

gaa cta aga aga ggg cct tgg act ttg gag gaa gac aca ctt ctc aca	152
Glu Leu Arg Arg Gly Pro Trp Thr Leu Glu Glu Asp Thr Leu Leu Thr	
20 25 30	
aat tac atc ctc cat aac ggt gag ggt cgt tgg aat cac gtc gcc aaa	200
Asn Tyr Ile Leu His Asn Gly Glu Gly Arg Trp Asn His Val Ala Lys	
35 40 45	
tgt gct ggg cta aag aga act ggg aaa agt tgt aga ttg aga tgg ttg	248
Cys Ala Gly Leu Lys Arg Thr Gly Lys Ser Cys Arg Leu Arg Trp Leu	
50 55 60 65	
aat tac ttg aaa ccc gac ata aga cga ggg aat ctt act cct caa gaa	296
Asn Tyr Leu Lys Pro Asp Ile Arg Arg Gly Asn Leu Thr Pro Gln Glu	
70 75 80	
cag ctt ttg atc ctt gag ctt cac tct aaa tgg ggt aat agg tgg tcc	344
Gln Leu Leu Ile Leu Glu Leu His Ser Lys Trp Gly Asn Arg Trp Ser	
85 90 95	
aag att gca cag tac ttg cca gga aga acg gat aac gag atc aag aac	392
Lys Ile Ala Gln Tyr Leu Pro Gly Arg Thr Asp Asn Glu Ile Lys Asn	
100 105 110	
tat tgg aga aca aga gtt caa aaa caa gct cgt caa ctc aac atc gaa	440
Tyr Trp Arg Thr Arg Val Gln Lys Gln Ala Arg Gln Leu Asn Ile Glu	
115 120 125	
tct aac agc gac aag ttc ttt gac gct gtt cgt agt ttt tgg gtc cct	488
Ser Asn Ser Asp Lys Phe Phe Asp Ala Val Arg Ser Phe Trp Val Pro	
130 135 140 145	
aga ttg atc gag aag atg gaa caa aac tca tcc act act act act tat	536
Arg Leu Ile Glu Lys Met Glu Gln Asn Ser Ser Thr Thr Thr Thr Tyr	
150 155 160	
tgt tgt ccc caa aac aac aac aac aac tct ctt ctt ctt cct tct caa	584
Cys Cys Pro Gln Asn Asn Asn Asn Asn Ser Leu Leu Leu Pro Ser Gln	
165 170 175	
tct cac gac tct tta agt atg caa aaa gat ata gat tac tcg ggt ttc	632
Ser His Asp Ser Leu Ser Met Gln Lys Asp Ile Asp Tyr Ser Gly Phe	
180 185 190	
agc aac ata gac ggt tct tct tca act tct act tgc atg tct cat cta	680
Ser Asn Ile Asp Gly Ser Ser Ser Thr Ser Thr Cys Met Ser His Leu	
195 200 205	
aca aca gtt cca cac ttt atg gat caa agc aac acc aat atc atc gat	728
Thr Thr Val Pro His Phe Met Asp Gln Ser Asn Thr Asn Ile Ile Asp	
210 215 220 225	
ggc tcg atg tgt ttc cat gaa ggc aat gtt caa gaa ttc gga gga tat	776
Gly Ser Met Cys Phe His Glu Gly Asn Val Gln Glu Phe Gly Gly Tyr	
230 235 240	
gtt cct ggc atg gag gat tac atg gta aac tcg gac atc tca atg gaa	824
Val Pro Gly Met Glu Asp Tyr Met Val Asn Ser Asp Ile Ser Met Glu	
245 250 255	
tgt cac gtg gcg gat ggt tat tca gcg tac gag gat gtt aca caa gat	872
Cys His Val Ala Asp Gly Tyr Ser Ala Tyr Glu Asp Val Thr Gln Asp	
260 265 270	
ccc atg tgg aat gtg gat gac att tgg cag ttt agg gag taa	914
Pro Met Trp Asn Val Asp Ile Trp Gln Phe Arg Glu	
275 280 285	
ttaagtcgtc aagagatgag atggtagagc ctaccactac ggttctatta tatggactaa	974
tatacttctt ttgcttaact aagcaaaaag tttcgaacct tttaccata ttatctcggg	1034
ttggagacta gaacatgtta aatttgtatc ttctttgttg cgagtactta ctaagtcatt	1094
ggataaatat ttataatgat agtttcttgt acaaaaaaaaaaaa aaa	1137

## MBI-20 Sequence Listing.ST25

<210> 24  
 <211> 286  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 24

Met Glu Asn Ser Met Lys Lys Lys Lys Ser Phe Lys Glu Ser Glu Asp  
 1 5 10 15

Glu Glu Leu Arg Arg Gly Pro Trp Thr Leu Glu Glu Asp Thr Leu Leu  
 20 25 30

Thr Asn Tyr Ile Leu His Asn Gly Glu Gly Arg Trp Asn His Val Ala  
 35 40 45

Lys Cys Ala Gly Leu Lys Arg Thr Gly Lys Ser Cys Arg Leu Arg Trp  
 50 55 60

Leu Asn Tyr Leu Lys Pro Asp Ile Arg Arg Gly Asn Leu Thr Pro Gln  
 65 70 75 80

Glu Gln Leu Leu Ile Leu Glu Leu His Ser Lys Trp Gly Asn Arg Trp  
 85 90 95

Ser Lys Ile Ala Gln Tyr Leu Pro Gly Arg Thr Asp Asn Glu Ile Lys  
 100 105 110

Asn Tyr Trp Arg Thr Arg Val Gln Lys Gln Ala Arg Gln Leu Asn Ile  
 115 120 125

Glu Ser Asn Ser Asp Lys Phe Phe Asp Ala Val Arg Ser Phe Trp Val  
 130 135 140

Pro Arg Leu Ile Glu Lys Met Glu Gln Asn Ser Ser Thr Thr Thr Thr  
 145 150 155 160

Tyr Cys Cys Pro Gln Asn Asn Asn Asn Asn Ser Leu Leu Leu Pro Ser  
 165 170 175

Gln Ser His Asp Ser Leu Ser Met Gln Lys Asp Ile Asp Tyr Ser Gly  
 180 185 190

Phe Ser Asn Ile Asp Gly Ser Ser Ser Thr Ser Thr Cys Met Ser His  
 195 200 205

Leu Thr Thr Val Pro His Phe Met Asp Gln Ser Asn Thr Asn Ile Ile  
 210 215 220

Asp Gly Ser Met Cys Phe His Glu Gly Asn Val Gln Glu Phe Gly Gly  
 225 230 235 240

Tyr Val Pro Gly Met Glu Asp Tyr Met Val Asn Ser Asp Ile Ser Met  
 245 250 255

Glu Cys His Val Ala Asp Gly Tyr Ser Ala Tyr Glu Asp Val Thr Gln  
 260 265 270

## MBI-20 Sequence Listing.ST25

Asp Pro Met Trp Asn Val Asp Asp Ile Trp Gln Phe Arg Glu  
 275 280 285

<210> 25  
 <211> 1630  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (97)..(1398)  
 <223> G1337

<400> 25  
 aatggatttg tcatcattct tctcaccgtc cttagtctct gaaaataaat tctgattttg 60  
 atttcgaatt ttagggattt tgagagagag tcagtt atg agt agt tcg gag aga 114  
 Met Ser Ser Ser Glu Arg  
 1 5  
 gta ccg tgc gat ttc tgc gcc gag cgt acg gcg gtt ttg ttt tgt aga 162  
 Val Pro Cys Asp Phe Cys Gly Glu Arg Thr Ala Val Leu Phe Cys Arg  
 10 15 20  
 gcc gat acg gcg aag ctg tgt ttg cct tgt gat cag caa gtt cac acg 210  
 Ala Asp Thr Ala Lys Leu Cys Leu Pro Cys Asp Gln Gln Val His Thr  
 25 30 35  
 gcg aat ctg ttg tcg agg aag cac gtg cga tct cag atc tgc gat aat 258  
 Ala Asn Leu Leu Ser Arg Lys His Val Arg Ser Gln Ile Cys Asp Asn  
 40 45 50  
 tgc ggt aac gag cca gtc tct gtt cgg tgt ttc acc gat aat ctg att 306  
 Cys Gly Asn Glu Pro Val Ser Val Arg Cys Phe Thr Asp Asn Leu Ile  
 55 60 65 70  
 ttg tgt cag gag tgt gat tgg gat gtt cac gga agt tgt tca gtt tcc 354  
 Leu Cys Gln Glu Cys Asp Trp Asp Val His Gly Ser Cys Ser Val Ser  
 75 80 85  
 gat gct cat gtt cga tcc gcc gtg gaa ggt ttt tcc ggt tgt cca tcg 402  
 Asp Ala His Val Arg Ser Ala Val Glu Gly Phe Ser Gly Cys Pro Ser  
 90 95 100  
 gcg ttg gag ctt gct gct tta tgg gga ctt gat ttg gag caa ggg agg 450  
 Ala Leu Glu Leu Ala Ala Leu Trp Gly Leu Asp Leu Gln Gly Arg  
 105 110 115  
 aaa gat gaa gag aat caa gtt ccg atg atg gcg atg atg atg gat aat 498  
 Lys Asp Glu Glu Asn Gln Val Pro Met Met Ala Met Met Met Asp Asn  
 120 125 130  
 ttc ggg atg cag ttg gat tct tgg gtt ttg gga tct aat gaa ttg att 546  
 Phe Gly Met Gln Leu Asp Ser Trp Val Leu Gly Ser Asn Glu Leu Ile  
 135 140 145 150  
 gtt ccc agc gat acg acg ttt aag aag cgt gga tct tgt gga tct agt 594  
 Val Pro Ser Asp Thr Thr Phe Lys Lys Arg Gly Ser Cys Gly Ser Ser  
 155 160 165  
 tgt ggg agg tat aag cag gta ttg tgt aag cag ctt gag gag ttg ctt 642  
 Cys Gly Arg Tyr Lys Gln Val Leu Cys Lys Gln Leu Glu Leu Leu  
 170 175 180  
 aag agt ggt gtt gtc ggt ggt gat gcc gat gat ggt gat cgt gac cgt 690  
 Lys Ser Gly Val Val Gly Gly Asp Gly Asp Asp Gly Asp Arg Asp Arg  
 185 190 195  
 gat tgt gac cgt gag ggt gct tgt gat gga gat gga gat gga gaa gca 738  
 Asp Cys Asp Arg Glu Gly Ala Cys Asp Gly Asp Gly Asp Gly Glu Ala  
 200 205 210

## MBI-20 Sequence Listing.ST25

```

gga gag ggg ctt atg gtt ccg gag atg tca gag aga ttg aaa tgg tca      786
Gly Glu Gly Leu Met Val Pro Glu Met Ser Glu Arg Leu Lys Trp Ser
215                      220                      225                      230

aga gat gtt gag gag atc aat ggt ggc gga gga gga ggt aac cag      834
Arg Asp Val Glu Glu Ile Asn Gly Gly Gly Gly Gly Val Asn Gln
235                      240                      245

cag tgg aat gct act act act aat cct agt ggt ggc cag agt tct cag      882
Gln Trp Asn Ala Thr Thr Thr Asn Pro Ser Gly Gly Gln Ser Ser Gln
250                      255                      260

ata tgg gat ttt aac ttg gga cag tca cgg gga cct gag gat acg agt      930
Ile Trp Asp Phe Asn Leu Gly Gln Ser Arg Gly Pro Glu Asp Thr Ser
265                      270                      275

cga gtg gaa gct gca tat gta ggg aaa ggt gct gct tct tca ttc aca      978
Arg Val Glu Ala Ala Tyr Val Gly Lys Gly Ala Ala Ser Ser Phe Thr
280                      285                      290

atc aac aat ttt gtt gac cat atg aat gaa act tgt tcc act aat gtg      1026
Ile Asn Asn Phe Val Asp His Met Asn Glu Thr Cys Ser Thr Asn Val
295                      300                      305                      310

aaa ggt gtc aaa gag att aaa aag gat gac tac aag cga tca act tca      1074
Lys Gly Val Lys Glu Ile Lys Lys Asp Asp Tyr Lys Arg Ser Thr Ser
315                      320                      325

ggc cag gta caa cca aca aaa tct gag agc aac aat cgt cca att acc      1122
Gly Gln Val Gln Pro Thr Lys Ser Glu Ser Asn Asn Arg Pro Ile Thr
330                      335                      340

ttt ggc tct gag aaa ggt tcg aac tcc tcc agt gac ttg cat ttc aca      1170
Phe Gly Ser Glu Lys Gly Ser Asn Ser Ser Ser Asp Leu His Phe Thr
345                      350                      355

gag cat att gct gga act agt tgt aag acc aca aga cta gtt gca act      1218
Glu His Ile Ala Gly Thr Ser Cys Lys Thr Thr Arg Leu Val Ala Thr
360                      365                      370

aag gct gat ctg gag cgg ctg gct cag aac aga gga gat gca atg cag      1266
Lys Ala Asp Leu Glu Arg Leu Ala Gln Asn Arg Gly Asp Ala Met Gln
375                      380                      385                      390

cgt tac aag gaa aag agg aag aca cgg aga tat gat aag acc ata agg      1314
Arg Tyr Lys Glu Lys Arg Lys Thr Arg Arg Tyr Asp Lys Thr Ile Arg
395                      400                      405

tat gaa tcg agg aag gca aga gct gac act agg ttg cgt gtc aga ggc      1362
Tyr Glu Ser Arg Lys Ala Arg Ala Asp Thr Arg Leu Arg Val Arg Gly
410                      415                      420

aga ttt gtg aaa gct agt gaa gct cct tac cct taa ccttaagttt      1408
Arg Phe Val Lys Ala Ser Glu Ala Pro Tyr Pro
425                      430

tttcacatag gcttcctttt agctacaaac ttagttactt tttttactcc actgcctcat      1468

aaatgtacag accggtctcg tttcatctgg ccgcccttct tgttttattg ccttatctgg      1528

cccttttatg taccttgga tcttatctag tttaaaaaag attgtaacct tctagaaaac      1588

catattctgt tgacagtata tacatgtcta tccaagcaaa aa                        1630

```

<210> 26  
 <211> 433  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 26

Met Ser Ser Ser Glu Arg Val Pro Cys Asp Phe Cys Gly Glu Arg Thr  
 1 5 10 15

## MBI-20 Sequence Listing.ST25

Ala Val Leu Phe Cys Arg Ala Asp Thr Ala Lys Leu Cys Leu Pro Cys  
                   20                                  25                                  30

Asp Gln Gln Val His Thr Ala Asn Leu Leu Ser Arg Lys His Val Arg  
                   35                                  40                                  45

Ser Gln Ile Cys Asp Asn Cys Gly Asn Glu Pro Val Ser Val Arg Cys  
           50                                  55                                  60

Phe Thr Asp Asn Leu Ile Leu Cys Gln Glu Cys Asp Trp Asp Val His  
   65                                  70                                  75                                  80

Gly Ser Cys Ser Val Ser Asp Ala His Val Arg Ser Ala Val Glu Gly  
                                   85                                  90                                  95

Phe Ser Gly Cys Pro Ser Ala Leu Glu Leu Ala Ala Leu Trp Gly Leu  
                                   100                                  105                                  110

Asp Leu Glu Gln Gly Arg Lys Asp Glu Glu Asn Gln Val Pro Met Met  
           115                                  120                                  125

Ala Met Met Met Asp Asn Phe Gly Met Gln Leu Asp Ser Trp Val Leu  
   130                                  135                                  140

Gly Ser Asn Glu Leu Ile Val Pro Ser Asp Thr Thr Phe Lys Lys Arg  
   145                                  150                                  155                                  160

Gly Ser Cys Gly Ser Ser Cys Gly Arg Tyr Lys Gln Val Leu Cys Lys  
                                   165                                  170                                  175

Gln Leu Glu Glu Leu Leu Lys Ser Gly Val Val Gly Gly Asp Gly Asp  
                                   180                                  185                                  190

Asp Gly Asp Arg Asp Arg Asp Cys Asp Arg Glu Gly Ala Cys Asp Gly  
           195                                  200                                  205

Asp Gly Asp Gly Glu Ala Gly Glu Gly Leu Met Val Pro Glu Met Ser  
   210                                  215                                  220

Glu Arg Leu Lys Trp Ser Arg Asp Val Glu Glu Ile Asn Gly Gly Gly  
   225                                  230                                  235                                  240

Gly Gly Gly Val Asn Gln Gln Trp Asn Ala Thr Thr Thr Asn Pro Ser  
                                   245                                  250                                  255

Gly Gly Gln Ser Ser Gln Ile Trp Asp Phe Asn Leu Gly Gln Ser Arg  
                                   260                                  265                                  270

Gly Pro Glu Asp Thr Ser Arg Val Glu Ala Ala Tyr Val Gly Lys Gly  
           275                                  280                                  285

Ala Ala Ser Ser Phe Thr Ile Asn Asn Phe Val Asp His Met Asn Glu  
   290                                  295                                  300

Thr Cys Ser Thr Asn Val Lys Gly Val Lys Glu Ile Lys Lys Asp Asp



305					310						315						320
Tyr	Lys	Arg	Ser	Thr	Ser	Gly	Gln	Val	Gln	Pro	Thr	Lys	Ser	Glu	Ser		
				325					330					335			
Asn	Asn	Arg	Pro	Ile	Thr	Phe	Gly	Ser	Glu	Lys	Gly	Ser	Asn	Ser	Ser		
			340					345					350				
Ser	Asp	Leu	His	Phe	Thr	Glu	His	Ile	Ala	Gly	Thr	Ser	Cys	Lys	Thr		
		355					360					365					
Thr	Arg	Leu	Val	Ala	Thr	Lys	Ala	Asp	Leu	Glu	Arg	Leu	Ala	Gln	Asn		
	370					375					380						
Arg	Gly	Asp	Ala	Met	Gln	Arg	Tyr	Lys	Glu	Lys	Arg	Lys	Thr	Arg	Arg		
385					390					395					400		
Tyr	Asp	Lys	Thr	Ile	Arg	Tyr	Glu	Ser	Arg	Lys	Ala	Arg	Ala	Asp	Thr		
				405					410					415			
Arg	Leu	Arg	Val	Arg	Gly	Arg	Phe	Val	Lys	Ala	Ser	Glu	Ala	Pro	Tyr		
			420					425					430				

```
<210> 27
<211> 768
<212> DNA
<213> Arabidopsis thaliana

<220>
<221> CDS
<222> (58)..(657)
<223> G975
```

<400>	27																	
attactcatc atcaagttcc tactttctct ctgacaaaca tcacagagta agtaaga																		57
atg gta cag acg aag aag ttc aga ggt gtc agg caa cgc cat tgg ggt	105																	
Met Val Gln Thr Lys Lys Phe Arg Gly Val Arg Gln Arg His Trp Gly																		
1 5 10 15																		
tct tgg gtc gct gag att cgt cat cct ctc ttg aaa cgg agg att tgg	153																	
Ser Trp Val Ala Glu Ile Arg His Pro Leu Leu Lys Arg Arg Ile Trp																		
20 25 30																		
cta ggg acg ttc gag acc gca gag gag gca gca aga gca tac gac gag	201																	
Leu Gly Thr Phe Glu Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Glu																		
35 40 45																		
gcc gcc gtt tta atg agc ggc cgc aac gcc aaa acc aac ttt ccc ctc	249																	
Ala Ala Val Leu Met Ser Gly Arg Asn Ala Lys Thr Asn Phe Pro Leu																		
50 55 60																		
aac aac aac aac acc gga gaa act tcc gag ggc aaa acc gat att tca	297																	
Asn Asn Asn Asn Thr Gly Glu Thr Ser Glu Gly Lys Thr Asp Ile Ser																		
65 70 75 80																		
gct tcg tcc aca atg tca tcc tca aca tca tct tca tcg ctc tct tcc	345																	
Ala Ser Ser Thr Met Ser Ser Ser Thr Ser Ser Ser Ser Leu Ser Ser																		
85 90 95																		
atc ctc agc gcc aaa ctg agg aaa tgc tgc aag tct cct tcc cca tcc	393																	
Ile Leu Ser Ala Lys Leu Arg Lys Cys Cys Lys Ser Pro Ser Pro Ser																		

## MBI-20 Sequence Listing.ST25

```

100          105          110
ctc acc tgc ctc cgt ctt gac aca gcc agc tcc cat atc ggc gtc tgg      441
Leu Thr Cys Leu Arg Leu Asp Thr Ala Ser Ser His Ile Gly Val Trp
115          120          125

cag aaa cgg gcc ggt tca aag tct gac tcc agc tgg gtc atg acg gtg      489
Gln Lys Arg Ala Gly Ser Lys Ser Asp Ser Ser Trp Val Met Thr Val
130          135          140

gag cta ggt ccc gca agc tcc tcc caa gag act act agt aaa gct tca      537
Glu Leu Gly Pro Ala Ser Ser Ser Gln Glu Thr Thr Ser Lys Ala Ser
145          150          155

caa gac gct att ctt gct ccg acc act gaa gtt gaa att ggt ggc agc      585
Gln Asp Ala Ile Leu Ala Pro Thr Thr Glu Val Glu Ile Gly Gly Ser
165          170          175

aga gaa gaa gta ttg gat gag gaa gaa aag gtt gct ttg caa atg ata      633
Arg Glu Glu Val Leu Asp Glu Glu Glu Lys Val Ala Leu Gln Met Ile
180          185          190

gag gag ctt ctc aat aca aac taa atcttatttg cttatatata tgtacctatt      687
Glu Glu Leu Leu Asn Thr Asn
195

ttcattgctg atttacagcc aaaataatca attataccgt gtattttata gatgtttat      747
attaaaagggt tgtagatat a      768

<210> 28
<211> 199
<212> PRT
<213> Arabidopsis thaliana

<400> 28
Met Val Gln Thr Lys Lys Phe Arg Gly Val Arg Gln Arg His Trp Gly
1      5      10      15

Ser Trp Val Ala Glu Ile Arg His Pro Leu Leu Lys Arg Arg Ile Trp
20      25      30

Leu Gly Thr Phe Glu Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Glu
35      40      45

Ala Ala Val Leu Met Ser Gly Arg Asn Ala Lys Thr Asn Phe Pro Leu
50      55      60

Asn Asn Asn Asn Thr Gly Glu Thr Ser Glu Gly Lys Thr Asp Ile Ser
65      70      75      80

Ala Ser Ser Thr Met Ser Ser Ser Thr Ser Ser Ser Ser Leu Ser Ser
85      90      95

Ile Leu Ser Ala Lys Leu Arg Lys Cys Cys Lys Ser Pro Ser Pro Ser
100     105     110

Leu Thr Cys Leu Arg Leu Asp Thr Ala Ser Ser His Ile Gly Val Trp
115     120     125

Gln Lys Arg Ala Gly Ser Lys Ser Asp Ser Ser Trp Val Met Thr Val
130     135     140

Glu Leu Gly Pro Ala Ser Ser Ser Gln Glu Thr Thr Ser Lys Ala Ser

```

## MBI-20 Sequence Listing.ST25

```

145                               150                               155                               160
Gln Asp Ala Ile Leu Ala Pro Thr Thr Glu Val Glu Ile Gly Gly Ser
      165                               170                               175

Arg Glu Glu Val Leu Asp Glu Glu Glu Lys Val Ala Leu Gln Met Ile
      180                               185                               190

Glu Glu Leu Leu Asn Thr Asn
      195

<210> 29
<211> 2526
<212> DNA
<213> Arabidopsis thaliana

<220>
<221> CDS
<222> (338)..(2275)
<223> G680

<400> 29
cagttatctt cttccttctt ctctctgttt tttaaattta ttttttagaga attttttttg      60
ttttgcttcc gatttgatta tttccgggaa cgatgacttc tccggggagt tcccggtgag      120
atgataagtc agattgcata cttgtctcct ccatggctac tctcaagggt tttggctgcg      180
gtggattcgt ttggtttctc tagaatctaa agaggttatc acaacggctt tgcaatttga      240
aaactttcat gtttggggag atcaaagatg gtttcttttt tatactttac ttgtagaga      300
ggatttgaag cagcgaatag ctgcaaccgg tcctggtt atg gat act aat aca tct      355
                               Met Asp Thr Asn Thr Ser
                               1                               5

gga gaa gaa tta tta gct aag gca aga aag cca tat aca ata aca aag      403
Gly Glu Glu Leu Leu Ala Lys Ala Arg Lys Pro Tyr Thr Ile Thr Lys
      10                               15                               20

cag cga gag cga tgg act gag gat gag cat gag agg ttt cta gaa gcc      451
Gln Arg Glu Arg Trp Thr Glu Asp Glu His Glu Arg Phe Leu Glu Ala
      25                               30                               35

ttg agg ctt tat gga aga gct tgg caa cga att gaa gaa cat att ggg      499
Leu Arg Leu Tyr Gly Arg Ala Trp Gln Arg Ile Glu Glu His Ile Gly
      40                               45                               50

aca aag act gct gtt cag atc aga agt cat gca caa aag ttc ttc aca      547
Thr Lys Thr Ala Val Gln Ile Arg Ser His Ala Gln Lys Phe Phe Thr
      55                               60                               65                               70

aag ttg gag aaa gag gct gaa gtt aaa ggc atc cct gtt tgc caa gct      595
Lys Leu Glu Lys Glu Ala Glu Val Lys Gly Ile Pro Val Cys Gln Ala
      75                               80                               85

ttg gac ata gaa att ccg cct cct cgt cct aaa cga aaa ccc aat act      643
Leu Asp Ile Glu Ile Pro Pro Pro Arg Pro Lys Arg Lys Pro Asn Thr
      90                               95                               100

cct tat cct cga aaa cct ggg aac aac ggt aca tct tcc tct caa gta      691
Pro Tyr Pro Arg Lys Pro Gly Asn Asn Gly Thr Ser Ser Ser Gln Val
      105                               110                               115

tca tca gca aaa gat gca aaa ctt gtt tca tcg gcc tct tct tca cag      739
Ser Ser Ala Lys Asp Ala Lys Leu Val Ser Ser Ala Ser Ser Ser Gln
      120                               125                               130

ttg aat cag gcg ttc ttg gat ttg gaa aaa atg ccg ttc tct gag aaa      787
Leu Asn Gln Ala Phe Leu Asp Leu Glu Lys Met Pro Phe Ser Glu Lys

```

MBI-20 Sequence Listing.ST25

135	140	145	150	
aca tca act gga aaa gaa aat caa gat gag aat tgc tcg ggt gtt tct				835
Thr Ser Thr Gly Lys Glu Asn Gln Asp Glu Asn Cys Ser Gly Val Ser				
	155	160	165	
act gtg aac aag tat ccc tta cca acg aaa cag gta agt ggc gac att				883
Thr Val Asn Lys Tyr Pro Leu Pro Thr Lys Gln Val Ser Gly Asp Ile				
	170	175	180	
gaa aca agt aag acc tca act gtg gac aac gcg gtt caa gat gtt ccc				931
Glu Thr Ser Lys Thr Ser Thr Val Asp Asn Ala Val Gln Asp Val Pro				
	185	190	195	
aag aag aac aaa gac aaa gat ggt aac gat ggt act act gtg cac agc				979
Lys Lys Asn Lys Asp Lys Asp Gly Asn Asp Gly Thr Thr Val His Ser				
	200	205	210	
atg caa aac tac cct tgg cat ttc cac gca gat att gtg aac ggg aat				1027
Met Gln Asn Tyr Pro Trp His Phe His Ala Asp Ile Val Asn Gly Asn				
	215	220	225	230
ata gca aaa tgc cct caa aat cat ccc tca ggt atg gta tct caa gac				1075
Ile Ala Lys Cys Pro Gln Asn His Pro Ser Gly Met Val Ser Gln Asp				
	235	240	245	
ttc atg ttt cat cct atg aga gaa gaa act cac ggg cac gca aat ctt				1123
Phe Met Phe His Pro Met Arg Glu Thr His Gly His Ala Asn Leu				
	250	255	260	
caa gct aca aca gca tct gct act act aca gct tct cat caa gcg ttt				1171
Gln Ala Thr Thr Ala Ser Ala Thr Thr Thr Ala Ser His Gln Ala Phe				
	265	270	275	
cca gct tgt cat tca cag gat gat tac cgt tcg ttt ctc cag ata tca				1219
Pro Ala Cys His Ser Gln Asp Asp Tyr Arg Ser Phe Leu Gln Ile Ser				
	280	285	290	
tct act ttc tcc aat ctt att atg tca act ctc cta cag aat cct gca				1267
Ser Thr Phe Ser Asn Leu Ile Met Ser Thr Leu Leu Gln Asn Pro Ala				
	295	300	305	310
gct cat gct gca gct aca ttc gct gct tcg gtc tgg cct tat gcg agt				1315
Ala His Ala Ala Thr Phe Ala Ala Ser Val Trp Pro Tyr Ala Ser				
	315	320	325	
gtc ggg aat tct ggt gat tca tca acc cca atg agc tct tct cct cca				1363
Val Gly Asn Ser Gly Asp Ser Ser Thr Pro Met Ser Ser Ser Pro Pro				
	330	335	340	
agt ata act gcc att gcc gct gct aca gta gct gct gca act gct tgg				1411
Ser Ile Thr Ala Ile Ala Ala Ala Thr Val Ala Ala Thr Ala Trp				
	345	350	355	
tgg gct tct cat gga ctt ctt cct gta tgc gct cca gct cca ata aca				1459
Trp Ala Ser His Gly Leu Leu Pro Val Cys Ala Pro Ala Pro Ile Thr				
	360	365	370	
tgt gtt cca ttc tca act gtt gca gtt cca act cca gca atg act gaa				1507
Cys Val Pro Phe Ser Thr Val Ala Val Pro Thr Pro Ala Met Thr Glu				
	375	380	385	390
atg gat acc gtt gaa aat act caa ccg ttt gag aaa caa aac aca gct				1555
Met Asp Thr Val Glu Asn Thr Gln Pro Phe Glu Lys Gln Asn Thr Ala				
	395	400	405	
ctg caa gat caa acc ttg gct tcg aaa tct cca gct tca tca tct gat				1603
Leu Gln Asp Gln Thr Leu Ala Ser Lys Ser Pro Ala Ser Ser Ser Asp				
	410	415	420	
gat tca gat gag act gga gta acc aag cta aat gcc gac tca aaa acc				1651
Asp Ser Asp Glu Thr Gly Val Thr Lys Leu Asn Ala Asp Ser Lys Thr				
	425	430	435	
aat gat gat aaa att gag gag gtt gtt gtt act gcc gct gtg cat gac				1699

## MBI-20 Sequence Listing.ST25

Asn	Asp	Asp	Lys	Ile	Glu	Glu	Val	Val	Val	Thr	Ala	Ala	Val	His	Asp	
440						445					450					
tca	aac	act	gcc	cag	aag	aaa	aat	ctt	gtg	gac	cgc	tca	tcg	tgt	ggc	1747
Ser	Asn	Thr	Ala	Gln	Lys	Lys	Asn	Leu	Val	Asp	Arg	Ser	Ser	Cys	Gly	
455					460					465					470	
tca	aat	aca	cct	tca	ggg	agt	gac	gca	gaa	act	gat	gca	tta	gat	aaa	1795
Ser	Asn	Thr	Pro	Ser	Gly	Ser	Asp	Ala	Glu	Thr	Asp	Ala	Leu	Asp	Lys	
				475					480					485		
atg	gag	aaa	gat	aaa	gag	gat	gtg	aag	gag	aca	gat	gag	aat	cag	cca	1843
Met	Glu	Lys	Asp	Lys	Glu	Asp	Val	Lys	Glu	Thr	Asp	Glu	Asn	Gln	Pro	
			490					495					500			
gat	gtt	att	gag	tta	aat	aac	cgt	aag	att	aaa	atg	aga	gac	aac	aac	1891
Asp	Val	Ile	Glu	Leu	Asn	Asn	Arg	Lys	Ile	Lys	Met	Arg	Asp	Asn	Asn	
		505					510					515				
agc	aac	aac	aat	gca	act	act	gat	tcg	tgg	aag	gaa	gtc	tcc	gaa	gag	1939
Ser	Asn	Asn	Asn	Ala	Thr	Thr	Asp	Ser	Trp	Lys	Glu	Val	Ser	Glu	Glu	
			520			525					530					
ggc	cgt	ata	gag	ttt	cag	gct	ctc	ttt	gca	aga	gaa	aga	ttg	cct	caa	1987
Gly	Arg	Ile	Ala	Phe	Gln	Ala	Leu	Phe	Ala	Arg	Glu	Arg	Leu	Pro	Gln	
535					540				545					550		
agc	ttt	tcg	cct	cct	caa	gtg	gca	gag	aat	gtg	aat	aga	aaa	caa	agt	2035
Ser	Phe	Ser	Pro	Pro	Gln	Val	Ala	Glu	Asn	Val	Asn	Arg	Lys	Gln	Ser	
				555					560					565		
gac	acg	tca	atg	cca	ttg	gct	cct	aat	ttc	aaa	agc	cag	gat	tct	tgt	2083
Asp	Thr	Ser	Met	Pro	Leu	Ala	Pro	Asn	Phe	Lys	Ser	Gln	Asp	Ser	Cys	
			570					575					580			
gct	gca	gac	caa	gaa	gga	gta	gta	atg	atc	ggc	gtt	gga	aca	tgc	aag	2131
Ala	Ala	Asp	Gln	Glu	Gly	Val	Val	Met	Ile	Gly	Val	Gly	Thr	Cys	Lys	
		585					590					595				
agt	ctt	aaa	acg	aga	cag	aca	gga	ttt	aag	cca	tac	aag	aga	tgt	tca	2179
Ser	Leu	Lys	Thr	Arg	Gln	Thr	Gly	Phe	Lys	Pro	Tyr	Lys	Arg	Cys	Ser	
	600					605					610					
atg	gaa	gtg	aaa	gag	agc	caa	gtt	ggg	aac	ata	aac	aat	caa	agt	gat	2227
Met	Glu	Val	Lys	Glu	Ser	Gln	Val	Gly	Asn	Ile	Asn	Asn	Gln	Ser	Asp	
615					620					625					630	
gaa	aaa	gtc	tgc	aaa	agg	ctt	cga	ttg	gaa	gga	gaa	gct	tct	aca	tga	2275
Glu	Lys	Val	Cys	Lys	Arg	Leu	Arg	Leu	Glu	Gly	Glu	Ala	Ser	Thr		
				635					640					645		
cagacttgga	ggtaaaaaaa	aaacatccac	attttttatca	atatcttttaa	atctagtgtt											2335
agtagtttgc	ttctccaatc	tttatgaaag	agacttttaa	ttttccttcc	gaacatttct											2395
ttggctcatgt	cagggtctgt	accatattac	cccatgtctt	gtctcttctg	tctgtttgtg											2455
tatgctactt	gtggctctata	tgctcatctgc	tactactgtt	aattaacat	taagcaatgg											2515
atttgccttt	a															2526
<210>	30															
<211>	645															
<212>	PRT															
<213>	Arabidopsis thaliana															
<400>	30															
Met	Asp	Thr	Asn	Thr	Ser	Gly	Glu	Glu	Leu	Leu	Ala	Lys	Ala	Arg	Lys	
1				5					10					15		
Pro	Tyr	Thr	Ile	Thr	Lys	Gln	Arg	Glu	Arg	Trp	Thr	Glu	Asp	Glu	His	
			20					25					30			

## MBI-20 Sequence Listing.ST25

Glu Arg Phe Leu Glu Ala Leu Arg Leu Tyr Gly Arg Ala Trp Gln Arg  
 35 40 45  
 Ile Glu Glu His Ile Gly Thr Lys Thr Ala Val Gln Ile Arg Ser His  
 50 55 60  
 Ala Gln Lys Phe Phe Thr Lys Leu Glu Lys Glu Ala Glu Val Lys Gly  
 65 70 75 80  
 Ile Pro Val Cys Gln Ala Leu Asp Ile Glu Ile Pro Pro Pro Arg Pro  
 85 90 95  
 Lys Arg Lys Pro Asn Thr Pro Tyr Pro Arg Lys Pro Gly Asn Asn Gly  
 100 105 110  
 Thr Ser Ser Ser Gln Val Ser Ser Ala Lys Asp Ala Lys Leu Val Ser  
 115 120 125  
 Ser Ala Ser Ser Ser Gln Leu Asn Gln Ala Phe Leu Asp Leu Glu Lys  
 130 135 140  
 Met Pro Phe Ser Glu Lys Thr Ser Thr Gly Lys Glu Asn Gln Asp Glu  
 145 150 155 160  
 Asn Cys Ser Gly Val Ser Thr Val Asn Lys Tyr Pro Leu Pro Thr Lys  
 165 170 175  
 Gln Val Ser Gly Asp Ile Glu Thr Ser Lys Thr Ser Thr Val Asp Asn  
 180 185 190  
 Ala Val Gln Asp Val Pro Lys Lys Asn Lys Asp Lys Asp Gly Asn Asp  
 195 200 205  
 Gly Thr Thr Val His Ser Met Gln Asn Tyr Pro Trp His Phe His Ala  
 210 215 220  
 Asp Ile Val Asn Gly Asn Ile Ala Lys Cys Pro Gln Asn His Pro Ser  
 225 230 235 240  
 Gly Met Val Ser Gln Asp Phe Met Phe His Pro Met Arg Glu Glu Thr  
 245 250 255  
 His Gly His Ala Asn Leu Gln Ala Thr Thr Ala Ser Ala Thr Thr Thr  
 260 265 270  
 Ala Ser His Gln Ala Phe Pro Ala Cys His Ser Gln Asp Asp Tyr Arg  
 275 280 285  
 Ser Phe Leu Gln Ile Ser Ser Thr Phe Ser Asn Leu Ile Met Ser Thr  
 290 295 300  
 Leu Leu Gln Asn Pro Ala Ala His Ala Ala Ala Thr Phe Ala Ala Ser  
 305 310 315 320  
 Val Trp Pro Tyr Ala Ser Val Gly Asn Ser Gly Asp Ser Ser Thr Pro

## MBI-20 Sequence Listing.ST25

325

330

335

Met Ser Ser Ser Pro Pro Ser Ile Thr Ala Ile Ala Ala Thr Val  
 340 345 350

Ala Ala Ala Thr Ala Trp Trp Ala Ser His Gly Leu Leu Pro Val Cys  
 355 360 365

Ala Pro Ala Pro Ile Thr Cys Val Pro Phe Ser Thr Val Ala Val Pro  
 370 375 380

Thr Pro Ala Met Thr Glu Met Asp Thr Val Glu Asn Thr Gln Pro Phe  
 385 390 395 400

Glu Lys Gln Asn Thr Ala Leu Gln Asp Gln Thr Leu Ala Ser Lys Ser  
 405 410 415

Pro Ala Ser Ser Ser Asp Asp Ser Asp Glu Thr Gly Val Thr Lys Leu  
 420 425 430

Asn Ala Asp Ser Lys Thr Asn Asp Asp Lys Ile Glu Glu Val Val Val  
 435 440 445

Thr Ala Ala Val His Asp Ser Asn Thr Ala Gln Lys Lys Asn Leu Val  
 450 455 460

Asp Arg Ser Ser Cys Gly Ser Asn Thr Pro Ser Gly Ser Asp Ala Glu  
 465 470 475 480

Thr Asp Ala Leu Asp Lys Met Glu Lys Asp Lys Glu Asp Val Lys Glu  
 485 490 495

Thr Asp Glu Asn Gln Pro Asp Val Ile Glu Leu Asn Asn Arg Lys Ile  
 500 505 510

Lys Met Arg Asp Asn Asn Ser Asn Asn Asn Ala Thr Thr Asp Ser Trp  
 515 520 525

Lys Glu Val Ser Glu Glu Gly Arg Ile Ala Phe Gln Ala Leu Phe Ala  
 530 535 540

Arg Glu Arg Leu Pro Gln Ser Phe Ser Pro Pro Gln Val Ala Glu Asn  
 545 550 555 560

Val Asn Arg Lys Gln Ser Asp Thr Ser Met Pro Leu Ala Pro Asn Phe  
 565 570 575

Lys Ser Gln Asp Ser Cys Ala Ala Asp Gln Glu Gly Val Val Met Ile  
 580 585 590

Gly Val Gly Thr Cys Lys Ser Leu Lys Thr Arg Gln Thr Gly Phe Lys  
 595 600 605

Pro Tyr Lys Arg Cys Ser Met Glu Val Lys Glu Ser Gln Val Gly Asn  
 610 615 620

## MBI-20 Sequence Listing.ST25

Ile Asn Asn Gln Ser Asp Glu Lys Val Cys Lys Arg Leu Arg Leu Glu  
 625 630 635 640

Gly Glu Ala Ser Thr  
 645

<210> 31  
 <211> 1195  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (67)..(1041)  
 <223> G883

<400> 31  
 ctctctcgtc ttcgtcttct tcttcttcaa cgctcctctc caaaatcctc agaccaagaa 60  
 atcatc atg gcc gtc gat cta atg cgt ttc cct aag ata gat gat caa 108  
 Met Ala Val Asp Leu Met Arg Phe Pro Lys Ile Asp Asp Gln  
 1 5 10  
 acg gct att cag gaa gct gca tcg caa ggt tta caa agt atg gaa cat 156  
 Thr Ala Ile Gln Glu Ala Ala Ser Gln Gly Leu Gln Ser Met Glu His  
 15 20 25 30  
 ctg atc cgt gtc ctc tct aac cgt ccc gaa caa caa cac aac gtt gac 204  
 Leu Ile Arg Val Ser Asn Arg Pro Glu Gln Gln His Asn Val Asp  
 35 40 45  
 tgc tcc gag atc act gac ttc acc gtt tct aaa ttc aaa acc gtc att 252  
 Cys Ser Glu Ile Thr Asp Phe Thr Val Ser Lys Phe Lys Thr Val Ile  
 50 55 60  
 tct ctc ctt aac cgt act ggt cac gct cgg ttc aga cgc gga ccg gtt 300  
 Ser Leu Leu Asn Arg Thr Gly His Ala Arg Phe Arg Arg Gly Pro Val  
 65 70 75  
 cac tcc act tcc tct gcc gca tct cag aaa cta cag agt cag atc gtt 348  
 His Ser Thr Ser Ser Ala Ala Ser Gln Lys Leu Gln Ser Gln Ile Val  
 80 85 90  
 aaa aat act caa cct gag gct ccg ata gtg aga aca act acg aat cac 396  
 Lys Asn Thr Gln Pro Glu Ala Pro Ile Val Arg Thr Thr Thr Asn His  
 95 100 105 110  
 cct caa atc gtt cct cca ccg tct agt gta aca ctc gat ttc tct aaa 444  
 Pro Gln Ile Val Pro Pro Pro Ser Ser Val Thr Leu Asp Phe Ser Lys  
 115 120 125  
 cca agc atc ttc ggc acc aaa gct aag agc gcc gag ctg gaa ttc tcc 492  
 Pro Ser Ile Phe Gly Thr Lys Ala Lys Ser Ala Glu Leu Glu Phe Ser  
 130 135 140  
 aaa gaa aac ttc agt gtt tct tta aac tcc tca ttc atg tcg tcg gcg 540  
 Lys Glu Asn Phe Ser Val Ser Leu Asn Ser Ser Phe Met Ser Ser Ala  
 145 150 155  
 ata acc gga gac ggc agc gtc tcc aat gga aaa atc ttc ctt gct tct 588  
 Ile Thr Gly Asp Gly Ser Val Ser Asn Gly Lys Ile Phe Leu Ala Ser  
 160 165 170  
 gct ccg tcg cag cct gtt aac tct tcc gga aaa cca ccg ttg gct ggt 636  
 Ala Pro Ser Gln Pro Val Asn Ser Ser Gly Lys Pro Pro Leu Ala Gly  
 175 180 185 190  
 cat cct tac aga aag aga tgt ctc gag cat gag cac tca gag agt ttc 684  
 His Pro Tyr Arg Lys Arg Cys Leu Glu His Glu His Ser Glu Ser Phe  
 195 200 205  
 tcc gga aaa gtc tcc ggc tcc gcc tac gga aag tgc cat tgc aag aaa 732



## MBI-20 Sequence Listing.ST25

Ser Gly Lys Val Ser Gly Ser Ala Tyr Gly Lys Cys His Cys Lys Lys  
 210 215 220

agg aaa aat cgg atg aag aga acc gtg aga gta ccg gcg ata agt gca 780  
 Arg Lys Asn Arg Met Lys Arg Thr Val Arg Val Pro Ala Ile Ser Ala  
 225 230 235

aag atc gcc gat att cca ccg gac gaa tat tcg tgg agg aag tac gga 828  
 Lys Ile Ala Asp Ile Pro Pro Asp Glu Tyr Ser Trp Arg Lys Tyr Gly  
 240 245 250

caa aaa ccg atc aag ggc tca cca cac cca cgt ggt tac tac aag tgc 876  
 Gln Lys Pro Ile Lys Gly Ser Pro His Pro Arg Gly Tyr Tyr Lys Cys  
 255 260 265 270

agt aca ttc aga gga tgt cca gcg agg aaa cac gtg gaa cga gca tta 924  
 Ser Thr Phe Arg Gly Cys Pro Ala Arg Lys His Val Glu Arg Ala Leu  
 275 280 285

gat gat cca gcg atg ctt att gtg aca tac gaa gga gag cac cgt cat 972  
 Asp Asp Pro Ala Met Leu Ile Val Thr Tyr Glu Gly Glu His Arg His  
 290 295 300

aac caa tcc gcg atg cag gag aat att tct tct tca ggc att aat gat 1020  
 Asn Gln Ser Ala Met Gln Glu Asn Ile Ser Ser Ser Gly Ile Asn Asp  
 305 310 315

tta gtg ttt gcc tcg gct tga cttttttttg tactatttgt tttttgattt 1071  
 Leu Val Phe Ala Ser Ala  
 320

tttgagtact ttagatggat tgaaatttgt aaattttttt attaagaaat caatttaaat 1131

agagaaaaat tagtggtggt gcaaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa 1191

aaaa 1195

<210> 32  
 <211> 324  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 32

Met Ala Val Asp Leu Met Arg Phe Pro Lys Ile Asp Asp Gln Thr Ala  
 1 5 10 15

Ile Gln Glu Ala Ala Ser Gln Gly Leu Gln Ser Met Glu His Leu Ile  
 20 25 30

Arg Val Leu Ser Asn Arg Pro Glu Gln Gln His Asn Val Asp Cys Ser  
 35 40 45

Glu Ile Thr Asp Phe Thr Val Ser Lys Phe Lys Thr Val Ile Ser Leu  
 50 55 60

Leu Asn Arg Thr Gly His Ala Arg Phe Arg Arg Gly Pro Val His Ser  
 65 70 75 80

Thr Ser Ser Ala Ala Ser Gln Lys Leu Gln Ser Gln Ile Val Lys Asn  
 85 90 95

Thr Gln Pro Glu Ala Pro Ile Val Arg Thr Thr Thr Asn His Pro Gln  
 100 105 110

Ile Val Pro Pro Pro Ser Ser Val Thr Leu Asp Phe Ser Lys Pro Ser  
 115 120 125

## MBI-20 Sequence Listing.ST25

Ile Phe Gly Thr Lys Ala Lys Ser Ala Glu Leu Glu Phe Ser Lys Glu  
130 135 140

Asn Phe Ser Val Ser Leu Asn Ser Ser Phe Met Ser Ser Ala Ile Thr  
145 150 155 160

Gly Asp Gly Ser Val Ser Asn Gly Lys Ile Phe Leu Ala Ser Ala Pro  
165 170 175

Ser Gln Pro Val Asn Ser Ser Gly Lys Pro Pro Leu Ala Gly His Pro  
180 185 190

Tyr Arg Lys Arg Cys Leu Glu His Glu His Ser Glu Ser Phe Ser Gly  
195 200 205

Lys Val Ser Gly Ser Ala Tyr Gly Lys Cys His Cys Lys Lys Arg Lys  
210 215 220

Asn Arg Met Lys Arg Thr Val Arg Val Pro Ala Ile Ser Ala Lys Ile  
225 230 235 240

Ala Asp Ile Pro Pro Asp Glu Tyr Ser Trp Arg Lys Tyr Gly Gln Lys  
245 250 255

Pro Ile Lys Gly Ser Pro His Pro Arg Gly Tyr Tyr Lys Cys Ser Thr  
260 265 270

Phe Arg Gly Cys Pro Ala Arg Lys His Val Glu Arg Ala Leu Asp Asp  
275 280 285

Pro Ala Met Leu Ile Val Thr Tyr Glu Gly Glu His Arg His Asn Gln  
290 295 300

Ser Ala Met Gln Glu Asn Ile Ser Ser Ser Gly Ile Asn Asp Leu Val  
305 310 315 320

Phe Ala Ser Ala

<210> 33  
<211> 1902  
<212> DNA  
<213> Arabidopsis thaliana

<220>  
<221> CDS  
<222> (1)..(1902)  
<223> G1855

<400> 33  
atg gcg aaa gag aac agt ggt cat cat cac caa aca gaa gca aga aga 48  
Met Ala Lys Glu Asn Ser Gly His His His Gln Thr Glu Ala Arg Arg  
1 5 10 15  
aag aaa cta act ttg att ctt ggt gta agt gga ctc tgc att ttg ttc 96  
Lys Lys Leu Thr Leu Ile Leu Gly Val Ser Gly Leu Cys Ile Leu Phe  
20 25 30  
tat gtt tta ggt gca tgg caa gcc aat acc gtc cca tct tct atc tcg 144

MBI-20 Sequence Listing.ST25																
Tyr	Val	Leu	Gly	Ala	Trp	Gln	Ala	Asn	Thr	Val	Pro	Ser	Ser	Ile	Ser	
	35						40					45				
aag	ctc	gga	tgc	gag	acg	caa	tca	aac	cct	tct	tcg	tcc	tct	tcc	tct	192
Lys	Leu	Gly	Cys	Glu	Thr	Gln	Ser	Asn	Pro	Ser	Ser	Ser	Ser	Ser	Ser	
	50					55					60					
tcc	tca	tct	tca	gag	tca	gct	gaa	cta	gat	ttc	aaa	agc	cat	aat	cag	240
Ser	Ser	Ser	Ser	Glu	Ser	Ala	Glu	Leu	Asp	Phe	Lys	Ser	His	Asn	Gln	
	65				70					75					80	
att	gag	tta	aag	gaa	aca	aac	caa	acc	att	aag	tac	ttt	gaa	cca	tgt	288
Ile	Glu	Leu	Lys		Thr	Asn	Gln	Thr	Ile	Lys	Tyr	Phe	Glu	Pro	Cys	
				85					90					95		
gaa	tta	tct	ctc	agt	gag	tac	act	cct	tgt	gaa	gac	cga	caa	aga	gga	336
Glu	Leu	Ser	Leu	Ser	Glu	Tyr	Thr	Pro	Cys	Glu	Asp	Arg	Gln	Arg	Gly	
			100					105					110			
aga	aga	ttc	gat	agg	aac	atg	atg	aaa	tat	aga	gaa	aga	cat	tgt	cct	384
Arg	Arg	Phe	Asp	Arg	Asn	Met	Met	Lys	Tyr	Arg	Glu	Arg	His	Cys	Pro	
		115				120						125				
gta	aaa	gat	gag	ctt	ctt	tat	tgt	ttg	att	cct	cct	cca	cca	aac	tac	432
Val	Lys	Asp	Glu	Leu	Leu	Tyr	Cys	Leu	Ile	Pro	Pro	Pro	Pro	Asn	Tyr	
	130					135					140					
aag	att	cca	ttt	aaa	tgg	cca	caa	agt	aga	gac	tat	gct	tgg	tat	gac	480
Lys	Ile	Pro	Phe	Lys		Pro	Gln	Ser	Arg	Asp	Tyr	Ala	Trp	Tyr	Asp	
	145				150					155					160	
aat	atc	cct	cac	aag	gaa	ctt	agt	gtt	gag	aaa	gca	gtt	caa	aac	tgg	528
Asn	Ile	Pro	His	Lys	Glu	Leu	Ser	Val	Glu	Lys	Ala	Val	Gln	Asn	Trp	
				165					170					175		
att	caa	gtt	gaa	ggt	gac	cgc	ttt	aga	ttc	cct	ggt	ggt	ggt	act	atg	576
Ile	Gln	Val	Glu	Gly	Asp	Arg	Phe	Arg	Phe	Pro	Gly	Gly	Gly	Thr	Met	
			180				185						190			
ttt	cct	cgt	gga	gct	gat	gct	tat	atc	gat	gat	att	gct	agg	ctt	att	624
Phe	Pro	Arg	Gly	Ala	Asp	Ala	Tyr	Ile	Asp	Asp	Ile	Ala	Arg	Leu	Ile	
		195					200					205				
cct	ctt	act	gat	ggt	gga	atc	aga	aca	gct	att	gac	act	gga	tgt	ggt	672
Pro	Leu	Thr	Asp	Gly	Gly	Ile	Arg	Thr	Ala	Ile	Asp	Thr	Gly	Cys	Gly	
		210				215					220					
gtt	gca	agt	ttt	ggt	gct	tac	ctc	ttg	aag	aga	gac	att	atg	gct	gtg	720
Val	Ala	Ser	Phe	Gly	Ala	Tyr	Leu	Leu	Lys	Arg	Asp	Ile	Met	Ala	Val	
				230						235				240		
tct	ttt	gct	cca	aga	gac	act	cat	gaa	gct	cag	gta	cag	ttt	gct	tta	768
Ser	Phe	Ala	Pro	Arg	Asp	Thr	His	Glu	Ala	Gln	Val	Gln	Phe	Ala	Leu	
				245					250					255		
gaa	cgc	gga	gtt	cct	gcg	ata	atc	ggg	att	atg	gga	tca	aga	aga	ctt	816
Glu	Arg	Gly	Val	Pro	Ala	Ile	Ile	Gly	Ile	Met	Gly	Ser	Arg	Arg	Leu	
			260					265					270			
cct	tat	cca	gct	aga	gct	ttt	gat	ctt	gct	cat	tgt	tct	cgt	tgt	ttg	864
Pro	Tyr	Pro	Ala	Arg	Ala	Phe	Asp	Leu	Ala	His	Cys	Ser	Arg	Cys	Leu	
		275					280					285				
atc	cct	tgg	ttt	aaa	aat	gat	ggt	ttg	tac	ctt	atg	gag	gtc	gac	cgg	912
Ile	Pro	Trp	Phe	Lys	Asn	Asp	Gly	Leu	Tyr	Leu	Met	Glu	Val	Asp	Arg	
		290				295					300					
gtt	tta	aga	ccg	ggc	ggt	tac	tgg	atc	ctc	tcg	gga	cca	ccg	att	aac	960
Val	Leu	Arg	Pro	Gly	Gly	Tyr	Trp	Ile	Leu	Ser	Gly	Pro	Pro	Ile	Asn	
				310						315				320		
tgg	aaa	cag	tac	tgg	aga	ggg	tgg	gag	aga	aca	gag	gag	gat	ttg	aag	1008
Trp	Lys	Gln	Tyr	Trp	Arg	Gly	Trp	Glu	Arg	Thr	Glu	Glu	Asp	Leu	Lys	
				325					330					335		

## MBI-20 Sequence Listing.ST25

aaa gag caa gat tca ata gaa gat gta gca aag agt ctt tgc tgg aag Lys Glu Gln Asp Ser Ile Glu Asp Val Ala Lys Ser Leu Cys Trp Lys 340 345 350	1056
aaa gta act gaa aaa ggt gac tta tca att tgg caa aag cct ctc aat Lys Val Thr Glu Lys Gly Asp Leu Ser Ile Trp Gln Lys Pro Leu Asn 355 360 365	1104
cac att gag tgt aaa aag ctc aaa caa aac aat aag tca cct ccg ata His Ile Glu Cys Lys Lys Leu Lys Gln Asn Asn Lys Ser Pro Pro Ile 370 375 380	1152
tgc agc tca gat aac gcg gat tcc gct tgg tac aaa gac ttg gaa act Cys Ser Ser Asp Asn Ala Asp Ser Ala Trp Tyr Lys Asp Leu Glu Thr 385 390 395 400	1200
tgt ata aca cca tta cca gaa aca aac aat cca gat gat tca gca ggc Cys Ile Thr Pro Leu Pro Glu Thr Asn Asn Pro Asp Asp Ser Ala Gly 405 410 415	1248
ggg gca ctc gag gat tgg cca gac cga gca ttc gcg gta cct cca aga Gly Ala Leu Glu Asp Trp Pro Asp Arg Ala Phe Ala Val Pro Pro Arg 420 425 430	1296
atc atc aga gga act ata cca gaa atg aac gcg gag aaa ttt aga gaa Ile Ile Arg Gly Thr Ile Pro Glu Met Asn Ala Glu Lys Phe Arg Glu 435 440 445	1344
gac aac gag gtt tgg aaa gag aga ata gca cat tac aag aag ata gtc Asp Asn Glu Val Trp Lys Glu Arg Ile Ala His Tyr Lys Lys Ile Val 450 455 460	1392
cct gag ctt tca cat gga aga ttc agg aac att atg gac atg aac gct Pro Glu Leu Ser His Gly Arg Phe Arg Asn Ile Met Asp Met Asn Ala 465 470 475 480	1440
ttt ctc ggc gga ttc gct gct tcc atg ctg aaa tat ccc tca tgg gtc Phe Leu Gly Gly Phe Ala Ala Ser Met Leu Lys Tyr Pro Ser Trp Val 485 490 495	1488
atg aac gtt gtc ccg gtc gat gca gag aaa caa acg tta ggt gtg atc Met Asn Val Val Pro Val Asp Ala Glu Lys Gln Thr Leu Gly Val Ile 500 505 510	1536
tac gaa cgt gga ttg ata ggg acg tat caa gat tgg tgt gaa gga ttc Tyr Glu Arg Gly Leu Ile Gly Thr Tyr Gln Asp Trp Cys Glu Gly Phe 515 520 525	1584
tca acg tat cca aga act tat gat atg att cat gca gga gga ttg ttc Ser Thr Tyr Pro Arg Thr Tyr Asp Met Ile His Ala Gly Gly Leu Phe 530 535 540	1632
agc tta tac gaa cat agg tgt gat ttg acg ttg ata ttg ttg gag atg Ser Leu Tyr Glu His Arg Cys Asp Leu Thr Leu Ile Leu Leu Glu Met 545 550 555 560	1680
gat cga att ttg aga cca gaa gga aca gtt gtg ttg aga gat aat gtg Asp Arg Ile Leu Arg Pro Glu Gly Thr Val Val Leu Arg Asp Asn Val 565 570 575	1728
gag acg ttg aat aag gta gag aag ata gtg aag gga atg aag tgg aag Glu Thr Leu Asn Lys Val Glu Lys Ile Val Lys Gly Met Lys Trp Lys 580 585 590	1776
agt caa att gtt gat cat gag aaa ggt cct ttt aat cct gag aag att Ser Gln Ile Val Asp His Glu Lys Gly Pro Phe Asn Pro Glu Lys Ile 595 600 605	1824
ctt gtt gct gtt aaa act tat tgg act ggt caa cct tct gac aag aac Leu Val Ala Val Lys Thr Tyr Trp Thr Gly Gln Pro Ser Asp Lys Asn 610 615 620	1872
aac aac aac aac aac aac aac aac aac tag Asn Asn Asn Asn Asn Asn Asn Asn Asn 625 630	1902

## MBI-20 Sequence Listing.ST25

<210> 34  
 <211> 633  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 34

Met Ala Lys Glu Asn Ser Gly His His His Gln Thr Glu Ala Arg Arg  
 1 5 10 15

Lys Lys Leu Thr Leu Ile Leu Gly Val Ser Gly Leu Cys Ile Leu Phe  
 20 25 30

Tyr Val Leu Gly Ala Trp Gln Ala Asn Thr Val Pro Ser Ser Ile Ser  
 35 40 45

Lys Leu Gly Cys Glu Thr Gln Ser Asn Pro Ser Ser Ser Ser Ser  
 50 55 60

Ser Ser Ser Ser Glu Ser Ala Glu Leu Asp Phe Lys Ser His Asn Gln  
 65 70 75 80

Ile Glu Leu Lys Glu Thr Asn Gln Thr Ile Lys Tyr Phe Glu Pro Cys  
 85 90 95

Glu Leu Ser Leu Ser Glu Tyr Thr Pro Cys Glu Asp Arg Gln Arg Gly  
 100 105 110

Arg Arg Phe Asp Arg Asn Met Met Lys Tyr Arg Glu Arg His Cys Pro  
 115 120 125

Val Lys Asp Glu Leu Leu Tyr Cys Leu Ile Pro Pro Pro Pro Asn Tyr  
 130 135 140

Lys Ile Pro Phe Lys Trp Pro Gln Ser Arg Asp Tyr Ala Trp Tyr Asp  
 145 150 155 160

Asn Ile Pro His Lys Glu Leu Ser Val Glu Lys Ala Val Gln Asn Trp  
 165 170 175

Ile Gln Val Glu Gly Asp Arg Phe Arg Phe Pro Gly Gly Gly Thr Met  
 180 185 190

Phe Pro Arg Gly Ala Asp Ala Tyr Ile Asp Asp Ile Ala Arg Leu Ile  
 195 200 205

Pro Leu Thr Asp Gly Gly Ile Arg Thr Ala Ile Asp Thr Gly Cys Gly  
 210 215 220

Val Ala Ser Phe Gly Ala Tyr Leu Leu Lys Arg Asp Ile Met Ala Val  
 225 230 235 240

Ser Phe Ala Pro Arg Asp Thr His Glu Ala Gln Val Gln Phe Ala Leu  
 245 250 255

Glu Arg Gly Val Pro Ala Ile Ile Gly Ile Met Gly Ser Arg Arg Leu  
 260 265 270

## MBI-20 Sequence Listing.ST25

Pro Tyr Pro Ala Arg Ala Phe Asp Leu Ala His Cys Ser Arg Cys Leu  
 275 280 285  
 Ile Pro Trp Phe Lys Asn Asp Gly Leu Tyr Leu Met Glu Val Asp Arg  
 290 295 300  
 Val Leu Arg Pro Gly Gly Tyr Trp Ile Leu Ser Gly Pro Pro Ile Asn  
 305 310 315 320  
 Trp Lys Gln Tyr Trp Arg Gly Trp Glu Arg Thr Glu Glu Asp Leu Lys  
 325 330 335  
 Lys Glu Gln Asp Ser Ile Glu Asp Val Ala Lys Ser Leu Cys Trp Lys  
 340 345 350  
 Lys Val Thr Glu Lys Gly Asp Leu Ser Ile Trp Gln Lys Pro Leu Asn  
 355 360 365  
 His Ile Glu Cys Lys Lys Leu Lys Gln Asn Asn Lys Ser Pro Pro Ile  
 370 375 380  
 Cys Ser Ser Asp Asn Ala Asp Ser Ala Trp Tyr Lys Asp Leu Glu Thr  
 385 390 395 400  
 Cys Ile Thr Pro Leu Pro Glu Thr Asn Asn Pro Asp Asp Ser Ala Gly  
 405 410 415  
 Gly Ala Leu Glu Asp Trp Pro Asp Arg Ala Phe Ala Val Pro Pro Arg  
 420 425 430  
 Ile Ile Arg Gly Thr Ile Pro Glu Met Asn Ala Glu Lys Phe Arg Glu  
 435 440 445  
 Asp Asn Glu Val Trp Lys Glu Arg Ile Ala His Tyr Lys Lys Ile Val  
 450 455 460  
 Pro Glu Leu Ser His Gly Arg Phe Arg Asn Ile Met Asp Met Asn Ala  
 465 470 475 480  
 Phe Leu Gly Gly Phe Ala Ala Ser Met Leu Lys Tyr Pro Ser Trp Val  
 485 490 495  
 Met Asn Val Val Pro Val Asp Ala Glu Lys Gln Thr Leu Gly Val Ile  
 500 505 510  
 Tyr Glu Arg Gly Leu Ile Gly Thr Tyr Gln Asp Trp Cys Glu Gly Phe  
 515 520 525  
 Ser Thr Tyr Pro Arg Thr Tyr Asp Met Ile His Ala Gly Gly Leu Phe  
 530 535 540  
 Ser Leu Tyr Glu His Arg Cys Asp Leu Thr Leu Ile Leu Leu Glu Met  
 545 550 555 560  
 Asp Arg Ile Leu Arg Pro Glu Gly Thr Val Val Leu Arg Asp Asn Val

## MBI-20 Sequence Listing.ST25

565

570

575

Glu Thr Leu Asn Lys Val Glu Lys Ile Val Lys Gly Met Lys Trp Lys  
580 585 590

Ser Gln Ile Val Asp His Glu Lys Gly Pro Phe Asn Pro Glu Lys Ile  
595 600 605

Leu Val Ala Val Lys Thr Tyr Trp Thr Gly Gln Pro Ser Asp Lys Asn  
610 615 620

Asn Asn Asn Asn Asn Asn Asn Asn Asn  
625 630

<210> 35  
<211> 2324  
<212> DNA  
<213> Arabidopsis thaliana

<220>  
<221> CDS  
<222> (209)..(2020)  
<223> G1190

<400> 35  
tctctgtccca aaaccaaag gcttgagagt gtgtctttag agagagatct tctctctttt 60  
atcttacgac tctcacttct tatctcaaat ctacttcaac tctatttcca gtctccacat 120  
tttccacaaa atttcaactc ttgttctctt catccaaagt aaaaaacaaa tcgttgcaag 180  
tgaggttttg ttttggtgtt atagaatt atg aag agc ggg aag caa tct tcg 232  
Met Lys Ser Gly Lys Gln Ser Ser  
1 5  
caa cct gaa aag ggt act tcc agg atc ttg tca ctg act gtc ctg ttt 280  
Gln Pro Glu Lys Gly Thr Ser Arg Ile Leu Ser Leu Thr Val Leu Phe  
10 15 20  
atc gca ttt tgc ggt ttc tcc ttc tac ctc ggt ggt ata ttt tgc tct 328  
Ile Ala Phe Cys Gly Phe Ser Phe Tyr Leu Gly Gly Ile Phe Cys Ser  
25 30 35 40  
gag aga gac aag att gta gcc aag gat gtc aca agg acg act aca aag 376  
Glu Arg Asp Lys Ile Val Ala Lys Asp Val Thr Arg Thr Thr Thr Lys  
45 50 55  
gct gta gct tcc cct aaa gaa cct aca gct act cct att caa atc aaa 424  
Ala Val Ala Ser Pro Lys Glu Pro Thr Ala Thr Pro Ile Gln Ile Lys  
60 65 70  
tcc gtt tct ttc ccg gag tgc ggg tca gag ttc caa gat tac acc ccg 472  
Ser Val Ser Phe Pro Glu Cys Gly Ser Glu Phe Gln Asp Tyr Thr Pro  
75 80 85  
tgc acc gat cca aag agg tgg aag aag tat ggt gtc cat cgc tta agt 520  
Cys Thr Asp Pro Lys Arg Trp Lys Lys Tyr Gly Val His Arg Leu Ser  
90 95 100  
ttc ttg gag cgt cat tgt cct ccg gta tat gaa aag aat gag tgt ttg 568  
Phe Leu Glu Arg His Cys Pro Pro Val Tyr Glu Lys Asn Glu Cys Leu  
105 110 115 120  
att cca cca cca gac ggg tat aaa ccg cct ata aga tgg ccc aag agc 616  
Ile Pro Pro Pro Asp Gly Tyr Lys Pro Pro Ile Arg Trp Pro Lys Ser  
125 130 135  
cga gaa cag tgt tgg tac agg aac gtg cct tat gat tgg atc aat aag 664  
Arg Glu Gln Cys Trp Tyr Arg Asn Val Pro Tyr Asp Trp Ile Asn Lys

## MBI-20 Sequence Listing.ST25

140	145	150	
caa aag tct aac cag cat tgg ctt aag aaa gaa gga gat aag ttc cat Gln Lys Ser Asn Gln His Trp Leu Lys Lys Glu Gly Asp Lys Phe His 155 160 165			712
ttc cct ggt ggt ggt acc atg ttc cct cgt gga gtt agt cac tat gtt Phe Pro Gly Gly Gly Thr Met Phe Pro Arg Gly Val Ser His Tyr Val 170 175 180			760
gat ttg atg caa gat ctg att cct gaa atg aaa gac gga aca gtc agg Asp Leu Met Gln Asp Leu Ile Pro Glu Met Lys Asp Gly Thr Val Arg 185 190 195 200			808
acc gcc att gat act ggc tgt ggg gtt gcg agc tgg gga ggc gat ctt Thr Ala Ile Asp Thr Gly Cys Gly Val Ala Ser Trp Gly Gly Asp Leu 205 210 215			856
ttg gac cgt ggg ata cta tca ctc tct ctt gct cca aga gat aac cat Leu Asp Arg Gly Ile Leu Ser Leu Ser Leu Ala Pro Arg Asp Asn His 220 225 230			904
gaa gct cag gtt caa ttt gct ctt gaa cgt gga att cct gcg att ctc Glu Ala Gln Val Gln Phe Ala Leu Glu Arg Gly Ile Pro Ala Ile Leu 235 240 245			952
ggg atc atc tct acg caa cgt ctc cct ttt cct tca aat gca ttt gat Gly Ile Ile Ser Thr Gln Arg Leu Pro Phe Pro Ser Asn Ala Phe Asp 250 255 260			1000
atg gct cat tgt tca aga tgt ctt att ccc tgg aca gaa ttt ggt gga Met Ala His Cys Ser Arg Cys Leu Ile Pro Trp Thr Glu Phe Gly Gly 265 270 275 280			1048
atc tat tta ctt gag att cac cgt ata gtt cga cct gga ggt ttt tgg Ile Tyr Leu Leu Glu Ile His Arg Ile Val Arg Pro Gly Gly Phe Trp 285 290 295			1096
gtt ctt tct ggt cca cct gtg aac tat aat aga cga tgg cgt gga tgg Val Leu Ser Gly Pro Pro Val Asn Tyr Asn Arg Arg Trp Arg Gly Trp 300 305 310			1144
aac aca acc atg gaa gat cag aaa tct gac tac aac aag ctt cag tca Asn Thr Thr Met Glu Asp Gln Lys Ser Asp Tyr Asn Lys Leu Gln Ser 315 320 325			1192
ctt cta acc tcc atg tgt ttc aaa aag tac gct caa aaa gat gac ata Leu Leu Thr Ser Met Cys Phe Lys Lys Tyr Ala Gln Lys Asp Asp Ile 330 335 340			1240
gcc gtg tgg cag aaa ctc tca gac aaa tct tgc tat gac aaa atc gct Ala Val Trp Gln Lys Leu Ser Asp Lys Ser Cys Tyr Asp Lys Ile Ala 345 350 355 360			1288
aag aac atg gaa gct tac cct ccc aaa tgt gac gac agt ata gaa cct Lys Asn Met Glu Ala Tyr Pro Pro Lys Cys Asp Asp Ser Ile Glu Pro 365 370 375			1336
gat tct gct tgg tac act cca ctc cgt cct tgc gtg gtt gcc ccg aca Asp Ser Ala Trp Tyr Thr Pro Leu Arg Pro Cys Val Val Ala Pro Thr 380 385 390			1384
cct aaa gtc aag aag tct ggt ctc gga tca atc cca aaa tgg ccc gag Pro Lys Val Lys Lys Ser Gly Leu Gly Ser Ile Pro Lys Trp Pro Glu 395 400 405			1432
agg tta cat gtc gcg ccc gag aga atc ggt gat gtt cac gga ggg agt Arg Leu His Val Ala Pro Glu Arg Ile Gly Asp Val His Gly Gly Ser 410 415 420			1480
gcg aac agt ttg aaa cac gat gat ggt aaa tgg aag aac aga gtt aag Ala Asn Ser Leu Lys His Asp Asp Gly Lys Trp Lys Asn Arg Val Lys 425 430 435 440			1528
cat tac aag aaa gtt tta cca gct ctt ggg aca gac aag ata aga aat			1576



## MBI-20 Sequence Listing.ST25

His Tyr Lys Lys Val Leu Pro Ala Leu Gly Thr Asp Lys Ile Arg Asn  
 445 450 455  
 gtt atg gat atg aac act gtt tat gga ggt ttc tct gcg gcc ctc att 1624  
 Val Met Asp Met Asn Thr Val Tyr Gly Gly Phe Ser Ala Ala Leu Ile  
 460 465 470  
 gag gat ccc att tgg gtc atg aac gtt gta tca tcg tac agc gca aat 1672  
 Glu Asp Pro Ile Trp Val Met Asn Val Val Ser Ser Tyr Ser Ala Asn  
 475 480 485  
 tcg ctt cct gtt gtc ttt gat cgc ggt ctc atc ggg act tac cac gac 1720  
 Ser Leu Pro Val Val Phe Asp Arg Gly Leu Ile Gly Thr Tyr His Asp  
 490 495 500  
 tgg tgc gaa gct ttc tca acg tat cca aga aca tat gat ctt ctt cac 1768  
 Trp Cys Glu Ala Phe Ser Thr Tyr Pro Arg Thr Tyr Asp Leu Leu His  
 505 510 515 520  
 ctc gac agt ctt ttt acc ttg gag agt cac agg tgt gag atg aag tac 1816  
 Leu Asp Ser Leu Phe Thr Leu Glu Ser His Arg Cys Glu Met Lys Tyr  
 525 530 535  
 att ttg cta gag atg gac agg atc ttg cgg ccg agt gga tat gtt ata 1864  
 Ile Leu Leu Glu Met Asp Arg Ile Leu Arg Pro Ser Gly Tyr Val Ile  
 540 545 550  
 atc cga gaa tcg agt tat ttc atg gac gca atc aca acg tta gcg aaa 1912  
 Ile Arg Glu Ser Ser Tyr Phe Met Asp Ala Ile Thr Thr Leu Ala Lys  
 555 560 565  
 ggg ata agg tgg agt tgc cgg aga gag gag act gag tat gca gtc aaa 1960  
 Gly Ile Arg Trp Ser Cys Arg Arg Glu Glu Thr Glu Tyr Ala Val Lys  
 570 575 580  
 agt gag aag att ctg gtt tgc cag aaa aag cta tgg ttt tcg tca aac 2008  
 Ser Glu Lys Ile Leu Val Cys Gln Lys Lys Leu Trp Phe Ser Ser Asn  
 585 590 595 600  
 caa acc tct tga tgagaccacc tgtatcatag tgtttatcat ctctctgtgat 2060  
 Gln Thr Ser  
 gcacactaca gagagaagga tctagtcttt tgagtcgaag atatagctct ataaacaatc 2120  
 tcctttttttt gttctcttta atttcttggg tatttcacgg tatagattga tattatatat 2180  
 tttttaatta tatttttaat atatagatat attagtatgt gggttaaaca ctattattat 2240  
 caaggtctta aagatttgct ttgcaagagt taaaaaatgt tggagtaagg acctcttgat 2300  
 taataaattg actgacgcag caaa 2324  
 <210> 36  
 <211> 603  
 <212> PRT  
 <213> Arabidopsis thaliana  
 <400> 36  
 Met Lys Ser Gly Lys Gln Ser Ser Gln Pro Glu Lys Gly Thr Ser Arg  
 1 5 10 15  
 Ile Leu Ser Leu Thr Val Leu Phe Ile Ala Phe Cys Gly Phe Ser Phe  
 20 25 30  
 Tyr Leu Gly Gly Ile Phe Cys Ser Glu Arg Asp Lys Ile Val Ala Lys  
 35 40 45  
 Asp Val Thr Arg Thr Thr Thr Lys Ala Val Ala Ser Pro Lys Glu Pro  
 50 55 60

## MBI-20 Sequence Listing.ST25

Thr Ala Thr Pro Ile Gln Ile Lys Ser Val Ser Phe Pro Glu Cys Gly  
 65 70 75 80  
 Ser Glu Phe Gln Asp Tyr Thr Pro Cys Thr Asp Pro Lys Arg Trp Lys  
 85 90 95  
 Lys Tyr Gly Val His Arg Leu Ser Phe Leu Glu Arg His Cys Pro Pro  
 100 105 110  
 Val Tyr Glu Lys Asn Glu Cys Leu Ile Pro Pro Pro Asp Gly Tyr Lys  
 115 120 125  
 Pro Pro Ile Arg Trp Pro Lys Ser Arg Glu Gln Cys Trp Tyr Arg Asn  
 130 135 140  
 Val Pro Tyr Asp Trp Ile Asn Lys Gln Lys Ser Asn Gln His Trp Leu  
 145 150 155 160  
 Lys Lys Glu Gly Asp Lys Phe His Phe Pro Gly Gly Gly Thr Met Phe  
 165 170 175  
 Pro Arg Gly Val Ser His Tyr Val Asp Leu Met Gln Asp Leu Ile Pro  
 180 185 190  
 Glu Met Lys Asp Gly Thr Val Arg Thr Ala Ile Asp Thr Gly Cys Gly  
 195 200 205  
 Val Ala Ser Trp Gly Gly Asp Leu Leu Asp Arg Gly Ile Leu Ser Leu  
 210 215 220  
 Ser Leu Ala Pro Arg Asp Asn His Glu Ala Gln Val Gln Phe Ala Leu  
 225 230 235 240  
 Glu Arg Gly Ile Pro Ala Ile Leu Gly Ile Ile Ser Thr Gln Arg Leu  
 245 250 255  
 Pro Phe Pro Ser Asn Ala Phe Asp Met Ala His Cys Ser Arg Cys Leu  
 260 265 270  
 Ile Pro Trp Thr Glu Phe Gly Gly Ile Tyr Leu Leu Glu Ile His Arg  
 275 280 285  
 Ile Val Arg Pro Gly Gly Phe Trp Val Leu Ser Gly Pro Pro Val Asn  
 290 295 300  
 Tyr Asn Arg Arg Trp Arg Gly Trp Asn Thr Thr Met Glu Asp Gln Lys  
 305 310 315 320  
 Ser Asp Tyr Asn Lys Leu Gln Ser Leu Leu Thr Ser Met Cys Phe Lys  
 325 330 335  
 Lys Tyr Ala Gln Lys Asp Asp Ile Ala Val Trp Gln Lys Leu Ser Asp  
 340 345 350  
 Lys Ser Cys Tyr Asp Lys Ile Ala Lys Asn Met Glu Ala Tyr Pro Pro

MBI-20 Sequence Listing.ST25  
360 365

355

Lys Cys Asp Asp Ser Ile Glu Pro Asp Ser Ala Trp Tyr Thr Pro Leu  
370 375 380

Arg Pro Cys Val Val Ala Pro Thr Pro Lys Val Lys Lys Ser Gly Leu  
385 390 395 400

Gly Ser Ile Pro Lys Trp Pro Glu Arg Leu His Val Ala Pro Glu Arg  
405 410 415

Ile Gly Asp Val His Gly Gly Ser Ala Asn Ser Leu Lys His Asp Asp  
420 425 430

Gly Lys Trp Lys Asn Arg Val Lys His Tyr Lys Lys Val Leu Pro Ala  
435 440 445

Leu Gly Thr Asp Lys Ile Arg Asn Val Met Asp Met Asn Thr Val Tyr  
450 455 460

Gly Gly Phe Ser Ala Ala Leu Ile Glu Asp Pro Ile Trp Val Met Asn  
465 470 475 480

Val Val Ser Ser Tyr Ser Ala Asn Ser Leu Pro Val Val Phe Asp Arg  
485 490 495

Gly Leu Ile Gly Thr Tyr His Asp Trp Cys Glu Ala Phe Ser Thr Tyr  
500 505 510

Pro Arg Thr Tyr Asp Leu Leu His Leu Asp Ser Leu Phe Thr Leu Glu  
515 520 525

Ser His Arg Cys Glu Met Lys Tyr Ile Leu Leu Glu Met Asp Arg Ile  
530 535 540

Leu Arg Pro Ser Gly Tyr Val Ile Ile Arg Glu Ser Ser Tyr Phe Met  
545 550 555 560

Asp Ala Ile Thr Thr Leu Ala Lys Gly Ile Arg Trp Ser Cys Arg Arg  
565 570 575

Glu Glu Thr Glu Tyr Ala Val Lys Ser Glu Lys Ile Leu Val Cys Gln  
580 585 590

Lys Lys Leu Trp Phe Ser Ser Asn Gln Thr Ser  
595 600

<210> 37  
<211> 1951  
<212> DNA  
<213> Arabidopsis thaliana

<220>  
<221> CDS  
<222> (196)..(1794)  
<223> G308

<400> 37

## MBI-20 Sequence Listing.ST25

```

agtaatttag tttttttttt ttttttttac aattttttt gttattagaa gtggtagtgg      60
. agtgaaaaaa caaatcctaa gcagtcctaa ccgatccccc aagctaaaga ttcttcacct      120
tcccaaataa agcaaaaacct agatccgaca ttgaaggaaa aaccttttag atccatctct      180
gaaaaaaacc caacc atg aag aga gat cat cat cat cat cat caa gat aag      231
                Met Lys Arg Asp His His His His His Gln Asp Lys
                1          5          10

aag act atg atg atg aat gaa gaa gac gac ggt aac ggc atg gat gag      279
Lys Thr Met Met Met Asn Glu Glu Asp Asp Gly Asn Gly Met Asp Glu
                15          20          25

ctt cta gct gtt ctt ggt tac aag gtt agg tca tcg gaa atg gct gat      327
Leu Leu Ala Val Leu Gly Tyr Lys Val Arg Ser Ser Glu Met Ala Asp
                30          35          40

ggt gct cag aaa ctc gag cag ctt gaa gtt atg atg tct aat gtt caa      375
Val Ala Gln Lys Leu Glu Gln Leu Glu Val Met Met Ser Asn Val Gln
                45          50          55          60

gaa gac gat ctt tct caa ctc gct act gag act gtt cac tat aat ccg      423
Glu Asp Asp Leu Ser Gln Leu Ala Thr Glu Thr Val His Tyr Asn Pro
                65          70          75

gcg gag ctt tac acg tgg ctt gat tct atg ctc acc gac ctt aat cct      471
Ala Glu Leu Tyr Thr Trp Leu Asp Ser Met Leu Thr Asp Leu Asn Pro
                80          85          90

ccg tcg tct aac gcc gag tac gat ctt aaa gct att ccc ggt gac gcg      519
Pro Ser Ser Asn Ala Glu Tyr Asp Leu Lys Ala Ile Pro Gly Asp Ala
                95          100          105

att ctc aat cag ttc gct atc gat tcg gct tct tcg tct aac caa ggc      567
Ile Leu Asn Gln Phe Ala Ile Asp Ser Ala Ser Ser Ser Asn Gln Gly
                110          115          120

ggc gga gga gat acg tat act aca aac aag cgg ttg aaa tgc tca aac      615
Gly Gly Gly Asp Thr Tyr Thr Thr Asn Lys Arg Leu Lys Cys Ser Asn
                125          130          135          140

ggc gtc gtg gaa acc acc aca gcg acg gct gag tca act cgg cat gtt      663
Gly Val Val Glu Thr Thr Thr Ala Thr Ala Glu Ser Thr Arg His Val
                145          150          155

gtc ctg gtt gac tcg cag gag aac ggt gtg cgt ctc gtt cac gcg ctt      711
Val Leu Val Asp Ser Gln Glu Asn Gly Val Arg Leu Val His Ala Leu
                160          165          170

ttg gct tgc gct gaa gct gtt cag aag gag aat ctg act gtg gcg gaa      759
Leu Ala Cys Ala Glu Ala Val Gln Lys Glu Asn Leu Thr Val Ala Glu
                175          180          185

gct ctg gtg aag caa atc gga ttc tta gct gtt tct caa atc gga gct      807
Ala Leu Val Lys Gln Ile Gly Phe Leu Ala Val Ser Gln Ile Gly Ala
                190          195          200

atg aga caa gtc gct act tac ttc gcc gaa gct ctc gcg cgg cgg att      855
Met Arg Gln Val Ala Thr Tyr Phe Ala Glu Ala Leu Ala Arg Arg Ile
                205          210          215          220

tac cgt ctc tct ccg tcg cag agt cca atc gac cac tct ctc tcc gat      903
Tyr Arg Leu Ser Pro Ser Gln Ser Pro Ile Asp His Ser Leu Ser Asp
                225          230          235

act ctt cag atg cac ttc tac gag act tgt cct tat ctc aag ttc gct      951
Thr Leu Gln Met His Phe Tyr Glu Thr Cys Pro Tyr Leu Lys Phe Ala
                240          245          250

cac ttc acg gcg aat caa gcg att ctc gaa gct ttt caa ggg aag aaa      999
His Phe Thr Ala Asn Gln Ala Ile Leu Glu Ala Phe Gln Gly Lys Lys
                255          260          265

aga gtt cat gtc att gat ttc tct atg agt caa ggt ctt caa tgg ccg      1047

```

MBI-20 Sequence Listing.ST25															
Arg	Val	His	Val	Ile	Asp	Phe	Ser	Met	Ser	Gln	Gly	Leu	Gln	Trp	Pro
270						275					280				
gcg	ctt	atg	cag	gct	ctt	gcg	ctt	cga	cct	ggt	ggt	cct	cct	gtt	ttc
Ala	Leu	Met	Gln	Ala	Leu	Ala	Leu	Arg	Pro	Gly	Gly	Pro	Pro	Val	Phe
285				290					295						300
cgg	tta	acc	gga	att	ggt	cca	ccg	gca	ccg	gat	aat	ttc	gat	tat	ctt
Arg	Leu	Thr	Gly	Ile	Gly	Pro	Pro	Ala	Pro	Asp	Asn	Phe	Asp	Tyr	Leu
				305					310					315	
cat	gaa	gtt	ggg	tgt	aag	ctg	gct	cat	tta	gct	gag	gcg	att	cac	gtt
His	Glu	Val	Gly	Cys	Lys	Leu	Ala	His	Leu	Ala	Glu	Ala	Ile	His	Val
			320					325					330		
gag	ttt	gag	tac	aga	gga	ttt	gtg	gct	aac	act	tta	gct	gat	ctt	gat
Glu	Phe	Glu	Tyr	Arg	Gly	Phe	Val	Ala	Asn	Thr	Leu	Ala	Asp	Leu	Asp
		335					340					345			
gct	tcg	atg	ctt	gag	ctt	aga	cca	agt	gag	att	gaa	tct	gtt	gcg	gtt
Ala	Ser	Met	Leu	Glu	Leu	Arg	Pro	Ser	Glu	Ile	Glu	Ser	Val	Ala	Val
		350				355					360				
aac	tct	gtt	ttc	gag	ctt	cac	aag	ctc	ttg	gga	cga	cct	ggt	gcg	atc
Asn	Ser	Val	Phe	Glu	Leu	His	Lys	Leu	Leu	Gly	Arg	Pro	Gly	Ala	Ile
365				370					375						380
gat	aag	gtt	ctt	ggt	gtg	gtg	aat	cag	att	aaa	ccg	gag	att	ttc	act
Asp	Lys	Val	Leu	Gly	Val	Val	Asn	Gln	Ile	Lys	Pro	Glu	Ile	Phe	Thr
			385					390						395	
gtg	gtt	gag	cag	gaa	tcg	aac	cat	aat	agt	ccg	att	ttc	tta	gat	cgg
Val	Val	Glu	Gln	Glu	Ser	Asn	His	Asn	Ser	Pro	Ile	Phe	Leu	Asp	Arg
			400					405					410		
ttt	act	gag	tcg	ttg	cat	tat	tac	tcg	acg	ttg	ttt	gac	tcg	ttg	gaa
Phe	Thr	Glu	Ser	Leu	His	Tyr	Tyr	Ser	Thr	Leu	Phe	Asp	Ser	Leu	Glu
		415				420						425			
ggt	gta	ccg	agt	ggt	caa	gac	aag	gtc	atg	tcg	gag	gtt	tac	ttg	ggt
Gly	Val	Pro	Ser	Gly	Gln	Asp	Lys	Val	Met	Ser	Glu	Val	Tyr	Leu	Gly
	430					435					440				
aaa	cag	atc	tgc	aac	gtt	gtg	gct	tgt	gat	gga	cct	gac	cga	gtt	gag
Lys	Gln	Ile	Cys	Asn	Val	Val	Ala	Cys	Asp	Gly	Pro	Asp	Arg	Val	Glu
445				450						455					460
cgt	cat	gaa	acg	ttg	agt	cag	tgg	agg	aac	cgg	ttc	ggg	tct	gct	ggg
Arg	His	Glu	Thr	Leu	Ser	Gln	Trp	Arg	Asn	Arg	Phe	Gly	Ser	Ala	Gly
				465					470					475	
ttt	gcg	gct	gca	cat	att	ggt	tcg	aat	gcg	ttt	aag	caa	gcg	agt	atg
Phe	Ala	Ala	Ala	His	Ile	Gly	Ser	Asn	Ala	Phe	Lys	Gln	Ala	Ser	Met
			480					485					490		
ctt	ttg	gct	ctg	ttc	aac	ggc	ggt	gag	ggt	tat	cgg	gtg	gag	gag	agt
Leu	Leu	Ala	Leu	Phe	Asn	Gly	Gly	Glu	Gly	Tyr	Arg	Val	Glu	Glu	Ser
		495				500						505			
gac	ggc	tgt	ctc	atg	ttg	ggt	tgg	cac	aca	cga	ccg	ctc	ata	gcc	acc
Asp	Gly	Cys	Leu	Met	Leu	Gly	Trp	His	Thr	Arg	Pro	Leu	Ile	Ala	Thr
	510					515					520				
tcg	gct	tgg	aaa	ctc	tcc	acc	aat	tag	atggtggctc	aatgaattga					
Ser	Ala	Trp	Lys	Leu	Ser	Thr	Asn								
525					530										
tctgttgaac	cggttatgat	gatagatttc	cgaccgaagc	caaaactaaat	cctactgttt										
1874															
ttccctttgt	cacttgtaa	gatcttatct	ttcattatat	taggtaattg	aaaaatttta										
1934															
atctcgcta	aattact														
1951															

## MBI-20 Sequence Listing.ST25

&lt;211&gt; 532

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 38

```

Met Lys Arg Asp His His His His His Gln Asp Lys Lys Thr Met Met
1      5      10      15

Met Asn Glu Glu Asp Asp Gly Asn Gly Met Asp Glu Leu Leu Ala Val
      20      25      30

Leu Gly Tyr Lys Val Arg Ser Ser Glu Met Ala Asp Val Ala Gln Lys
      35      40      45

Leu Glu Gln Leu Glu Val Met Met Ser Asn Val Gln Glu Asp Asp Leu
      50      55      60

Ser Gln Leu Ala Thr Glu Thr Val His Tyr Asn Pro Ala Glu Leu Tyr
      65      70      75      80

Thr Trp Leu Asp Ser Met Leu Thr Asp Leu Asn Pro Pro Ser Ser Asn
      85      90      95

Ala Glu Tyr Asp Leu Lys Ala Ile Pro Gly Asp Ala Ile Leu Asn Gln
      100      105      110

Phe Ala Ile Asp Ser Ala Ser Ser Ser Asn Gln Gly Gly Gly Gly Asp
      115      120      125

Thr Tyr Thr Thr Asn Lys Arg Leu Lys Cys Ser Asn Gly Val Val Glu
      130      135      140

Thr Thr Thr Ala Thr Ala Glu Ser Thr Arg His Val Val Leu Val Asp
      145      150      155      160

Ser Gln Glu Asn Gly Val Arg Leu Val His Ala Leu Leu Ala Cys Ala
      165      170      175

Glu Ala Val Gln Lys Glu Asn Leu Thr Val Ala Glu Ala Leu Val Lys
      180      185      190

Gln Ile Gly Phe Leu Ala Val Ser Gln Ile Gly Ala Met Arg Gln Val
      195      200      205

Ala Thr Tyr Phe Ala Glu Ala Leu Ala Arg Arg Ile Tyr Arg Leu Ser
      210      215      220

Pro Ser Gln Ser Pro Ile Asp His Ser Leu Ser Asp Thr Leu Gln Met
      225      230      235      240

His Phe Tyr Glu Thr Cys Pro Tyr Leu Lys Phe Ala His Phe Thr Ala
      245      250      255

Asn Gln Ala Ile Leu Glu Ala Phe Gln Gly Lys Lys Arg Val His Val
      260      265      270

Ile Asp Phe Ser Met Ser Gln Gly Leu Gln Trp Pro Ala Leu Met Gln

```

MBI-20 Sequence Listing.ST25  
280 285

275

Ala Leu Ala Leu Arg Pro Gly Gly Pro Pro Val Phe Arg Leu Thr Gly  
290 295 300

Ile Gly Pro Pro Ala Pro Asp Asn Phe Asp Tyr Leu His Glu Val Gly  
305 310 315 320

Cys Lys Leu Ala His Leu Ala Glu Ala Ile His Val Glu Phe Glu Tyr  
325 330 335

Arg Gly Phe Val Ala Asn Thr Leu Ala Asp Leu Asp Ala Ser Met Leu  
340 345 350

Glu Leu Arg Pro Ser Glu Ile Glu Ser Val Ala Val Asn Ser Val Phe  
355 360 365

Glu Leu His Lys Leu Leu Gly Arg Pro Gly Ala Ile Asp Lys Val Leu  
370 375 380

Gly Val Val Asn Gln Ile Lys Pro Glu Ile Phe Thr Val Val Glu Gln  
385 390 395 400

Glu Ser Asn His Asn Ser Pro Ile Phe Leu Asp Arg Phe Thr Glu Ser  
405 410 415

Leu His Tyr Tyr Ser Thr Leu Phe Asp Ser Leu Glu Gly Val Pro Ser  
420 425 430

Gly Gln Asp Lys Val Met Ser Glu Val Tyr Leu Gly Lys Gln Ile Cys  
435 440 445

Asn Val Val Ala Cys Asp Gly Pro Asp Arg Val Glu Arg His Glu Thr  
450 455 460

Leu Ser Gln Trp Arg Asn Arg Phe Gly Ser Ala Gly Phe Ala Ala Ala  
465 470 475 480

His Ile Gly Ser Asn Ala Phe Lys Gln Ala Ser Met Leu Leu Ala Leu  
485 490 495

Phe Asn Gly Gly Glu Gly Tyr Arg Val Glu Glu Ser Asp Gly Cys Leu  
500 505 510

Met Leu Gly Trp His Thr Arg Pro Leu Ile Ala Thr Ser Ala Trp Lys  
515 520 525

Leu Ser Thr Asn  
530

<210> 39  
<211> 1445  
<212> DNA  
<213> Arabidopsis thaliana

<220>  
<221> CDS  
<222> (236)..(1306)

## MBI-20 Sequence Listing.ST25

&lt;223&gt; G1944

&lt;400&gt; 39

tcgaccttcc taatttccaa cctctgttct tagcaatata ttttttctcc aaaaataatt	60
ctcagtttga ttttcttctt cttagctctta agtatatttc tttgttgta tttatctttt	120
aatcctttaa tctcatcttt gtttatcttt aatcaaaacc caaaatttac atgggttctt	180
gaaaatctag aagaaataaa ggaaacataa caaaaataga aagaaaaaga agcta atg Met 1	238
gtc tta aat atg gag tct acc gga gaa gct gtt aga tca acc acc ggt Val Leu Asn Met Glu Ser Thr Gly Glu Ala Val Arg Ser Thr Thr Gly 5 10 15	286
aac gac ggt ggt att acg gtg gtt aga tcc gac gcg ccg tca gat ttc Asn Asp Gly Gly Ile Thr Val Val Arg Ser Asp Ala Pro Ser Asp Phe 20 25 30	334
cac gta gct caa aga tca gaa agc tca aac caa tct ccc acc tct gtc His Val Ala Gln Arg Ser Glu Ser Ser Asn Gln Ser Pro Thr Ser Val 35 40 45	382
act cct cct cca cca cag cca tcg tct cat cac aca gct cct ccg ccg Thr Pro Pro Pro Pro Glu Pro Ser Ser His His Thr Ala Pro Pro Pro 50 55 60 65	430
ctg caa att tcg acg gtg acg act acg act acg acg gcc gcg atg gaa Leu Gln Ile Ser Thr Val Thr Thr Thr Thr Thr Ala Ala Met Glu 70 75 80	478
ggt atc tcc ggt gga ctg atg aag aag aag cgt gga cgg cca agg aag Gly Ile Ser Gly Gly Leu Met Lys Lys Lys Arg Gly Arg Pro Arg Lys 85 90 95	526
tat gga ccg gac ggg act gtt gta gcg tta tct cct aaa ccg att tca Tyr Gly Pro Asp Gly Thr Val Val Ala Leu Ser Pro Lys Pro Ile Ser 100 105 110	574
tca gcg ccg gcg ccg tcg cat ctt ccg ccg ccg agt tca cac gtc atc Ser Ala Pro Ala Pro Ser His Leu Pro Pro Pro Ser Ser His Val Ile 115 120 125	622
gat ttc tcc gct tct gag aaa cgt agc aaa gtg aaa cca acg aac tcg Asp Phe Ser Ala Ser Glu Lys Arg Ser Lys Val Lys Pro Thr Asn Ser 130 135 140 145	670
ttt aac aga aca aag tat cat cac caa gtt gag aat ttg ggt gaa tgg Phe Asn Arg Thr Lys Tyr His His Gln Val Glu Asn Leu Gly Glu Trp 150 155 160	718
gct cct tgc tcc gtc ggt ggt aat ttc aca cct cat ata atc aca gtc Ala Pro Cys Ser Val Gly Gly Asn Phe Thr Pro His Ile Ile Thr Val 165 170 175	766
aac acc ggc gag gat gta aca atg aag ata atc tcg ttt tcg caa caa Asn Thr Gly Glu Asp Val Thr Met Lys Ile Ile Ser Phe Ser Gln Gln 180 185 190	814
gga cct cgc tct att tgt gtt ctg tca gca aac ggt gtt att tca agc Gly Pro Arg Ser Ile Cys Val Leu Ser Ala Asn Gly Val Ile Ser Ser 195 200 205	862
gtt aca ctt cgt cag cca gat tcc tct ggc ggc aca ttg aca tac gaa Val Thr Leu Arg Gln Pro Asp Ser Ser Gly Gly Thr Leu Thr Tyr Glu 210 215 220 225	910
ggt cgg ttt gag ata tta tca tta tcc ggg tca ttc atg cct aat gat Gly Arg Phe Glu Ile Leu Ser Leu Ser Gly Ser Phe Met Pro Asn Asp 230 235 240	958
tca ggc gga aca cga agt aga acg gga gga atg agt gta tcg tta gca	1006



## MBI-20 Sequence Listing.ST25

Ser Gly Gly Thr Arg Ser Arg Thr Gly Gly Met Ser Val Ser Leu Ala  
 245 250 255

agt ccc gat gga cgt gta gta ggc ggt ggc ctc gcc ggt tta cta gta 1054  
 Ser Pro Asp Gly Arg Val Val Gly Gly Gly Leu Ala Gly Leu Leu Val  
 260 265 270

gcc gcg agt ccg gtt cag gtg gtt gta gga agt ttt tta gcg gcc act 1102  
 Ala Ala Ser Pro Val Gln Val Val Gly Ser Phe Leu Ala Gly Thr  
 275 280 285

gac cat caa gat cag aaa ccg aaa aag aac aaa cat gat ttc atg ttg 1150  
 Asp His Gln Asp Gln Lys Pro Lys Lys Asn Lys His Asp Phe Met Leu  
 290 295 300 305

tcg agt cct acc gct gca att cct atc tct agt gca gct gat cac cgg 1198  
 Ser Ser Pro Thr Ala Ala Ile Pro Ile Ser Ser Ala Ala Asp His Arg  
 310 315 320

aca atc cat tcg gtc tcg tct ctt ccg gtc aat aat aat aca tgg cag 1246  
 Thr Ile His Ser Val Ser Ser Leu Pro Val Asn Asn Asn Thr Trp Gln  
 325 330 335

act tct tta gct tcc gat cca aga aac aag cat acc gat att aat gtc 1294  
 Thr Ser Leu Ala Ser Asp Pro Arg Asn Lys His Thr Asp Ile Asn Val  
 340 345 350

aat gta act tga aatccaatct ttctctgtat ttctctgttaa caagtttgat 1346  
 Asn Val Thr  
 355

ttgggtgttt atctacatta ggattttact aaaatggtag tattatttat aggggttttag 1406

ggctctttatt ttggttccac tggtgtcact tgtaggata 1445

<210> 40  
 <211> 356  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 40

Met Val Leu Asn Met Glu Ser Thr Gly Glu Ala Val Arg Ser Thr Thr  
 1 5 10 15

Gly Asn Asp Gly Gly Ile Thr Val Val Arg Ser Asp Ala Pro Ser Asp  
 20 25 30

Phe His Val Ala Gln Arg Ser Glu Ser Ser Asn Gln Ser Pro Thr Ser  
 35 40 45

Val Thr Pro Pro Pro Pro Gln Pro Ser Ser His His Thr Ala Pro Pro  
 50 55 60

Pro Leu Gln Ile Ser Thr Val Thr Thr Thr Thr Thr Thr Ala Ala Met  
 65 70 75 80

Glu Gly Ile Ser Gly Gly Leu Met Lys Lys Lys Arg Gly Arg Pro Arg  
 85 90 95

Lys Tyr Gly Pro Asp Gly Thr Val Val Ala Leu Ser Pro Lys Pro Ile  
 100 105 110

Ser Ser Ala Pro Ala Pro Ser His Leu Pro Pro Pro Ser Ser His Val  
 115 120 125

## MBI-20 Sequence Listing.ST25

Ile Asp Phe Ser Ala Ser Glu Lys Arg Ser Lys Val Lys Pro Thr Asn  
 130 135 140

Ser Phe Asn Arg Thr Lys Tyr His His Gln Val Glu Asn Leu Gly Glu  
 145 150 155 160

Trp Ala Pro Cys Ser Val Gly Gly Asn Phe Thr Pro His Ile Ile Thr  
 165 170 175

Val Asn Thr Gly Glu Asp Val Thr Met Lys Ile Ile Ser Phe Ser Gln  
 180 185 190

Gln Gly Pro Arg Ser Ile Cys Val Leu Ser Ala Asn Gly Val Ile Ser  
 195 200 205

Ser Val Thr Leu Arg Gln Pro Asp Ser Ser Gly Gly Thr Leu Thr Tyr  
 210 215 220

Glu Gly Arg Phe Glu Ile Leu Ser Leu Ser Gly Ser Phe Met Pro Asn  
 225 230 235 240

Asp Ser Gly Gly Thr Arg Ser Arg Thr Gly Gly Met Ser Val Ser Leu  
 245 250 255

Ala Ser Pro Asp Gly Arg Val Val Gly Gly Gly Leu Ala Gly Leu Leu  
 260 265 270

Val Ala Ala Ser Pro Val Gln Val Val Val Gly Ser Phe Leu Ala Gly  
 275 280 285

Thr Asp His Gln Asp Gln Lys Pro Lys Lys Asn Lys His Asp Phe Met  
 290 295 300

Leu Ser Ser Pro Thr Ala Ala Ile Pro Ile Ser Ser Ala Ala Asp His  
 305 310 315 320

Arg Thr Ile His Ser Val Ser Ser Leu Pro Val Asn Asn Asn Thr Trp  
 325 330 335

Gln Thr Ser Leu Ala Ser Asp Pro Arg Asn Lys His Thr Asp Ile Asn  
 340 345 350

Val Asn Val Thr  
 355

<210> 41  
 <211> 1558  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (191)..(1396)  
 <223> G326

<400> 41  
 caattaatga catcttcttc ttctcctttc actgcaaaac cgaaagcttg agactttgag 60  
 attatgtcta tgcacatctc ttcttcttcc atcgatcact tcacacaccc tcgtcatctt 120

## MBI-20 Sequence Listing.ST25

gatcttatttc tccactgcat aaaatcagcg agattttaag ggattgtgaa ggtaccatct	180
taaacacaaaa atg ggt act tct act aca gag agt gtg gtg gcg tgt gaa	229
Met Gly Thr Ser Thr Thr Glu Ser Val Val Ala Cys Glu	
1 5 10	
ttt tgc ggc gag aga acg gcg gtt ctg ttt tgt aga gcc gat acg gcg	277
Phe Cys Gly Glu Arg Thr Ala Val Leu Phe Cys Arg Ala Asp Thr Ala	
15 20 25	
aag ctt tgt ttg cct tgt gac cag cac gtg cac tcg gcg aac ctt ctc	325
Lys Leu Cys Leu Pro Cys Asp Gln His Val His Ser Ala Asn Leu Leu	
30 35 40 45	
tcg agg aag cat gtt cgt tct cag atc tgt gat aac tgt agc aaa gag	373
Ser Arg Lys His Val Arg Ser Gln Ile Cys Asp Asn Cys Ser Lys Glu	
50 55 60	
ccg gtg tcc gta cgt tgc ttc aca gat aat ctc gta ttg tgt cag gag	421
Pro Val Ser Val Arg Cys Phe Thr Asp Asn Leu Val Leu Cys Gln Glu	
65 70 75	
tgt gat tgg gat gtt cac gga agc tgt tcc tcc tcc gcg acg cat gaa	469
Cys Asp Trp Asp Val His Gly Ser Cys Ser Ser Ser Ala Thr His Glu	
80 85 90	
cgc tcc gcc gtg gaa ggg ttt tca ggt tgt cct tcg gtt ttg gag ctt	517
Arg Ser Ala Val Glu Gly Phe Ser Gly Cys Pro Ser Val Leu Glu Leu	
95 100 105	
gct gct gtg tgg gga atc gat tta aag ggt aag aag aaa gaa gat gac	565
Ala Ala Val Trp Gly Ile Asp Leu Lys Gly Lys Lys Lys Glu Asp Asp	
110 115 120 125	
gaa gac gaa ttg act aag aat ttt ggg atg ggg ttg gat tcg tgg ggt	613
Glu Asp Glu Leu Thr Lys Asn Phe Gly Met Gly Leu Asp Ser Trp Gly	
130 135 140	
tct gga tct aac atc gtt caa gaa ctg att gtt cct tat gat gtg tct	661
Ser Gly Ser Asn Ile Val Gln Glu Leu Ile Val Pro Tyr Asp Val Ser	
145 150 155	
tgc aaa aag caa agc ttt agc ttt ggg agg tct aag cag gta gtg ttt	709
Cys Lys Lys Gln Ser Phe Ser Phe Gly Arg Ser Lys Gln Val Val Phe	
160 165 170	
gaa cag ctt gag tta ctg aag aga ggc ttc gtt gaa ggc gaa gga gag	757
Glu Gln Leu Glu Leu Leu Lys Arg Gly Phe Val Glu Gly Glu Gly Glu	
175 180 185	
att atg gtt ccg gag gga atc aat ggc gga gga agc att tct cag cca	805
Ile Met Val Pro Glu Gly Ile Asn Gly Gly Gly Ser Ile Ser Gln Pro	
190 195 200 205	
tct ccg acg acg tcg ttt act tct ttg ctt atg tct caa agt ctt tgt	853
Ser Pro Thr Thr Ser Phe Thr Ser Leu Leu Met Ser Gln Ser Leu Cys	
210 215 220	
ggg aat ggt atg caa tgg aat gct act aat cat agc act ggc cag aac	901
Gly Asn Gly Met Gln Trp Asn Ala Thr Asn His Ser Thr Gly Gln Asn	
225 230 235	
act cag ata tgg gat ttt aac ttg gga cag tcg agg aac cct gat gaa	949
Thr Gln Ile Trp Asp Phe Asn Leu Gly Gln Ser Arg Asn Pro Asp Glu	
240 245 250	
cct agt cca gtc gaa act aaa ggc tct act ttc aca ttc aac aac gtt	997
Pro Ser Pro Val Glu Thr Lys Gly Ser Thr Phe Thr Phe Asn Asn Val	
255 260 265	
act cat ctc aag aac gat acc cga acc acc aat atg aat gct ttc aaa	1045
Thr His Leu Lys Asn Asp Thr Arg Thr Thr Asn Met Asn Ala Phe Lys	
270 275 280 285	

## MBI-20 Sequence Listing.ST25

gag agt tac cag gag gat tcc gtc cac tca act tct acc aag gga cag 1093  
 Glu Ser Tyr Gln Glu Asp Ser Val His Ser Thr Ser Thr Lys Gly Gln  
 290 295 300  
 gaa aca tct aag agc aac aat att cct gct gcc att cac tcg cat aaa 1141  
 Glu Thr Ser Lys Ser Asn Asn Ile Pro Ala Ala Ile His Ser His Lys  
 305 310 315  
 agt tct aac gac tcc tgt ggc ttg cat tgc acg gaa cat att gct att 1189  
 Ser Ser Asn Asp Ser Cys Gly Leu His Cys Thr Glu His Ile Ala Ile  
 320 325 330  
 act agt aat aga gcc aca aga ttg gtg gcg gta acg aat gct gat cta 1237  
 Thr Ser Asn Arg Ala Thr Arg Leu Val Ala Val Thr Asn Ala Asp Leu  
 335 340 345  
 gag cag atg gca cag aac aga gat aat gct atg cag cgg tac aag gaa 1285  
 Glu Gln Met Ala Gln Asn Arg Asp Asn Ala Met Gln Arg Tyr Lys Glu  
 350 355 360 365  
 aag aag aaa acg cgg aga tat gat aag acc ata aga tat gaa acg agg 1333  
 Lys Lys Lys Thr Arg Arg Tyr Asp Lys Thr Ile Arg Tyr Glu Thr Arg  
 370 375 380  
 aag gcg aga gcc gag acc agg ttg cgt gtt aag ggc aga ttt gtg aaa 1381  
 Lys Ala Arg Ala Glu Thr Arg Leu Arg Val Lys Gly Arg Phe Val Lys  
 385 390 395  
 gct aca gat cct tag atgtctctcc acgttaggtt ttacatttga gatcctaagt 1436  
 Ala Thr Asp Pro  
 400  
 taggaacttt ttttgttttt tctactttca actaccttgt aaatgtaaat gatcgatctt 1496  
 cagctgcata atgtgtggcc agatttttgt aatttttacg ttttaaccttc taaaaaaaaa 1556  
 aa 1558

<210> 42  
 <211> 401  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 42

Met Gly Thr Ser Thr Thr Glu Ser Val Val Ala Cys Glu Phe Cys Gly  
 1 5 10 15  
 Glu Arg Thr Ala Val Leu Phe Cys Arg Ala Asp Thr Ala Lys Leu Cys  
 20 25 30  
 Leu Pro Cys Asp Gln His Val His Ser Ala Asn Leu Leu Ser Arg Lys  
 35 40 45  
 His Val Arg Ser Gln Ile Cys Asp Asn Cys Ser Lys Glu Pro Val Ser  
 50 55 60  
 Val Arg Cys Phe Thr Asp Asn Leu Val Leu Cys Gln Glu Cys Asp Trp  
 65 70 75 80  
 Asp Val His Gly Ser Cys Ser Ser Ser Ala Thr His Glu Arg Ser Ala  
 85 90 95  
 Val Glu Gly Phe Ser Gly Cys Pro Ser Val Leu Glu Leu Ala Ala Val  
 100 105 110  
 Trp Gly Ile Asp Leu Lys Gly Lys Lys Lys Glu Asp Asp Glu Asp Glu

MBI-20 Sequence Listing.ST25  
 120 125

115

Leu Thr Lys Asn Phe Gly Met Gly Leu Asp Ser Trp Gly Ser Gly Ser  
 130 135 140

Asn Ile Val Gln Glu Leu Ile Val Pro Tyr Asp Val Ser Cys Lys Lys  
 145 150 155 160

Gln Ser Phe Ser Phe Gly Arg Ser Lys Gln Val Val Phe Glu Gln Leu  
 165 170 175

Glu Leu Leu Lys Arg Gly Phe Val Glu Gly Glu Gly Glu Ile Met Val  
 180 185 190

Pro Glu Gly Ile Asn Gly Gly Gly Ser Ile Ser Gln Pro Ser Pro Thr  
 195 200 205

Thr Ser Phe Thr Ser Leu Leu Met Ser Gln Ser Leu Cys Gly Asn Gly  
 210 215 220

Met Gln Trp Asn Ala Thr Asn His Ser Thr Gly Gln Asn Thr Gln Ile  
 225 230 235 240

Trp Asp Phe Asn Leu Gly Gln Ser Arg Asn Pro Asp Glu Pro Ser Pro  
 245 250 255

Val Glu Thr Lys Gly Ser Thr Phe Thr Phe Asn Asn Val Thr His Leu  
 260 265 270

Lys Asn Asp Thr Arg Thr Thr Asn Met Asn Ala Phe Lys Glu Ser Tyr  
 275 280 285

Gln Glu Asp Ser Val His Ser Thr Ser Thr Lys Gly Gln Glu Thr Ser  
 290 295 300

Lys Ser Asn Asn Ile Pro Ala Ala Ile His Ser His Lys Ser Ser Asn  
 305 310 315 320

Asp Ser Cys Gly Leu His Cys Thr Glu His Ile Ala Ile Thr Ser Asn  
 325 330 335

Arg Ala Thr Arg Leu Val Ala Val Thr Asn Ala Asp Leu Glu Gln Met  
 340 345 350

Ala Gln Asn Arg Asp Asn Ala Met Gln Arg Tyr Lys Glu Lys Lys Lys  
 355 360 365

Thr Arg Arg Tyr Asp Lys Thr Ile Arg Tyr Glu Thr Arg Lys Ala Arg  
 370 375 380

Ala Glu Thr Arg Leu Arg Val Lys Gly Arg Phe Val Lys Ala Thr Asp  
 385 390 395 400

Pro

## MBI-20 Sequence Listing.ST25

<210> 43  
 <211> 844  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (89)..(658)  
 <223> G1387

<400> 43  
 tctctctccc actctcactt tctctcctat tcttagttcg tgtcagaaac acacagagaa 60  
 attaagaacc ctaatttaaa acagaaga atg gta cat tcg aag aag ttc cga 112  
 Met Val His Ser Lys Lys Phe Arg  
 1 5  
 ggt gtc cgc cag cgt cag tgg ggt tct tgg gtt tct gag att cgt cat 160  
 Gly Val Arg Gln Arg Gln Trp Gly Ser Trp Val Ser Glu Ile Arg His  
 10 15 20  
 cct ctc ttg aag aga aga gtg tgg cta gga aca ttc gac acg gcg gaa 208  
 Pro Leu Leu Lys Arg Arg Val Trp Leu Gly Thr Phe Asp Thr Ala Glu  
 25 30 35 40  
 aca gcg gct aga gcc tac gac caa gcc gcg gtt cta atg aac ggc cag 256  
 Thr Ala Ala Arg Ala Tyr Asp Gln Ala Val Leu Met Asn Gly Gln  
 45 50 55  
 agc gcg aag act aac ttc ccc gtc atc aaa tcg aac ggt tca aat tcc 304  
 Ser Ala Lys Thr Asn Phe Pro Val Ile Lys Ser Asn Gly Ser Asn Ser  
 60 65 70  
 ttg gag att aac tct gcg tta agg tct ccc aaa tca tta tcg gaa cta 352  
 Leu Glu Ile Asn Ser Ala Leu Arg Ser Pro Lys Ser Leu Ser Glu Leu  
 75 80 85  
 ttg aac gct aag cta agg aag aac tgt aaa gac cag aca ccg tat ctg 400  
 Leu Asn Ala Lys Leu Arg Lys Asn Cys Lys Asp Gln Thr Pro Tyr Leu  
 90 95 100  
 acg tgt ctc cgc ctc gac aac gac agc tca cac atc ggc gtc tgg cag 448  
 Thr Cys Leu Arg Leu Asp Asn Asp Ser Ser His Ile Gly Val Trp Gln  
 105 110 115 120  
 aaa cgc gcc ggg tca aaa acg agt cca aac tgg gtc aag ctt gtt gaa 496  
 Lys Arg Ala Gly Ser Lys Thr Ser Pro Asn Trp Val Lys Leu Val Glu  
 125 130 135  
 cta ggt gac aaa gtt aac gca cgt ccc ggt ggt gat att gag act aat 544  
 Leu Gly Asp Lys Val Asn Ala Arg Pro Gly Gly Asp Ile Glu Thr Asn  
 140 145 150  
 aag atg aag gta cga aac gaa gac gtt cag gaa gat gat caa atg gcg 592  
 Lys Met Lys Val Arg Asn Glu Asp Val Gln Glu Asp Asp Gln Met Ala  
 155 160 165  
 atg cag atg atc gag gag ttg ctt aac tgg acc tgt cct gga tct gga 640  
 Met Gln Met Ile Glu Glu Leu Leu Asn Trp Thr Cys Pro Gly Ser Gly  
 170 175 180  
 tcc att gca cag gtc taa aggagaatca ttgaattata tgatcaagat 688  
 Ser Ile Ala Gln Val  
 185  
 aataatatag ttgagggtta ataataatcg agggtaagta atttacgtgt agctaataat 748  
 taatataatt ttcgaacata tatatgaata tatgatagct ctagaaatga gtacgtatat 808  
 atacgtaaac atttttcctc aaatatagta tatgtg 844

<210> 44  
 <211> 189

## MBI-20 Sequence Listing.ST25

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 44

Met Val His Ser Lys Lys Phe Arg Gly Val Arg Gln Arg Gln Trp Gly  
 1 5 10 15

Ser Trp Val Ser Glu Ile Arg His Pro Leu Leu Lys Arg Arg Val Trp  
 20 25 30

Leu Gly Thr Phe Asp Thr Ala Glu Thr Ala Ala Arg Ala Tyr Asp Gln  
 35 40 45

Ala Ala Val Leu Met Asn Gly Gln Ser Ala Lys Thr Asn Phe Pro Val  
 50 55 60

Ile Lys Ser Asn Gly Ser Asn Ser Leu Glu Ile Asn Ser Ala Leu Arg  
 65 70 75 80

Ser Pro Lys Ser Leu Ser Glu Leu Leu Asn Ala Lys Leu Arg Lys Asn  
 85 90 95

Cys Lys Asp Gln Thr Pro Tyr Leu Thr Cys Leu Arg Leu Asp Asn Asp  
 100 105 110

Ser Ser His Ile Gly Val Trp Gln Lys Arg Ala Gly Ser Lys Thr Ser  
 115 120 125

Pro Asn Trp Val Lys Leu Val Glu Leu Gly Asp Lys Val Asn Ala Arg  
 130 135 140

Pro Gly Gly Asp Ile Glu Thr Asn Lys Met Lys Val Arg Asn Glu Asp  
 145 150 155 160

Val Gln Glu Asp Asp Gln Met Ala Met Gln Met Ile Glu Glu Leu Leu  
 165 170 175

Asn Trp Thr Cys Pro Gly Ser Gly Ser Ile Ala Gln Val  
 180 185

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/31344

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>												
IPC(7) : C12N 5/04, 5/10, 15/00, 15/09, 15/63, 15/70, 15/74, 15/82, 15/87; C07H 21/02, 21/04; A01H 1/00, 9/00, 11/00												
US CL : 435/320.1, 419, 468; 536/23.1; 800/ 278, 295												
<b>B. FIELDS SEARCHED</b>												
Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/320.1, 419, 468; 536/23.1; 800/ 278, 295												
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched												
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Continuation Sheet												
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>												
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.										
X	Database GenEmbl, Accession Number U28422, WANG et al., Arabidopsis thaliana Dna-binding protein CCAL (CCAL) mRNA, complete cds 14 January 1997.	4-6										
---		-----										
Y		1-3, 7-13, 25-27										
X	Database Geneseq., Accession Number V65382, THE REGENTS OF THE UNIVERSITY OF CALIFORNIA. 15 February 1999.	4-6										
---		-----										
Y		1-3, 7-13, 25-27										
X	Database PIR. Accession Number T02684, ROUNSLEY et al., DNA-binding protein CCA1 - Arabidopsis thaliana. 24 March 1999.	11										
---		-----										
Y		1-10, 12-13, 25-27										
X	Database Geneseq, Accession Number W79280, THE REGENTS OF THE UNIVERSITY OF CALIFORNIA. 15 February 1999.	11										
---		-----										
Y		1-10, 12-13, 25-27										
X	WO 98/48007 A1 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 28 October 1998, pages 26-33, pages 43-46 SEQ ID NO: 3.	1-13, 25-27										
P, Y	RIECHMANN et al. A genomic perspective on plant transcription factors. Current Opinion in Plant Biology. October 2000, Vol. 3, No. 5, pages 423-424, especially pages 427-428.	1-13, 25-27										
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.												
* Special categories of cited documents: <table border="0"> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E" earlier application or patent published on or after the international filing date</td> <td>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td>"&amp;" document member of the same patent family</td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	"P" document published prior to the international filing date but later than the priority date claimed	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention											
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone											
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art											
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family											
"P" document published prior to the international filing date but later than the priority date claimed												
Date of the actual completion of the international search 08 February 2001 (08.02.2001)		Date of mailing of the international search report <b>07 MAR 2001</b>										
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703)305-3230		Authorized officer <i>Cynthia Collins</i> Cynthia Collins Telephone No. (703) 308-0196										



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/31344

## C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	RIECHMANN et al. The AP2/EREBP family of plant transcription factors. Biological Chemistry, June 1998, Vol. 379, No. 6, pages 633-646.	1-13, 25-27
A	RIECHMANN et al. MADS domain proteins in plant development. Biological Chemistry, October 1997, Vol. 378, No. 10, pages 1079-1101.	1-13, 25-27
A	HEARD et al. Evolutionary diversity of symbiotically induced nodule MADS box genes: characterization of mntC5, a member of a novel subfamily. Molecular plant-microbe interactions: MPMI, July 1997, Vol. 10, No. 5, pages 665-676.	1-13, 25-27
A	HEARD et al. Symbiotic induction of a MADS-box gene during development of alfalfa root nodules. Proc. Natl. Acad. Sci. USA, 06 June 1995, Vol. 92, No. 12, pages 5273-5277.	1-13, 25-27

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/31344

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 14  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:  
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-13, 25-27 SEQ ID NOS: 1 and 2

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/31344

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING** This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Groups I-XXII, claim(s) 1-13 and 25-27, drawn to transgenic plants with modified biochemical characteristics, polynucleotides and vectors for producing said transgenic plants, and methods of making said transgenic plants. Applicant must elect one pair of sequences (one nucleotide sequence and its corresponding amino acid translation) per Group to be examined, *i.e.* SEQ ID NOS: 1 and 2 in Group I, SEQ ID NOS: 3 and 4 in Group II, SEQ ID NOS: 5 and 6 in Group III, etc.

Group XXIII, claim(s) 15-17, drawn to a method of identifying a factor that is modulated by or interacts with a polypeptide.

Group XXIV, claim(s) 18, drawn to a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest.

Group XXV, claim(s) 19 and 20, drawn to an integrated system, computer, or computer readable medium.

Group XXVI, claim(s) 21-23, drawn to a method of identifying a polynucleotide sequence.

The inventions listed as Groups I-XXVI do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The inventions listed as Groups I-XXVI do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Groups I-XXII are drawn to transgenic plants and methods of producing said plants with nucleic acid sequences. The methods of Groups I-XXII differ from each other in that they are directed to plant transformation methods and transgenic plants with structurally and functionally distinct nucleic acid sequences which encode structurally and functionally different amino acid sequences. In addition, Groups XXIII, XXIV, and XXVI are different methods from any of Groups I-XXII in that they have different method steps and different end products, and Group XXV requires a computer system. Thus, there is no single special technical feature which links the inventions of Groups I-XXVI under PCT Rule 13.2.

**Continuation of B. FIELDS SEARCHED Item 3:** STN (agricola, biosis, biotechno, biotechds, biotechabs, caba, caplus, embase, medline, uspatfull, wpids, pctfull, europatfull, japio) SEARCH TERMS: inventor names, plant transcription factor, fatty acid, chlorophyll, carotenoid; STIC: sequence search for SEQ ID NOS: 1 and 2

**THIS PAGE BLANK (USPTO)**